

(UNDATED)

HAID # 00015534

DATA EVALUATION RECORD

1. CHEMICAL: Metolachlor (108801)
2. FORMULATION: Technical
3. CITATION: Sachsse, K.; Ullman, L. (1974) Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA 24705. Project No. Siss 3516. Received Sep. 26, 1974 under 5G1553. (Unpublished report prepared by CIBA GEIGY Ltd., Basle, Switzerland; CDL:112840-N).
4. REASON FOR REVIEW: Generic Standard for Metolachlor
5. REVEIUED BY: H. T. Craven
Biologist
Efficacy and Ecological Effects Branch
Registration Division
6. DATA REVIEWED: 12/16/77
7. TEST TYPE: Cold Water Fish Acute 96 hr. (LC₅₀)
 - A. TEST ID: ES F 1
 - B. TEST SPECIES: Rainbow Trout (Salmo gairdneri)
 - C. TEST MATERIAL: Technical Metolachlor
 - D. REPORTED RESULTS: 96 hr (LC₅₀) = Approx. 2 ppm. In the report, a general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance, the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy.
 - E. CONCLUSIONS:

The aeration of a static bioassay may result in the volatilization of the toxicant from the medium, therefore it is impossible to assess the validity of the reported LC₅₀.

This study does not meet the requirement for a cold water fish acute LC₅₀.

Trout - INVALID

1

MATERIALS AND METHODS

A. Test Conditions: The study was described to only a limited extent as it relied on the statement:

"The procedure for testing followed that prescribed by the United States Federal Department of the Interior Fish and Wildlife Services: 'Procedures for evaluation of acute toxicity of Pesticides to fish and wildlife' 1964."

B. Statistical Analysis: The LC_{50} values were calculated by probit analysis according to Goulden A., 1960, Method of Statistical Analysis, John Wiley and Sons, third printing p. 404-408.

DISCUSSION/RESULTS

Reported Results: 96 hr (LC_{50}) = Approx. 2 ppm. In the report, a general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance, the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy.

REVIEWER'S EVALUATION

A. Test Procedure

Several deviations from the recommended protocol described in the proposed 1977 Guideline include: (1) Only four vs. a minimum of five dosage levels were tested; (2) Although acetone controls were run, no acetone free controls were established; (3) The test containers were aerated during the study. It is noted that the loading factor (1.9 g/liter) exceeded the recommended 1.0 g/liter thereby possibly necessitating aeration.

B. Statistical Analysis

The Environmental Safety Section did not attempt to validate the statistics portion of this study because aeration was performed, thereby negating any LC_{50} value.

C. Validation

1. Category: Invalid

2. Rationale: The aeration of a state bioassay may result in the volatilization of toxicant from the medium.

3. Repairability Rationale: The rainbow trout section of the study cannot be repaired even to supplemental.

CONCLUSIONS

The aeration of a static bioassay may result in the volatilization of the toxicant from the medium, therefore it is impossible to assess the validity of the reported LC₅₀; it is noted that the loading factor (1.9 g/liter) exceeded the recommended 1.0 g/liter thereby ^{possibly} necessitating aeration. This study does not meet the requirement for a cold water fish acute LC₅₀.

8. TEST TYPE: Warm Water Fish Acute 96 hr (LC₅₀)
- A. Test ID: ES G1
- B. Test Species: Crucian Carp (Carassius carassius), Guppy (Lebistes reticulatus), Bluegill (Lepomis macrochirus), Channel Catfish (Ictalurus ameriurus).
- C. Test Material: Technical Metolachlor
- D. Reported Results:

<u>Species</u>	<u>96 Hour LC₅₀ (ppm)</u>	<u>95% Confidence Limits</u>
Crucian Carp (<u>Carassius carassius</u>)	4.9	3.6 - 6.8
Channel Catfish (<u>Ictalurus ameriurus</u>)	4.9	3.6 - 6.8
Bluegill (<u>Lepomis macrochirus</u>)	15	*
Guppy (<u>Lebistes reticulatus</u>)	8.6	7.4 - 10.5

* No confidence limits were calculable

In the report, a general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy. These symptoms were seen at 2.1 ppm in carp and 6.5 ppm in guppy.

E. Results of Evaluation:

The LC₅₀ values reported for the guppy, crucian carp and channel catfish are scientifically sound. These LC₅₀ values indicate metalachlor is moderately toxic to fish.

warm water

?
bluegill
sunfish
fish and
A

(3)

MATERIALS AND METHODS

Test procedure and method of Statistical analysis was the same as previously cited in the rainbow trout portion of this study.

DISCUSSION/RESULTS

1. Guppy, Crucian carp and Channel Catfish

The 96 hour LC₅₀ values and 95% I.I. are respectively: Carp 4.9 (3.6-6.8) ppm, Channel Catfish 4.9 (3.6-6.8) and Guppy 8.6 (7.4-10.5) ppm. Where mortality occurred, those organisms displayed (after 4-6 hrs. exposure) hypersensitivity, loss of equilibrium and apathy. These symptoms were seen at 2.1 ppm in channel catfish and carp and 6.5 for guppy.

2. Bluegill

Four dosage levels were tested (1, 10, 21 and 49 ppm). No mortality occurred at the two lower levels; but the next two levels showed 75% and 100% mortality respectively. A 96 hour LC₅₀ of 15 ppm without confidence limits was reported.

REVIEWER'S EVALUATIONS

A. Test Procedure

Several deviations from the recommended protocol described in the proposed 1977 Guidelines include: (1) only four vs. a minimum of five dosage levels were tested; (2) although acetone controls were run, no acetone free controls were established, (3) in the case of the carp and the bluegill, the test temperature ($14^{\circ}\text{C} \pm 2^{\circ}\text{C}$) is below the recommended range ($19^{\circ}\text{C} - 26^{\circ}\text{C} \pm 1^{\circ}\text{C}$) for warm water species.

B. Statistical Analysis:

1. Carp and Guppy

Finney probit was performed on the guppy portion of the study; the resulting LC₅₀ 8.6 (see accompanying printout) is the same as the reported value. The same statistical analysis the Environmental Safety Section performed on the Channel Catfish study applies to the carp because the dosage levels and 96 hr % mortality are the same for both species (see xerox copy of catfish statistics).

2. Channel Catfish

The Environmental Safety Section performed a Finney probit analysis on the data (see accompanying printout). The determined LC₅₀ (4.8) compares favorably with the reported value of 4.9.

3. Bluegill

Environmental Safety Section did not perform a Finney probit analysis as the requirement for two partial mortality levels was not met. Instead, a linear regression line was constructed (see accompanying printout and graph). The LC₅₀ of 12.1 ppm cannot be confirmed by a test for Chi² and fails the Tab T test.

C. Validation

1. Carp and Guppy

- a. Category: Supplementary
- b. Rationale: Neither of these species are recommended test species. Furthermore, in the case of carp, the test temperature (14°C ± 2°C) is below the recommended range (19°C - 26°C) for warmwater species.
- c. Repairability rationale: This portion of the study cannot be upgraded to core.

2. Channel Catfish

- a. Category: ~~Supplementary~~ *core*
- b. Rationale: ~~The channel catfish Ictalurus punctatus is a recommended test species, however, the Environmental Safety Section was not able to determine if Ictalurus ameiurus is the same species.~~
- c. Repairability rationale: ~~This portion of the study can be upgraded to core providing Ictalures ameiurus is the same species as Ictalurus punctatus.~~

Personal communication with La-Deigz indicated that the species reported as ameiurus was actually punctatus.

3. Bluegill

- a. Category: Invalid
- b. Rationale: Conducting this portion of the study at too low a temperature (14°C ± 2°C) instead of (19°C - 26°C) prohibits this study from being classified as core. Secondly, the LC₅₀ value cannot be supported by statistical analysis.
- c. Repairability: This portion of the study can be upgraded to supplementary provided an appropriate statistical analysis is performed.

CONCLUSIONS

Carp and Guppy -

The studies are scientifically sound and indicate that metolachlor is moderately toxic to these species of fish. Neither the carp or the guppy are recommended test species therefore, while the studies augment the required data on a recommended warm water fish they do not serve as a substitute.

Channel catfish - This study indicates that metolachlor is moderately toxic to fish with an LC₅₀ of 4.9 ppm (95% C.L. 3.6 - 6.8 ppm). The study on channel catfish is sound and acceptable to meet the requirement for a warm water fish acute LC₅₀.

Bluegill Sunfish - The study on bluegill is not statistically sound. This is due to having too few partial mortality levels.

~~Table 6~~
 Catfish
 501553
 Done Dec 16
 Finney
 Probit
Dual

1.
 0.
 12.
 2.1
 1.
 12.
 6.5
 7.
 12.
 10.
 12.
 12.

Tot CH12 = 5.99 > 2.334

4.478	M
1.984	YINT
1.672	LW M
2.334	CHI2
4.838	LD50
3.633	LDCL
6.443	UPCL
2.503	LD10
1.552	LDCL
4.035	UPCL
9.353	LD90
6.480	LDCL
13.459	UPCL

Test Species Ictalurus gmelinus (?) PROBIT ANALYSIS WORK SHEET Chemical Metolachlor
 Source Aquarium Basel, Switzerland Date Tested 4 Jan. 1974
 o. Peribd. 96-hours Analysis by: Sachsse, K., + L. Ullmann; DVM, 1974
 (Name) (Title) (Date)

Concentration PPM	No. dead/ /No. tested	Observed % Mortality	Expected % Mortality	O-E	Contributions to Chi (Nomo. #1)
10	12/12	100			
6.5	7/12	58			
2.1	1/12	8			
1	0/12	0			

Total Fish Tested = _____
 Number of Doses (K) = _____
 Degrees of freedom (K-2) = _____

Chi² = Total Cont. \times $\frac{\text{Total fish}}{K}$ = 2.334
 to Chi
 Chi² (p=.05) for 2 deg of freedom = 5.99

DETERMINE FLC₅₀:

LC₈₄ _____
 LC₅₀ _____
 LC₁₆ _____

$S = \frac{LC_{84} - LC_{16}}{2}$
 $N' (\text{Fish used between } 16\% \text{ and } 84\% E) =$ _____
 $\sqrt{N'} =$ _____
 (Nomo. #2) = _____

$FLC_{50} = S^{2.77/\sqrt{N'}} = S$ _____

DETERMINE fS:

R (Largest/Smallest dose plotted) _____
 S (As determined above) _____
 A (Nomo. #3 using R and S) _____

$fS = A^{10(K-1)/K\sqrt{N'}} = A$ (Nomo. #2) = _____

DETERMINE FLC_y:

$(fS)^x = fS^{2.33 \text{ or } 1.30}$ (Table 3 and Nomo. #2) = _____

FLC_y (Nomo. #4 using $(fS)^x$ and FLC_{50}) = _____

RESULTS (LC_x and Confidence Limits at p = .05):


LC₁ = _____
 Lower Limit (LC₁/LC_y) _____
 Upper Limit (LC₁ X LC_y) _____

LC₅₀ = _____
 Lower Limit (LC₅₀/FLC₅₀) _____
 Upper Limit (LC₅₀ X FLC₅₀) _____

LC₉₉ = _____
 Lower Limit (LC₉₉/LC_y) _____
 Upper Limit (LC₉₉ X LC_y) _____

8

9



100m / 2.10m
5 6.5m 10 100m

Table 7
 Bluegill
 Dec 18, 77
 Dual
~~Tab~~ < Tab
 T.

1.
 C.
 10.
 C.
 21.
 75.
 45.
 100.

Cont
 do
 Funny
 Prohibit
 oo
 CH12

4.
 0.671
 0.310
 0.180

N
 R2
 S
 M

12.128
 0.042
 3475.075

LD50
 LDCL
 UPCL

7.120
 0.024
 2111.150

LD10
 LDCL
 UPCL

20.658
 0.060
 7155.062


LD90
 LDCL
 UPCL

95.
 2.
 4.303

XCON
 DF
 TVAL

0.671
 1.342
 1.
 0.671
 4.079027356
 4.079027356
 2.019664208

+
 C
 -
 =
 FX
 = +



Handwritten notes:
P1
P2
ppm

Table 8

~~guppy~~
 561553 1.
 Done Dec 19 12.
 Finney 6.5
 Profit 1.
 Dual 12.
 10.
 9.
 12.
 21.
 12.
 12.

Total
 Chi² = 5.997 0.00

11.001	M
-5.326	YINT
1.233	LW M
0.000	CHI ²
8.663	LD50
7.631	LDCL
9.879	UPCL
6.639	LD10
5.420	LDCL
8.133	UPCL
11.355	LD90
9.133	LDCL
14.118	UPCL

2/2/72

108801

VALIDATION SHEET

CRF #

PAGE 1 OF 1

FORMULATION:			IA	IB	T	FW	EC	R		
% a.i.	SC #	CHEMICAL NAME	Validator:				Date:			
Tech.		CGA-24705 (Metolachlor)	Labuda				02 December, 1977			
			Test Type:							
			Coldwater Fish Acute 96-hour LC ₅₀							
			Test ID.# ESG4							

CITATION: Sachesse, K., and L. Ullmann. 1974. Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy to Technical CGA-24705 Ciba-Geigy, Ltd.

VALIDATION CATEGORY: Invalid

RESULTS: 96-hour LC₅₀ for Salmo gairdneri was reported to be "approximately 2 ppm".

VALIDATION CATEGORY RATIONALE: Test aquaria were aerated throughout the treatment.

CATEGORY REPAIRABILITY/RATIONALE: No.

13

103301

VALIDATION SHEET

FORMULATION:			IA	IB	T	FW	EC	R		
% a.i.	SC #	CHEMICAL NAME	Validator:				Date:			
Tech.		CGA-24705 (Metolachlor)	Labuda				02 December, 1977			
			Test Type: MRID# 0003534							
			Warmwater Fish Acute 96-hour LC ₅₀							
			Test ID.#ESF5							

CITATION: Sachesse, K., and L. Ullmann. 1974. Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA-24705. Ciba-Geigy, Ltd.

VALIDATION CATEGORY:
Invalid ?

RESULTS: 96-hour LC₅₀ for Lepomis macrochirus was reported to be "approximately 15 ppm". (No 95% C. I. Was reported).

VALIDATION CATEGORY RATIONALE: Preferred test temperature for the Bluegill is not less than 18°C. This study was conducted at 14°C ± 2°.

CATEGORY REPAIRABILITY/RATIONALE: Yes to supplemental by supplying section with the appropriate statistical method used to derive the LC₅₀.

14

~~034001~~

MAIA# 00015534 108801

DATA EVALUATION RECORD

1. CHEMICAL: Metolachlor (108801)
2. FORMULATION: Technical
3. CITATION: Sachsse, K.; Ullman, L. (1974) Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA 24705. Project No. Siss 3516. Received Sep. 26, 1974 under 5G1553. (Unpublished report prepared by CIBA-GEIGY Ltd., Basle, Switzerland; CDL: 112840-N).
4. REASON FOR REVIEW: Generic Standard for Metolachlor
5. REVIEWED BY: H.T. Craven *H. T. Craven*
Biologist
Efficacy and Ecological Effects Branch
Registration Division
6. DATA REVIEWED: 12/16/77
7. TEST TYPE: Cold Water Fish Acute 96 hr. (LC₅₀)
 - A. TEST ID: ES F 1
 - B. TEST SPECIES: Rainbow Trout (Salmo gairdneri)
 - C. TEST MATERIAL: Technical Metolachlor
 - D. REPORTED RESULTS: 96 hr. (LC₅₀) = Approx. 2 ppm.
In the report, general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance, the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy.
 - E. COMMENTS: The aeration of a static bioassay may result in the volatilization of the toxicant from the medium; therefore, it is impossible to assess the validity of the reported LC₅₀. This study does not meet the requirement for a cold water fish acute LC₅₀.

15

MATERIALS AND METHODS

- A. Test Conditions: The study was described to only a limited extent as it relied on the statement:

"The procedure for testing followed that prescribed by the United States Federal Department of the Interior Fish and Wildlife Services: 'Procedures for evaluation of acute toxicity of Pesticides to fish and wildlife' 1964."

- B. Statistical Analysis: The LC_{50} values were calculated by probit analysis according to Goulden A., 1960, Method of Statistical Analysis, John Wiley and Sons, third printing pp. 404-408.

DISCUSSION/RESULTS

Reported Results: 96 hr. (LC_{50}) = Approx. 2 ppm. In the report, a general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance, the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy.

REVIEWER'S EVALUATION

- A. Test Procedure

Several deviations from the recommended protocol described in the proposed 1977 Guideline include: (1) Only four vs. a minimum of five dosage levels were tested; (2) Although acetone controls were run, no acetone-free controls were established; (3) The test containers were aerated during the study. It is noted that the loading factor (1.9 g/liter) exceeded the recommended 1.0 g/liter thereby possibly necessitating aeration.

- B. Statistical Analysis

The Environmental Safety Section did not attempt to validate the statistics portion of this study because aeration was performed, thereby negating any LC_{50} value.

- C. Validation

1. Category: Invalid
2. Rationale: The aeration of a state bioassay may result in the volatilization of the toxicant from the medium.

3. Repairability Rationale: The rainbow trout Section of the study cannot be repaired even to supplemental.

COMMENTS

The aeration of a static bioassay may result in the volatilization of the toxicant from the medium; therefore it is impossible to assess the validity of the reported LC₅₀; it is noted that the loading factor (1.9 g/liter) exceeded the recommended 1.0 g/liter thereby possibly necessitating aeration. This study does not meet the requirement for a cold water fish acute LC₅₀.

8. TEST TYPE: Warm Water Fish Acute 96 hr. (LC₅₀)

A. TEST ID: ES G1

B. TEST SPECIES: Crucian Carp (Carassius carassius), Guppy (Lebistes reticulatus), Bluegill (Lepomis macrochirus), Channel Catfish (Ictalurus ameriurus).

C. TEST MATERIAL: Technical Metolachlor

D. REPORTED RESULTS:

<u>SPECIES</u>	96 <u>LC₅₀</u> (ppm)	95% <u>CONFIDENCE LIMITS</u>
Crucian Carp (<u>Carassius carassius</u>)	4.9	3.6 - 6.8
Channel Catfish (<u>Ictalurus ameriurus</u>)	4.9	3.6 - 6.8
Bluegill (<u>Lepomis macrochirus</u>)	15	*
Guppy (<u>Lebistes reticulatus</u>)	8.6	7.4 - 10.5

*No confidence limits were calculable

In the report, a general comment was made with regard to all species tested: About 4 to 6 hours after adding the substance the fish in concentrations where mortality occurred showed hypersensitivity, loss of equilibrium and later apathy. These symptoms were seen at 2.1 ppm in channel catfish and carp and 6.5 ppm in guppy.

E. COMMENTS

The LC₅₀ values reported for the guppy, crucian carp, channel catfish and bluegill sunfish are scientifically sound. These LC₅₀ values indicate metolachlor is moderately toxic to warm water fish.

MATERIALS AND METHODS

Test procedure and method of statistical analysis was the same as previously cited in the rainbow trout portion of this study.

DISCUSSION/RESULTS

1. Guppy, Crucian Carp and Channel Catfish

The 96 hour LC₅₀ values and 95% 1.1 are respectively: carp 4.9 (3.6-6.8) ppm, channel catfish 4.9 (3.6-6.8) and guppy 8.6 (7.4-10.5) ppm. Where mortality occurred, those organisms displayed (after 4-6 hrs. exposure) hypersensitivity, loss of equilibrium and apathy. These symptoms were seen at 2.1 ppm in channel catfish and carp and 6.5 for guppy.

2. Bluegill

Four dosage levels were tested (1, 10, 21 and 49 ppm). No mortality occurred at the two lower levels; but the next two levels showed 75% and 100% mortality respectively. A 96 hour LC₅₀ of 15 ppm without confidence limits was reported.

REVIEWER'S EVALUATIONS

A. Test Procedure

Several deviations from the recommended protocol described in the proposed 1977 Guidelines include: (1) only four vs. a minimum of five dosage levels were tested, (2) although acetone controls were run, no acetone-free controls were established, (3) in the case of the carp and the bluegill, the test temperature ($14^{\circ}\text{C} \pm 2^{\circ}\text{C}$) is below the recommended range ($19^{\circ}\text{C} - 26^{\circ}\text{C} \pm 1^{\circ}\text{C}$) for warm water species.

B. Statistical Analysis:

1. Carp and Guppy

Finney probit was performed on the guppy portion of the study; the resulting LC₅₀ 8.6 (see accompanying printout) is the same as the reported value. The

same statistical analysis the Environmental safety Section performed on the channel catfish study applies to the carp because the dosage levels and 96 hr. % mortality are the same for both species (see Zerox copy of catfish statistics).

2. The Environmental Safety Section performed a Finney probit analysis on the data (see accompanying printout). The determined LC_{50} (4.8) compares favorably with the reported value of 4.9.
3. Bluegill

Environmental Safety Section did not perform a Finney probit analysis as the requirement for two partial mortality levels was not met. Instead, a linear regression line was constructed (see accompanying printout and graph). The LC_{50} of 12.1 ppm cannot be confirmed by a test for CHI^2 and fails the Tab T test.

C. Validation

1. Carp and Guppy

- a. Category: Supplementary
- b. Rationale: Neither of these species are recommended test species. Furthermore, in the case of carp, the test temperature ($14^{\circ}C + 2^{\circ}C$) is below the recommended range ($19^{\circ}C - 26^{\circ}C$) for warm water species.
- c. Repairability rationale: This portion of the study cannot be upgraded to core.

2. Channel Catfish

- a. Category: Core

Personal communication with Ciba-Geigy indicated that the species reported as ameriurus was actually punctatus.

3. Bluegill

- a. Category: Supplementary
- b. Rationale: Conducting this portion of the study at too low a temperature ($14^{\circ}C + 2^{\circ}C$) instead of

(19°C - 26°C) prohibits this study from being classified as core. Secondly, the LC₅₀ value has not been supported by statistical analysis.

- c. Repairability: Even if an appropriate statistical analysis is performed, this portion of the study cannot be upgraded to core because the temperature was too low.

COMMENTS

Carp and Guppy -

The studies are scientifically sound and indicate that metolachlor is moderately toxic to these species of fish. Neither the carp nor the guppy is recommended test species; therefore, while the studies augment the required data on a recommended warm water fish, they do not serve as a substitute.

Channel Catfish -

This study indicates that metolachlor is moderately toxic to fish with an LC₅₀ of 4.9 ppm (95% C.L. 3.6-6.8 ppm). The study on channel catfish is sound and acceptable to meet the requirement for a warm water fish acute LC₅₀.

Bluegill Sunfish -

The study on bluegill is not statistically sound. This is due to having too few partial mortality levels.



8
 guppy
 561553
 Done Dec 19
 Finney
 Profit
 Dual

1.
 C.
 12.
 6.5
 1.
 12.
 10.
 3.
 12.
 21.
 12.
 12.

Total
 Chi² = 5.997 0.00

11.001	M
-5.326	YINT
1.283	LW M
0.000	CHI ²
8.683	LD50
7.631	LDCL
9.879	UPCL
6.639	LD10
5.420	LDCL
8.133	UPCL
11.355	LD90
9.133	LDCL
14.118	UPCL

Table 6

Catfish (corp)

561553 1.
12.

Done Deal 12.

Funney 2.1

Probit 1.
12.

Dual 6.5

7.
12.

10.
12.
12.

Tab C#12 = 5.99 >

4.478
1.984
1.672
2.334

4.888
3.638
6.448

2.578
1.582
4.035

9.388
6.480
13.489

Table 6

Catfish

561553

Done Dec 16

Funney
Probit

Dual


1.
1.
12.
2.1
1.
12.
6.5
7.
12.
10.
12.
12.

Tot CH12 = 5.99 > 2.334

4.478	M
1.934	YINT
1.672	LW M
2.334	CHI2
4.838	LD50
3.633	LDCL
6.443	UPCL
2.503	LD10
1.552	LDCL
4.035	UPCL
9.353	LD90
6.480	LDCL
13.499	UPCL

33

Dual Tech (CGA-24705), Bluegill 96 hr. LC50




Mortality

M

95

Dual Tech (CGA-24705), Bluegel 46W. L-50



Molarity

ppm

a

b

10

21

108701

VALIDATION SHEET

CRF #

PAGE 1 OF 1

FORMULATION:			IA	IB	T	FW	EC	R		
% a.i.	SC #	CHEMICAL NAME	Validator:				Date:			
Tech.		CGA-24705 (Metolachlor)	Labuda				02 December, 1977			
			Test Type:							
			Warmwater Fish Acute 96-hour LC ₅₀ MID # 00015534							
			Test ID.# ESF6							

CITATION: Sachesse, K., and L. Ullmann. 1974. Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA 24705. Ciba Geigy, Ltd.

VALIDATION CATEGORY: Supplemental

RESULTS: 96-hour LC₅₀ for the guppy, Lebistes reticulatus, was reported to be 8.6 ppm (95% C.I. = 7.4-10.5 ppm).

VALIDATION CATEGORY RATIONALE: Lebistes reticulatus is not an acceptable test species for studies submitted in support of pesticide registration.

CATEGORY REPAIRABILITY/RATIONALE: No.

26

108801

VALIDATION SHEET

CRF #

PAGE 1 OF 1

FORMULATION:			IA	IB	T	FW	EC	R		
% a.i.	SC #	CHEMICAL NAME	Validator:				Date:			
Tech.		CGA-24705 (Metolachlor)	Labuda				02 December, 1977			
			Test Type:							
			Warmwater Fish Acute 96-hour LC ₅₀							
			Main # 00015534							
			Test ID.# ESF-4							

CITATION: Sachesse, K., and L. Ullman. 1974. Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA-24705. Ciba-Geigy, Ltd.

Validation Category:
Core

RESULTS: 96-hour LC₅₀ for Ictalurus punctatus (?) was reported to be 4.9 ppm (95% C.I. = 3.6-6.8 ppm).

VALIDATION CATEGORY RATIONALE: The species tested was Ictalurus punctatus rather than Ictalurus ameiurus as reported.

CATEGORY REPAIRABILITY/RATIONALE: NA

27

12830

FORMULATION:			IA	IB	T	FW	EC	R		
% a.i.	SC #	CHEMICAL NAME	Validator:				Date:			
Tech.		CGA-24705 (Metolachlor)	Labuda				02 December, 1977			
			Test Type:							
			Warmwater Fish Acute 96-hour LC ₅₀							
			MHID # 00015534							
			Test ID.# ESF3							

CITATION: Sachesse, K., and L. Ullmann. 1974. Acute Toxicity to Rainbow Trout, Crucian Carp, Channel Catfish, Bluegill, and Guppy of Technical CGA 24705. Ciba Geigy, Ltd.

VALIDATION CATEGORY: Supplemental

RESULTS: 96-hour LC₅₀ for Carassius carassius was reported to be 4.9 ppm (95% C.I. = 3.6 - 6.8 ppm).

VALIDATION CATEGORY RATIONALE: Crucian carp, considered a warmwater species, should not be tested at temperatures less than 18°C. In this study they were tested at 14°C ± 2°.

CATEGORY REPAIRABILITY/RATIONALE: No.