



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D.C., 20460

MEMORANDUM

DP Barcode: 365085

PC Code: 122804

February 24, 2010

SUBJECT: Ecological risk assessment evaluating Abamectin for the registration of a new end-use product (Agri-Mek®SC Miticide/Insecticide) for use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes

FROM: Ibrahim Abdel-Saheb, Environmental Scientist
Amy Blankinship, Chemist *Amy Blankinship*
Environmental Risk Branch 2
Environmental Fate and Effects Division (7507P)

2/24/10

TO: Venus Eagle, Risk Manager
Thomas Harris, Risk Manager Reviewer
Insecticide/Rodenticide Branch
Registration Division (7505P)

THRU: William Eckel, Acting Branch Chief *William P. Eckel*
Jean Holmes, RAPL *Jean Holmes*
Environmental Risk Branch 2
Environmental Fate and Effects Division (7507P)

2/24/10

The Environmental Fate and Effects Division (EFED) has completed the baseline ecological risk assessment for the proposed use of abamectin (PC Code 122804) as a new end-use product (Agri-Mek®SC Miticide/Insecticide) for use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes. Conclusions regarding the environmental fate and ecological effects and ecological risks associated with the proposed uses of the chemical can be found in the executive summary of the attached document.



ENVIRONMENTAL FATE AND EFFECTS SCIENCE CHAPTER

For The Proposed Registration of

**ABAMECTIN AS A NEW END-USE PRODUCT (AGRI-MEK® SC
MITICIDE/INSECTICIDE) FOR ALMONDS, WALNUTS, APPLES,
AVOCADOS, CELERIAC, CITRUS, COTTON, CUCURBIT,
FRUITING VEGETABLES, GRAPES, HERBS, HOPS, LEAFY
VEGETABLES, MINT, PEARS, PLUMS, PRUNES AND POTATOES**

USEPA PC Code: 122804

ERB 2 Team:

**Ibrahim Abdel-Saheb, Environmental Scientist
Amy Blankinship, Chemist
Environmental Fate and Effects Division (7507P)**

Branch Chief Approval:

**William Eckel, Acting Branch Chief
Environmental Risk Branch
Environmental Fate and Effects Division (7507P)**

Table of Contents

Table of Contents.....	3
List of Tables	5
List of Figures	6
1.0 Executive Summary	7
1.1 Nature of Chemical Stressor	7
1.2 Conclusions – Exposure Characterization	8
1.3 Conclusions – Effects Characterization	8
1.4 Potential Risks to Non-target Organisms.....	10
1.5 Key Uncertainties and Data Gaps	14
1.5.1 Key Uncertainties.....	14
1.5.2 Data Gaps	16
2.0 Problem Formulation	19
2.1 Nature of Regulatory Action.....	19
2.2 Stressor Source and Distribution	19
2.2.1 Nature of the Chemical Stressor	20
2.2.2 Proposed Label Crop Use Rates	21
2.2.3 Overview of Pesticide Use.....	24
2.2.4 Environmental Properties of Abamectin.....	25
2.3 Receptors	26
2.3.1 Aquatic and Terrestrial Effects	26
2.3.2 Incident Database Review.....	29
2.4 Ecosystems Potentially at Risk	30
2.5 Conceptual Model.....	31
2.5.1 Risk Hypothesis	31
2.5.2 Conceptual Diagram	31
2.6 Analysis Plan	32
2.6.1 Conclusions from Previous Risk Assessments	33
2.6.2 Preliminary Identification of Data Gaps	33
3.0 Analysis.....	35
3.1 Exposure Characterization.....	35
3.1.1 Measures of Aquatic Exposure	35
3.1.1.1 Aquatic Exposure Modeling	35
3.1.1.2 Aquatic Exposure Monitoring and Field Data	37
3.1.2 Measures of Terrestrial Exposure	38
3.2 Ecological Effects Characterization.....	41
3.2.1.1 Terrestrial Animals	41
3.2.1.2 Terrestrial Plants	43
3.2.2 Aquatic Effects Characterization	43
3.2.2.1 Aquatic Animals	43
3.2.2.2 Aquatic Plants	45
4.0 Risk Characterization.....	46
4.1 Risk Estimation – Integration of Exposure and Effects Data	46
4.1.1 Non-target Aquatic Animals and Plants	46
4.1.1.1 Non-target Aquatic Animals.....	46

4.1.1.2 Aquatic Plants	49
4.1.1.3 Non-target Terrestrial Animals	51
4.1.1.4 Non-target Terrestrial and Semi-Aquatic Plants	58
4.2 Risk Description	58
4.2.1 Risks to Aquatic Organisms.....	58
4.2.1.1 Fish and Aquatic Invertebrates	59
4.2.1.2 Aquatic Plants	60
4.2.2 Risks to Terrestrial Organisms.....	60
4.2.2.1 Terrestrial Animals	60
4.2.2.2 Terrestrial Plants	63
4.2.3 Federally Threatened and Endangered (Listed) Species Concerns.....	63
4.2.3.1 Taxonomic Groups potentially at Risk	63
4.2.3.2 Direct and Indirect Effects	64
4.3 Description of Assumptions, Limitations, Uncertainties and Data Gaps.	64
4.3.1 Related to Exposure for All Species	64
4.3.1.1 General Exposure Parameters	64
4.3.2 Related to Exposure Assessment	65
4.3.2.1 Related to Exposure for Aquatic Species.....	65
4.3.2.2 Related to Exposure for Terrestrial Species.....	65
4.3.3 Related to Effects Assessment	66
4.3.3.1 Age class and sensitivity of effects thresholds	66
4.3.3.2 Aquatic Studies Conducted Above Water Solubility.....	67
4.3.3.3 Lack of Effect Studies and Complete Review of Aquatic Plant Data	67
4.3.3.4 Uncertainty in LD50 for Mallards and NOAEC for Chronic Daphnia Study	67
4.3.3.5 Use of the Most Sensitive Species Tested	67
5.0 Literature Cited	68
 Appendix A. EIIS Incident Reports	 69
Appendix B. PRZM/EXAMS Output Files	71
Appendix C. T-REX Outputs.....	111
Appendix D. Summary of Toxicity Data for Abamectin.....	132
Appendix E. RQ Method and LOCs.....	139
Appendix F. Locates Output	140

List of Tables

Table 1. Proposed Application Rates for Crops Listed in Agri-Mek SC Label	22
Table 2. Physical and Chemical Properties of Abamectin.....	26
Table 3. Measures of Ecological Effects and Exposure for Abamectin	28
Table 4. Surface water exposure inputs for PRZM/EXAMS	36
Table 5. Tier II Surface Water 1-in10 Year EECs (ppb) of abamectin and its major soil degradate (a mixture of 8- α -hydroxy and a ring opened aldehyde derivative	37
Table 6. Avian Dose-Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Application of Abamectin.....	38
Table 7. Mammalian Dose-Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Application of Abamectin.....	39
Table 8. Dietary Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Exposure to Abamectin.....	40
Table 9. Summary of Most Sensitive Acute and Chronic Toxicity Data for Birds, Mammals and Terrestrial Invertebrates Exposed to Abamectin.....	42
Table 10. Summary of Selected Acute and Chronic Toxicity Data for Fish and Aquatic Invertebrates Exposed to Abamectin for use in Determining Risk.....	44
Table 11. Summary of Acute Toxicity Data for Aquatic Plants Exposed to Abamectin	45
Table 12. Acute Risk Quotients for Fish and Aquatic Invertebrates from Abamectin Applied to Various Crops	47
Table 13. Chronic Risk Quotients for Fish and Aquatic Invertebrates from Abamectin Applied to Various Crops	48
Table 14. Risk quotients for Aquatic Plants Exposed to Foliar Applications of Abamectin	50
Table 15. Upper bound acute dose-based RQ values for birds for foliar application of abamectin	51
Table 16. Upper Bound Acute Avian Dietary-based RQ values from Foliar Application of Abamectin to Celeriac, Cucurbit, Fruiting and Leafy Vegetables, Herbs and Potato	53
Table 17. Comparison of the Dietary EECs from Foliar Application of Abamectin to the Chronic Avian NOAEC	53
Table 18. Upper bound Mammalian Acute Dose-based RQ values for Foliar Application of Abamectin.....	54
Table 19. Upper bound Mammalian Chronic Dose-based RQ values for Foliar Application of Abamectin.....	55

Table 20. Upper bound Chronic Dietary-based RQ Values for Mammals for Foliar Application of Abamectin.....	56
Table 21. Comparisons of Small and Large Insect EECs from Foliar Application of Abamectin to the Extrapolated Acute Contact Honeybee Concentration.....	58

List of Figures

Figure 1. Chemical Structure of Abamectin	20
Figure 2. Estimated use of abamectin in 2002 (USGS)	25
Figure 3. Conceptual diagram for assessment of risks from abamectin use on various crops.....	32

1.0 Executive Summary

Syngenta Crop Protection, Inc. is seeking a registration of abamectin (PC Code 122804) and its new end-use product Agri-Mek®SC Miticide/Insecticide) for almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes for control of mites, thrips, leafminers, leafhoppers, psyllids, potato beetles, skeletonizer, and pinworms.

The new end-use product may be applied by ground application and also aerially for some crops, except for in New York. The maximum single application rate ranges from 0.014 to 0.023 lb ai/A, and the maximum seasonal application rate ranges from 0.038 to 0.056 lb ai/A.

1.1 Nature of Chemical Stressor

Abamectin (also known as avermectin) is a mixture of macrocyclic lactones and is a fermentation product of the soil fungus, *Streptomyces avermitilis*. The active ingredient abamectin is a mixture of avermectins containing at least 80% avermectin B_{1a} (5-0-demethyl avermectin A_{1a}) and at most 20% avermectin B_{1b} (5-0-demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A_{1a}). A major soil degrade is a mixture of 8- α -hydroxy and a ring opened aldehyde derivative.

Abamectin is a miticide/insecticide registered for use on almonds, walnuts, apples, avocados, citrus fruits, cucurbits, grapes, fruiting vegetables and other crops. It is also registered as a nematicide for use as a seed treatment for corn and cotton (Avicta™ 500FS) and as a seed treatment for cucurbits and tomatoes (Avicta™ 400 FS). It is also registered as a treatment for as an indoor and outdoor bait for insects such as ants and roaches, waterbugs, and palmetto bugs.

The proposed registration action is for a new formulation, Agri-Mek® SC Miticide/Insecticide, an aqueous suspension concentrate that contains abamectin (avermectin B_{1a} & B_{1b}), for use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes for control of mites, thrips, leafminers, leafhoppers, psyllids, potato beetles, skeletonizer, and pinworms. According to the registrant, abamectin is not dissolved in the new end-use product, rather the particles of abamectin are suspended in water. Also, depending on the crop, Agri-Mek SC must be mixed with a horticulture oil (not a dormant oil), non-ionic surfactant, spreading and penetrating surfactant, cucurbit approved adjuvant or organosilicone adjuvant (potatoes only) to avoid the possibility of exceeded established crop tolerances. Agri-Mek SC may be applied by ground application and by aerial application for avocados, cucurbit, fruiting and leafy vegetables, mint, and potatoes and for control of citrus leafminer in citrus fruit (not in California). Aerial application is not approved in New York. Agri-Mek SC can not be applied within 25 ft for ground application or 150 ft for aerial application of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries or commercial fish farm

ponds. In addition, the label restricts cultivation within 25 ft of the aquatic area to allow growth of a vegetative filter strip. The label states not to apply Agri-Mek SC or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Abamectin acts as a chlorine channel agonist in invertebrates (Fritz, *et al.*, 1979, Mellin *et al.*, 1983 and Arena *et al.*, 1991 in Sherma and Cairns, 1993), and may function as a gamma-aminobutyric acid (GABAergic) agonist (Kass *et al.*, 1980, 1984 in Sherma and Cairns, 1993). It acts by stimulating the release of gamma-aminobutyric acid, an inhibitory neurotransmitter, thus causing paralysis (Tomlin, 1994). The difference in toxicity between invertebrates and mammals may be partially due to different distribution of the GABAergic neurons (Turner and Schaeffer, 1989 in Sherma and Cairns, 1993).

1.2 Conclusions – Exposure Characterization

The new proposed use of abamectin may result in drift onto plants, soil, or water adjacent to a treated field. Any abamectin on the soil surface or in clear, shallow surface water should undergo rapid photodegradation (half-life <1 day). However, photodegradation is not likely to be significant where abamectin is incorporated or under canopy. In addition, in most surface waters, suspended sediments and lack of mixing would decrease the rate of photodegradation. In natural waters, abamectin residues are expected to be associated with the sediment, reducing aqueous concentrations. Abamectin slowly biodegrades in soil (90% upper confidence bound of mean half-life = 80.6 days). Abamectin is stable to hydrolytic degradation. Due to its low vapor pressure (1.5×10^{-9} Torr); it is not likely that volatilization will be a transport process for abamectin.

Laboratory studies indicate that abamectin has moderate to low mobility ($K_{ads} = 9.7$ to 160 mg kg^{-1}); adsorption was correlated with soil organic matter content. Submitted field dissipation studies are unacceptable; therefore, EFED can not determine if the behavior of abamectin in the laboratory is demonstrated in the field. Based upon the laboratory data, ground water effects are expected to be minimal.

1.3 Conclusions – Effects Characterization

Aquatic invertebrates are the aquatic species most sensitive to abamectin. It is very highly acutely toxic to aquatic invertebrates, with a 48-h EC_{50} value of $0.34 \text{ } \mu\text{g ai/L}$ in the freshwater waterflea, *Daphnia magna*, and a 96-h LC_{50} of $0.020 \text{ } \mu\text{g ai/L}$ (20 parts-per-trillion) in the estuarine/marine mysid shrimp, *Americamysis bahia*. Abamectin is highly toxic to the embryo/larval stages of mollusks with a 48-h EC_{50} of $430 \text{ } \mu\text{g ai/L}$ (total form (both dissolved and undissolved abamectin)) in the Eastern Oyster. This value is above the water solubility of abamectin (7.8 ppb in distilled water; $<1 \text{ ppb}$ in tap water) without the presence of a vehicle such as acetone to increase its water solubility. The life-cycle toxicity test with the *Daphnia magna* resulted in a reproductive NOAEC of $0.030 \text{ } \mu\text{g ai/L}$ which was the lowest concentration tested, but the adults in the two lowest treatment groups were observed to be pale and smaller compared to the controls (MRID 00153570) and growth was not measured in the study. Therefore, the reproductive NOAEC appears

to underestimate the true no-effect concentration for Daphnia from chronic exposure to abamectin, as the NOAEC appears to be lower than 0.030 µg ai/L (30 parts-per-trillion). An acute to chronic ratio using the mysid shrimp toxicity data was used to calculate a chronic no-effect concentration for the daphnia and is 0.006 µg ai/L (6 parts-per-trillion). The NOAEC value for the life-cycle toxicity test with the mysid shrimp (*Americamysis bahia*) was previously reported as 0.0035 µg ai/L based on reproduction when compared to the solvent control, but is 0.00035 µg ai/L (0.35 parts-per-trillion) based on reproduction when compared to the negative control as there was a difference between the negative and solvent control for reproduction. Current EFED policy is to compare treatment groups to the negative control, therefore, the NOAEC value of 0.00035 µg ai/L was used in the assessment.

Abamectin is also very highly toxic to freshwater fish with an acute 96-h LC₅₀ value of 3.2 µg ai/L (total form) for rainbow trout (*Oncorhynchus mykiss*), a 96-h LC₅₀ value of 9.6 µg ai/L (total form) for bluegill sunfish (*Lepomis macrochirus*) and an acute 96-h LC₅₀ value of 15.0 µg ai/L (total form) for sheepshead minnow (*Cyprinodon variegatus*). These values are above the water solubility of abamectin (7.8 µg/L in distilled water; <1 µg/L in tap water) without the presence of a solvent such as acetone or DMF to increase its water solubility. The freshwater fish chronic toxicity NOAEC is 0.52 µg ai/L, based on an early life stage study in rainbow trout based on growth (wet weight). There is no chronic estuarine-marine fish study for abamectin, therefore an acute to chronic ratio was used to determine a no-effect concentration. The extrapolated estuarine/marine fish chronic toxicity NOAEC is 2.41 µg/L.

In birds, the acute oral LD₅₀ for bobwhite quail (*Colinus virginianus*) is >2,000 mg/kg-bw (practically nontoxic), whereas the acute oral LD₅₀ for mallard ducks (*Anas platyrhynchos*) is 85 mg/kg-bw (highly toxic). The dietary LC₅₀ values obtained in short-term toxicity tests in bobwhite quail and mallard ducks are >3,102 and 383 mg ai/kg-diet, respectively. There were no statistically significant effects on growth, survival or reproduction in the mallard duck reproduction study at the highest concentration tested, 12 mg ai/kg-diet, therefore, the no observed adverse effect concentration (NOAEC) is at least 12 mg ai/kg-diet for the mallard duck chronic reproduction study (MRID 40318601). During the pilot study for the mallard duck reproduction study, the average number of eggs laid was markedly less in the 64 mg ai/kg treatment group.

In laboratory rats, abamectin has an acute toxicity LD₅₀ value of 13.6 mg/kg-bw, when dosed using a sesame oil vehicle, and a 2-generation reproductive NOAEC value of 0.12 mg/kg-bw based on increased retinal folds, increased dead pups at birth, decreased viability and lactation indices, and decreased pup body weight. Based on two rat carcinogenicity studies abamectin is not a carcinogen and based on five mutagenicity and a cytogenetics test abamectin is not a mutagen.

Abamectin is highly toxic to the Honey Bee with an acute dermal LD₅₀ of 0.41 µg/bee. A foliar residue study on citrus, demonstrates that residues are toxic for approximately 48 hours.

Abamectin has been tested for phytotoxicity in only two aquatic plant species. The growth or biomass inhibition nominal concentration IC_{50} values obtained in these studies are >100 mg ai/L (total form) and 3.9 mg ai/L (total form) for the green alga *Selenastrum capricornutum* and the vascular aquatic plant *Lemna gibba*, respectively. These values are above the water solubility of abamectin (7.8 µg/L distilled water; <1 µg/L in tap water) without the presence of a solvent such as acetone or DMF to increase its water solubility. These studies were conducted using acetone, which is a potential photosensitizer and abamectin is subject to photolysis. Bioavailable dissolved concentrations are unknown, as test solutions were not analyzed.

Abamectin does not bioaccumulate significantly in fish or in mammals. Terrestrial plant toxicity data was not available.

1.4 Potential Risks to Non-target Organisms

Non-Listed Organisms

Acute risk is not expected for non-listed fish, birds or mammals from application of the new end-use abamectin product. Acute risk is expected for non-listed freshwater and estuarine/marine invertebrates. The potential for adverse risk also exists for terrestrial invertebrates and plants from use of abamectin. The RQ values did not exceed the non-listed LOC for aquatic plants, but data for only two of the five recommended species were submitted, and there are technical issues with the submitted data.

Listed Organisms

There is a potential for adverse risk to listed freshwater fish, freshwater and estuarine/marine invertebrates, birds, reptiles, amphibians, and mammals. The potential for adverse risk also exists for terrestrial invertebrates and plants from use of abamectin. The RQ values did not exceed the listed LOC for aquatic plants, but data for only two of the five recommended species were submitted, and there are technical issues with the submitted data.

Aquatic Organisms

Acute

Non-Listed Species

- There were no acute non-listed LOC exceedances for either freshwater or estuarine/marine fish.
- RQ values did exceed the acute non-listed LOC of 0.5 for estuarine/marine invertebrates for all crops (RQs 1.45-32.6), and for freshwater aquatic invertebrates from abamectin use on apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes and potatoes.

Listed Species

- There were no acute listed LOC exceedances for estuarine/marine fish for any crop scenario.
- The acute freshwater and estuarine/marine invertebrate RQ values exceed the Agency's acute listed LOC of 0.05 for all crop scenarios (RQs 0.085-1.91 for freshwater and 1.45-32.6 for estuarine/marine).
- The acute freshwater fish RQ values exceed the Agency's acute listed LOC for abamectin application to apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes, and potatoes (RQs 0.087-0.203).
- RQ values for aquatic plants did not exceed the listed or non-listed LOC. However, data for only two of the five required species was available for review. In addition, submitted studies were conducted as nominal concentrations with the use of a potential photosensitizing solvent; therefore, risk may be underestimated.

Chronic

- The chronic RQ values for fish did not exceed the LOC for any crop scenario.
- Chronic freshwater and estuarine/marine invertebrate RQ's exceed the chronic LOC (1.0) for all crop scenarios (RQs 3.83-94.0 for freshwater and 65.7-1611 for estuarine/marine).
- The life-cycle toxicity test with the *Daphnia magna* resulted in a reproductive NOAEC of 0.030 µg ai/L which was the lowest concentration tested, but the adults in the two lowest treatment groups were observed to be pale and smaller compared to the controls (MRID 00153570) and length and weight were not measured. Therefore, the reproductive NOAEC appears to underestimate the true no-effect concentration for *Daphnia* from chronic exposure to abamectin, as the NOAEC appears to be lower than 0.030 µg ai/L which may be underestimating risk. Therefore, an extrapolated NOAEC value, based on an acute to chronic ratio using the mysid shrimp toxicity data

Terrestrial Organisms

Acute

Non-Listed Species

- The acute dose-based and dietary-based RQ values for birds and dose-based RQ values for mammals did not exceed the non-listed LOC of 0.5 for any crop scenario. However, regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity.

Listed Species

- The avian acute dietary-based RQ values did not exceed the acute listed LOC of 0.1 for any crop scenario.
- The acute avian dose-based RQ values exceed the acute listed LOC for small birds feeding on small and tall grass, broadleaf plants and small insects for all

crop scenarios, except for tall grasses for cotton, grapes and hops, and the LOC was exceeded for medium birds consuming short grasses for all crops except cotton, grapes and hops (RQs 0.10-0.30).

- Since birds are surrogates for reptiles and land-phase amphibians, the potential for direct effects may exist for these taxa as well.
- Acute dose-based RQ values exceeded the LOC for small and medium mammals consuming short and tall grass, broadleaf plants and small insects for all crops, except for medium mammals consuming tall grass for cotton, grapes and hops (RQs 0.11-0.38).
- The acute dose-based listed LOC was also exceeded for large mammals feeding on short grasses for all crop scenarios and broadleaf plants and small insects for abamectin application to celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes (RQs 0.10-0.17).
- There are no data regarding the toxicity of abamectin to terrestrial plants, therefore RQ values were not calculated. Due to the lack of data, and reported incidences for almonds and grapes indicated possible plant injury due to abamectin, risk can not be precluded.
- Abamectin is highly toxic to the honeybee. Calculated EECs were greater than the honeybee acute contact toxicity value, and there was an incidence reported that indicated honeybee mortality from abamectin use on avocados. Therefore, the proposed abamectin use is expected to be toxic to terrestrial invertebrates and beneficial insects.

Chronic

- Chronic dose-based and dietary-based RQ values exceed the Agency's chronic LOC (1.0) for mammals feeding on short and tall grass, broadleaf plants and small insects (RQs 5.74-42.64 for dose-based and 1.45-4.92 for dietary based).
- Chronic dose-based RQ values also exceeded the LOC for small and medium mammals consuming fruits, pods or large insects for all crops and for large mammals from abamectin use on celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes (RQs 1.22-2.67).
- No chronic dietary-based RQ values exceeded the chronic LOC for mammals consuming fruits, pods, seeds, or large insects or for seeds on a chronic dose basis.
- Chronic risk to birds is not expected as the calculated EECs are lower than the highest concentration tested in the mallard reproduction study.

Table 1. Potential Risks to Nonlisted and Listed Species Associated with Direct or Indirect Effects from the Proposed Application of abamectin for use on Crops

Taxonomic Group	Effects Endpoint	Direct Effects		Indirect Effects to Listed Species	
		Non-listed	Listed	Potential	Indirect Effects Due to Direct Effect to: ²
Dicot terrestrial plants	Survival and Growth	Data not available, risk can not be precluded		Yes	Mammals and birds

Taxonomic Group	Effects Endpoint	Direct Effects		Indirect Effects to Listed Species	
		Non-listed	Listed	Potential	Indirect Effects Due to Direct Effect to: ²
Monocot terrestrial plants	Survival and Growth	Data not available, risk can not be precluded		Yes	Mammals and birds
Mammals	Acute oral dose: mortality Chronic: growth and survival of offspring	Acute: No Chronic: Yes	Acute: Yes Chronic: Yes	Yes	Terrestrial plants, terrestrial insects
Birds ²	Acute oral dose: mortality Chronic: growth & reproduction	Acute: No Chronic: No	Acute: Yes Chronic: No	Yes	Terrestrial plants, terrestrial insects
Terrestrial invertebrates	Acute contact: mortality	Acute: Yes	Acute: Yes	Yes	Terrestrial plants, birds
Freshwater Fish	Acute dose: mortality Chronic: growth & survival	Acute: No Chronic: No	Acute: Yes Chronic: No	Yes	Freshwater invertebrates, terrestrial plants
Freshwater Invertebrates	Acute dose: mortality Chronic: growth & reproduction	Acute: Yes Chronic: Yes	Acute: Yes Chronic: Yes	Yes	Freshwater fish, birds, terrestrial plants
Estuarine-marine fish	Acute dose: mortality Chronic: growth & survival	Acute: No Chronic: No ³	Acute: No Chronic: No ³	Yes	Estuarine/marine invertebrates, terrestrial plants
Estuarine-marine Invertebrates	Acute dose: mortality Chronic: survival	Acute: Yes Chronic: Yes	Acute: Yes Chronic: Yes	Yes	Birds, terrestrial plants
Aquatic Vascular Plants	Growth ⁴	Acute: No Chronic: No	Acute: No Chronic: No	Yes	Birds, terrestrial plants
Aquatic Non-Vascular Plants	Growth ⁴	Acute: No Chronic: No	Acute: No Chronic: No	Yes	Freshwater & estuarine/marine invertebrates, terrestrial plants

¹ Direct effects to species may result in indirect effects to other species by changing availability of prey, habitat, and other factors important to survival and reproduction.

² Since birds are surrogates for reptiles and land-phase amphibians, potential risk to these groups may occur due to direct effects to birds.

³ RQ value calculated using ACR using freshwater fish chronic NOAEC and LC50 value.

⁴ Studies conducted as nominal concentrations with the use of a potential photosensitizer solvent, so risk may be underestimated.

1.5 Key Uncertainties and Data Gaps

1.5.1 Key Uncertainties

- A number of the acute toxicity tests were conducted as nominal concentration static studies and were above the reported solubility limit for abamectin (7.8 µg/L in distilled water (MRID 47051904) and <1.0 µg/L in tap water (D235416)). In addition, the studies were conducted with acetone which is a potential photosensitizer, and abamectin has an aqueous photolysis half-life of 12 hours. Therefore, the use of acetone may have contributed to possible degradation of abamectin in the test solutions especially in the aquatic plant studies. Overall, the dissolved bioavailable concentration of abamectin in these toxicity tests is unknown. Risk quotients calculated from these values may underestimate risks. The acute static daphnia study was also conducted using nominal concentrations. The current OPPTS 850.1075 (acute fish) guideline states that there must be evidence that test concentrations remained at least 80 percent of the nominal concentrations throughout the test or that mean measured concentrations are an accurate representation of exposure levels. The OPPTS 850.1010 (acute daphnia) guideline indicates that the concentration of the test chemical in the chambers should be measured as often as is feasible during the test. Also, the 850.5400 (algal toxicity) indicates the concentration of test chemical in the test containers is to be determined at the beginning and end of the definitive test by standard analytical methods which have been validated prior to the test. Since test solutions were not measured in the acute fish, daphnia, oyster and aquatic plant studies, the actual bioavailable abamectin concentration these organisms were exposed to is not known which increases the uncertainty of the toxicity values. Therefore, it is recommended that the acute fish (rainbow trout, bluegill, and sheepshead minnow), daphnia, oyster, and aquatic plant (duckweed and green algae) studies be repeated under current guidance which would involve the measurement of dissolved (bioavailable) abamectin in the test solutions.
- The registrant submitted *Daphnia magna* chronic life-cycle study with abamectin did not measure growth in the parental generation at the end of the study (total length or dry weight) (MRID 00153570). The current no-effect concentration is the lowest concentration tested based on survival. The study does indicate that at test termination, the surviving adult daphnia in the two lowest treatment groups were pale and appeared smaller compared to the controls which may suggest that the actual no-effect concentration is less than the lowest treatment group tested. Risk quotients calculated from the current no adverse effect concentration may underestimate risk. The current OPPTS 850.1300 guideline states that growth for each surviving adult should be determined (total body length or dry weight, or both). It is preferred that both measures be taken. Therefore, it is recommended that the chronic *Daphnia magna* life-cycle study be repeated. Since the actual no-effect concentration may be less than the lowest treatment group tested, the acute and chronic toxicity values from the mysid shrimp studies were used to calculate an acute to chronic ration for the daphnia. This ratio was used to determine a

chronic no-effect concentration for the daphnia and was used to calculate risk quotients which may be overestimating or underestimating risk.

- In the registrant submitted mysid chronic toxicity study with abamectin, reproduction in the solvent control was statistically significant compared to the negative control which may indicate that the solvent may have interfered with the integrity of the test. In the study, reproduction in the treatment groups was compared to the solvent control, but current EFED policy is to compare to the negative control regardless if the controls are statistically different. Comparison of reproduction resulted in a lower no-effect concentration than previously reported, and the lower no-effect concentration was used in this assessment.
- An early life-cycle study for estuarine-marine fish with abamectin was not available. Therefore, the acute and chronic toxicity values from the rainbow trout studies were used to develop an acute to chronic ratio for the sheepshead minnow. This ratio was used to determine a chronic no-effect concentration for the sheepshead and was used to calculate risk quotients which may overestimate or underestimate risk.
- Regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity.
- The label states that for a number of crops (celeriac, cucurbit, fruiting vegetable, leafy vegetable, mint and potatoes (for potato psyllid) not to make more than two sequential applications of Agri-Mek SC or any other foliar applied abamectin containing product, but the maximum seasonal amount allowed for these crops is greater than two applications at the maximum single application rate. The application interval for these crops is 7 days, and the label does not state how long to wait between the second sequential application and subsequent applications. Also, the maximum amount allowed per season for these crops, except mint, is slightly less (0.001 lb ai/A) than the amount applied using three applications at the maximum single application rate. Since the label does not specifically state the interval between the second sequential application and subsequent applications, three applications at seven day intervals using the maximum seasonal rate divided by three was modeled for environmental exposure.
- For application to herbs, the label states not to make more than two applications of Agri-Mek SC per single cutting (harvest), but the maximum amount allowed per cropping season is greater than two applications at the maximum single application rate but slightly less than three applications at the maximum single application rate. Therefore, environmental exposure concentrations were modeled in the same manner as discussed above.
- For application to almonds, walnuts, apples, avocados, citrus, pears, plums and prunes, the label states that for the maximum amount per season, not to apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar

applied abamectin containing product in a growing season. Based on the density of the formulation, 8.5 fl oz/A calculates to 0.04648 lb ai/A, therefore, it is not known if the reported 0.047 lb ai/A is a rounding issue or if another abamectin product can be applied at 0.001 lb ai/A. In addition, the single maximum application rate reported is 0.023 lb ai/A, and two applications would be 0.046 lb ai/A. For this assessment, abamectin was modeled at 0.0235 lb ai/A (0.047 divided by two applications). Abamectin was also modeled at 0.023 lb ai/A which resulted in the same LOC exceedances as the 0.0235 lb ai/A application.

- The maximum seasonal application rate for cotton, potatoes (for Colorado potato beetle) and grapes on the label is reported as 0.038 lb ai/A, but the label also indicates not to apply more than 6.75 fl oz/A of Agri-Mek SC per season which calculates to 0.0369 (0.037) lb ai/A. The maximum single application rate for cotton, potatoes and grapes is 0.019 lb ai/A, and if applied twice per season, the maximum seasonal application rate would be 0.038 lb ai/A. Therefore, a maximum seasonal application rate of 0.038 lb ai/A was used for determining environmental exposure concentrations.
- EFED believes that the inclusion of the suggested buffer zone of (25 ft, for ground application; and 150 ft for aerial application) will not appreciably change the outcome of the risk assessment.

1.5.2 Data Gaps

This assessment is potentially underestimating risk to both terrestrial and aquatic organisms from exposure to abamectin. This potential underestimation is due to a lack of available toxicity data as well as technical issues with the data submitted for some species. Therefore, the following toxicity studies are requested.

- OPPTS 850.1400- Early Life-Stage Toxicity Test. There are no chronic toxicity data available for the Agency to assess chronic risk of abamectin to estuarine/marine fish.
- OPPTS 850.4225 – Seedling Emergence, Tier II and OPPTS 850.4250 – Vegetative Vigor, Tier II. Seedling emergence and vegetative vigor toxicity data are not available for terrestrial plants.
- OPPTS 850.2300 - Avian reproduction Study. A reproduction study with bobwhite quail is not available.

- OPPTS 850.2100 - Acute Oral Toxicity with a Passerine Bird. An acute oral toxicity study with a passerine bird is not available. No species recommended at this point. Protocol should be submitted prior to test initiation.
- Whole Sediment Toxicity Test: Chronic Invertebrates Freshwater and Marine. Based on the physiochemical properties, abamectin may sorb to organic materials in sediment and may be toxic to organisms that dwell in and ingest sediment as abamectin is very highly toxic to other aquatic invertebrates. Since abamectin is a foliar application, spray drift to both freshwater and estuarine-marine environments is possible. The concentration of abamectin in water from spray drift from ground or aerial application is greater than the acute EC₅₀ value for the estuarine/marine mysid shrimp. 40 CFR Part 158.630 requires a chronic freshwater sediment study if the half-life is greater than or equal to 10 days and any of the following conditions exist: i. $K_d \geq 50$, ii. the $\log K_{ow} \geq 3$, or iii. the $K_{oc} \geq 1000$. Abamectin meets these criteria. A protocol should be submitted to the Agency for review prior to testing.
- OPPTS 850.1075 – Fish Acute Toxicity Test, freshwater and marine; 850.1010- Aquatic Invertebrate Acute Toxicity test with Daphnia; 850.1025 or 1055 – Oyster Acute Toxicity Test (shell deposition) or Bivalve Acute Toxicity Test (embryo-larvae). The registrant submitted test were conducted as static tests that were conducted above the reported water solubility, conducted using a potential photosensitizing solvent and test concentrations were not measured. As a result, the actual test concentrations (dissolved bioavailable abamectin) are not known which may be underestimating risk. Therefore, a new acute toxicity study for a coldwater and warmwater freshwater fish, estuarine-marine fish and *Daphnia magna* is requested. An oyster shell deposition or a bivalve embryo-larvae toxicity study is also requested.
- OPPTS 850.1300 – Daphnia Chronic Toxicity Test. The registrant submitted chronic daphnia toxicity test did not measure growth for the surviving adults at test termination. The study indicates that the surviving daphnia in the two lowest concentrations tested were pale and smaller than the control. Measurement of growth is required under the current guidance. Therefore, a new study is requested.
- OPPTS 850.5400 – Algal Toxicity and 850.4400 Aquatic Plant Toxicity Test using Lemna spp. There are limited studies (data on two of the five species available (duckweed and a green alga study)) addressing the toxicity of abamectin to aquatic plants; the studies conducted with duckweed and green algae were conducted above solubility, with a potential photosensitizing solvent, and test concentrations were not measured. Abamectin toxicity studies with a marine diatom, freshwater diatom and blue-green algae are requested as well as new studies for the green algae and duckweed.

- Submitted field dissipation studies are unacceptable; therefore, the behavior of abamectin in the field as compared to the laboratory cannot be demonstrated. In most cases we would expect dissipation in the field to be greater than that predicted by laboratory studies due to pesticide transport.

2.0 Problem Formulation

2.1 Nature of Regulatory Action

This ecological risk assessment evaluates the use of the insecticide/miticide abamectin (PC 122804) as a new aqueous suspension concentrate end-use product, Agri-Mek®SC Miticide/Insecticide. The assessment is based on the proposed label use of the new end-use product on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes for control of mites, thrips, leafminers, leafhoppers, psyllids, potato beetles, skeletonizers, and pinworms. The proposed label is listed as a restricted use pesticide and may only be used by certified applicators or persons under their direct supervision, and only for the uses covered by the certified applicator's certificate.

The new end-use product may be applied by ground application and also aerially for some crops, except for in New York. The maximum single application rate ranges from 0.014 to 0.023 lb ai/A, and the maximum seasonal application rate ranges from 0.038 to 0.056 lb ai/A.

2.2 Stressor Source and Distribution

Abamectin (Figure 1) is a fermentation product of the soil fungus, *Streptomyces avermitilis*. Abamectin has been registered since the 1980s as an insecticide/miticide to be used for crop protection in numerous fruit and vegetable crops. Some of the active registrations are under trade names Avid®, Zephyr®, Agri-Mek®, Abamectin, Epi-Mek®, Abacide™, and Abasol™. It is also registered as a treatment for Fire Ants (Varsity™); turf, lawns, and other non-crop areas such as parks and golf courses, and in and around residential, commercial (food and non-food establishments) and industrial structures¹ for Fire Ants, Pharaoh Ants and related ants (Ascend and TC); as an indoor and outdoor ant² and insect pest³ crack and crevice treatment for residential, commercial (food and non-food establishments) and industrial structures⁴, and transportation equipment⁵ (AVERT® and TC); as an indoor and outdoor bait for ants and pests⁶ (Raid Baits); and for use as a cotton and corn seed treatment (Avicta™ 500 F) and as a seed

¹ Warehouses, hotels, food storage areas, meat packing plants, motels, schools, supermarkets, hospitals and nursing homes

² Includes but not limited to acrobat, allegheny, argentine, bigheaded, carpenter, soybeans field, crazy, fire, ghost, harvester, little black, odorous house, pavement, pharaoh, and pyramid

³ Booklice, carpet beetles, cockroaches, crickets, drugstore beetles, earwigs, flour beetles, grain weevils, pillbugs, and sowbugs

⁴ Apartments, campgrounds, garages, food storage areas, homes, hospitals and nursing homes (non-occupied patient areas), hotels, meat packing and food processing plants, motels, resorts, restaurants and other food handling establishments, schools, supermarkets, utilities, warehouses, and other commercial and industrial buildings

⁵ Buses, boats, ships, trains, trucks, planes

⁶ Roaches, waterbugs, palmetto bugs

treatment for cucurbits and tomatoes (Avicta™ 400 FS) to control nematodes. It is also used as a veterinary antihelminthic (destroys or causes expulsion of parasitic intestinal worms).




Figure 1. Chemical Structure of Abamectin

The proposed registration action is for a new formulation, Agri-Mek® SC Miticide/Insecticide, an aqueous suspension concentrate that contains abamectin (avermectin B1a & B1b), for use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes for control of mites, thrips, leafminers, leafhoppers, psyllids, potato beetles, skeletonizers, and pinworms. According to the registrant, abamectin is not dissolved in the new end-use product, rather the particles of abamectin are suspended in water. Also, depending on the crop, Agri-Mek SC must be mixed with a horticulture oil (not a dormant oil), non-ionic surfactant, spreading and penetrating surfactant, cucurbit approved adjuvant or organosilicone adjuvant (potatoes only) to avoid the possibility of exceeding established crop tolerances. Agri-Mek SC may be applied by ground application and by aerial application for avocados, cucurbit, fruiting and leafy vegetables, mint, and potatoes and for control of citrus leafminer in citrus fruit (not in California). Aerial application is not approved in New York. Agri-Mek SC can not be applied within 25 ft for ground application or 150 ft for aerial application of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries or commercial fish farm ponds. In addition, the label restricts cultivation within 25 ft of the aquatic area to allow growth of a vegetative filter strip. The label states not to apply Agri-Mek SC or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

2.2.1 Nature of the Chemical Stressor

The active ingredient abamectin is a mixture of avermectins containing at least 80% avermectin B_{1a} (5-0-demethyl avermectin A_{1a}) and up to 20% avermectin B_{1b} (5-0-demethyl-25-de(1-methylpropyl)-25-(1-methylethyl) avermectin A_{1a}).

Abamectin acts as a chlorine channel agonist in invertebrates (Fritz, *et al.*, 1979, Mellin *et al.*, 1983 and Arena *et al.*, 1991 in Sherma and Cairns, 1993), and may function as a gamma-aminobutyric acid (GABAergic) agonist (Kass *et al.*, 1980, 1984 in Sherma and Cairns, 1993). It acts by stimulating the release of gamma-aminobutyric acid, an

inhibitory neurotransmitter, thus causing paralysis (Tomlin, 1994). The difference in toxicity between invertebrates and mammals may be partially due to different distribution of the GABAergic neurons (Turner and Schaeffer, 1989 in Sherma and Cairns, 1993).

2.2.2 Proposed Label Crop Use Rates

The new end-use product may be applied by ground application and also aerially for some crops, except for in New York. The maximum single application rate ranges from 0.014 to 0.023 lb ai/A, and the maximum seasonal application rate ranges from 0.038 to 0.056 lb ai/A. Agri-Mek SC must be mixed with a horticulture oil (not a dormant oil), non-ionic surfactant, spreading and penetrating surfactant, cucurbit approved adjuvant or organosilicone adjuvant (potatoes only) to avoid the possibility of exceeding established crop tolerances.

There are a few uncertainties regarding the label language in terms of maximum seasonal application rate and application intervals:

- The label states that for a number of crops (celeriac, cucurbit, fruiting vegetable, leafy vegetable, mint and potatoes (for potato psyllid) not to make more than two sequential applications of Agri-Mek SC or any other foliar applied abamectin containing product, but the maximum seasonal amount allowed for these crops is greater than two applications at the maximum single application rate. The application interval for these crops is 7 days, and the label does not state how long to wait between the second sequential application and subsequent applications. Also, the maximum amount allowed per season for these crops, except mint, is slightly less (0.001 lb ai/A) than the amount applied using three applications at the maximum single application rate. Since the label does not specifically state the interval between the second sequential application and subsequent applications, three applications at seven day intervals using the maximum seasonal rate divided by three was modeled for environmental exposure.
- For application to herbs, the label states not to make more than two applications of Agri-Mek SC per single cutting (harvest), but the maximum amount allowed per cropping season is greater than two applications at the maximum single application rate but slightly less than three applications at the maximum single application rate. Therefore, environmental exposure concentrations were modeled in the same manner as discussed above.
- For application to almonds, walnuts, apples, avocados, citrus, pears, plums and prunes, the label states that for the maximum amount per season, not to apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar applied abamectin containing product in a growing season. Based on the density of the formulation, 8.5 fl oz/A calculates to 0.04648 lb ai/A, therefore, it is not known if the reported 0.047 lb ai/A is a rounding issue or if another abamectin product can be applied at 0.001 lb ai/A. In addition, the single maximum application rate reported is 0.023 lb ai/A, and two applications would be 0.046 lb ai/A. For this assessment, abamectin was modeled at 0.0235 lb ai/A (0.047

divided by two applications). Abamectin was also modeled at 0.023 lb ai/A which resulted in the same LOC exceedances as the 0.0235 lb ai/A application.

- The maximum seasonal application rate for cotton, potatoes (for Colorado potato beetle) and grapes on the label is reported as 0.038 lb ai/A, but the label also indicates not to apply more than 6.75 fl oz/A of Agri-Mek SC per season which calculates to 0.0369 (0.037) lb ai/A. The maximum single application rate for cotton, potatoes and grapes is 0.019 lb ai/A, and if applied twice per season, the maximum seasonal application rate would be 0.038 lb ai/A. Therefore, a maximum seasonal application rate of 0.038 lb ai/A was used for determining environmental exposure concentrations.

The maximum single and seasonal application rate, application rate interval and method of application for each of the crops listed in the Agri-Mek SC label is presented below in Table 1.

Table 1. Proposed Application Rates for Crops Listed in Agri-Mek SC Label

Crop	Max. Application rate lbs. a.i./A	No. Applications	Max. Seasonal Application rate lb ai/A ¹	Application Interval (days)	Application Method ³
Almonds & Walnuts	0.023	2	0.047 (Max seasonal app of 8.5 fl oz/A)	21	Ground
Apples	0.023	2	0.047 (Max seasonal app of 8.5 fl oz/A)	21	Ground
Avocados	0.023	2	0.047 (Max seasonal app of 8.5 fl oz/A)	30	Ground & Aerial
Celeriac	0.019	* ²	0.056 (Max seasonal app of 10.25 fl oz/A)	7	Ground
Citrus (calamondin, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin, sour orange, sweet orange, pummelo, Satsuma mandarin)	0.023	3	0.047 (Max seasonal app of 8.5 fl oz/A)	30	Ground; Aerial (citrus leafminer, not in CA)
Cotton	0.019	Not Reported	0.038 (reported on	21	Ground & Aerial

			label) (Max seasonal app of 6.75 fl oz/A)		
Cucurbits (Chayote, chinese waxgourd, citron melon, cucumber, gherkin, edible gourd, momordica spp, muskmelon, pumpkin, summer and winter squash, watermelon)	0.019	* ²	0.056 (Max seasonal app of 10.25 fl oz/A)	7	Ground & Aerial
Fruiting Vegetables (eggplant, groundcherry, pepino, peppers, tomatillo, tomato)	0.019	* ²	0.056 (Max seasonal app of 10.25 fl oz/A)	7	Ground & Aerial
Grapes	0.019	Not Reported	0.038 (reported on label) (Max seasonal app of 6.75 fl oz/A)	21	Ground & Aerial
Herb Crop Subgroup (except chives)	0.019	2 (per single cutting)	0.056 (Max seasonal app of 10.25 fl oz/A)	7	Ground
Hops (not in CA)	0.019	2	0.038	21	Ground
Leafy vegetables (amaranth, arugula, cardoon, celery, celtuce, chervil, chinese celery, chrysanthemum edible, corn salad, cress, dandelion, dock, endive, fennel, lettuce, New Zealand spinach, orach, parsley, purslane, radicchio, rhubarb, spinach, Swiss chard)	0.019	* ²	0.056 (Max seasonal app of 10.25 fl oz/A)	7	Ground & Aerial
Mint	0.014	* ² only 3 per season	0.042 (Max seasonal app of 7.75 fl oz/A)	7	Ground & Aerial
Pears (including Oriental pear trees)	0.023	2	0.047 (Max seasonal app of 8.5 fl oz/A)	21	Ground
Plums and Prunes	0.023	2	0.047 (Max seasonal app of 8.5 fl oz/A)	21	Ground
Potatoes	0.019	* ²	0.038;	7	Ground &

			0.056 (Max seasonal app of 6.75 fl oz/A for CO beetle, 10.25 fl oz/A for leafminer		Aerial
--	--	--	--	--	--------

¹ One gallon of Agri-Mek SC contains 0.7 lb abamectin

² * = label states not to make more than 2 sequential applications of Agri-Mek SC or any other foliar applied abamectin containing product.

³ Aerial application not approved in New York.

2.2.3 Overview of Pesticide Use

The current proposed registration is for the new end-use product Agri-Flex for use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes.

Abamectin is currently registered for use on these crops, except cotton, using the emulsifiable concentrate end-use product Agri-Mek 0.15 EC (EPA Reg. # 100-898) which was first registered in 1989.

Data are available which display the estimated annual use of abamectin (Figure 2).




Figure 2 Estimated use of abamectin in 2002 (USGS)

2.2.4 Environmental Properties of Abamectin

A summary of the physical and chemical properties are listed in Table 2. Based on fate properties and application methods, it is expected that abamectin will persist long enough to be available for transport to non-target environments. However, strong sorption to soil is expected to significantly reduce concentrations in the water column and in runoff water.

The results from reviewed studies indicate that abamectin should undergo rapid photodegradation (half-life <1 day) on the soil surface and in clear, shallow surface water. Photodegradation is not likely to be significant where abamectin is incorporated or under canopy. In addition, in most surface waters, suspended sediments and lack of mixing would decrease the rate of photodegradation. In natural waters, abamectin residues are expected to be associated with the sediment, reducing aqueous concentrations. Abamectin slowly biodegrades in soil (90% upper confidence bound of mean half-life = 80.6 days). Abamectin is stable to hydrolytic degradation. Due to its low vapor pressure (1.5×10^{-9} Torr); it is not likely that volatilization will be a transport process for abamectin.

Abamectin is nearly insoluble in water (7.8 ppb at pH 9 in distilled water; <1 ppb in tap water (D235416)). Laboratory studies indicate that abamectin has moderate to

low mobility ($K_{ads} = 9.7$ to 160 mg kg^{-1}); adsorption was correlated with soil organic matter content. Submitted field dissipation studies are unacceptable; therefore, EFED can not determine if the behavior of abamectin in the laboratory is demonstrated in the field. Based upon the laboratory data, ground water effects are expected to be minimal. Surface water contamination could occur from runoff events that occur soon after application.

Table 2 Physical and Chemical Properties of Abamectin

	Value	Source
Common name	Abamectin, Avermectin	
Pesticide type	Insecticide, Acaricide, Nematicide	
CAS number	71751-41-2	
Empirical formula	$C_{48}H_{72}O_{14} + C_{47}H_{70}O_{14}$	
Molecular mass (g/mol)	866.6	
Vapor pressure (Torr)	1.5×10^{-9}	MRID# 47051904
Henry's Law Constant ($\text{atm}\cdot\text{m}^3/\text{mol}$)	2.6×10^{-8}	MRID# 47051904
Solubility in water ($\mu\text{g/L}$)	7.8 (distilled water); <1 (tap water)	MRID# 47051904; D235416
Log Kow	4.4 at 25°C (pH aqueous phase 7.2)	MRID# 47051904
pK _a	No pK _a in aqueous solutions in the range of 1-12	MRID# 47051904

2.3 Receptors

2.3.1 Aquatic and Terrestrial Effects

In order for a chemical to pose an ecological risk, it must reach ecological receptors in biologically significant concentrations. An exposure pathway is the means by which a contaminant moves in the environment from a source to an ecological receptor. For an ecological exposure pathway to be complete, it must have a source, a release mechanism, an environmental transport medium, a point of exposure for ecological receptors, and a feasible route of exposure. In addition, the potential mechanisms of transformation (i.e., which degradates may form in the environment, in which media, and how much) must be known, especially for a chemical whose metabolites/degradates are of greater toxicological concern. The assessment of ecological exposure pathways, therefore, includes an examination of the source and potential migration pathways for constituents,

and the determination of potential exposure routes (e.g., ingestion, inhalation, and dermal absorption).

Ecological receptors that may potentially be exposed to abamectin on-field or off-field from spray drift or run-off include terrestrial wildlife (i.e., invertebrates, mammals, birds, and reptiles), and terrestrial and semi-aquatic plants. In addition to terrestrial ecological receptors, aquatic receptors (e.g., freshwater and estuarine/marine fish and invertebrates, amphibians, aquatic plants) may also be exposed to potential migration of pesticides from the site of application to various watersheds and other aquatic environments via runoff and drift.

Consistent with the process described in the Overview Document (EPA, 2004), this risk assessment uses a surrogate species approach in its evaluation of the proposed new end-use product of abamectin. Data generated from surrogate test species, which are intended to be representative of broad taxonomic groups, are used to extrapolate to potential effects on a variety of species (receptors) included under these taxonomic groupings.

A summary of the assessment and measurement endpoints selected to characterize potential ecological risks associated with exposure to abamectin is provided in Table 3.

Table 3. Measures of Ecological Effects and Exposure for Abamectin

Assessment Endpoint		Selected Surrogate Species and Measure of Ecological Effect ¹	Measures of Exposure
Birds ²	Acute Survival	Mallard (<i>Anas platyrhynchos</i>) acute oral LD ₅₀ (most sensitive avian acute oral LD ₅₀)	Maximum residues on dietary food items (dietary Estimated Environmental Concentrations (EEC))
	Survival, reproduction and growth	Mallard (<i>A. platyrhynchos</i>) Reproduction NOAEC (no statistical effects noted at highest concentration tested) (single study available)	
Mammals	Acute Survival	Lab Rat (<i>Rattus norvegicus</i>) acute oral LD ₅₀ (most sensitive acute oral study)	Maximum residues on dietary food items (dietary Estimated Environmental Concentrations (EEC))
	Survival, reproduction and growth	Lab Rat (<i>Rattus norvegicus</i>) 2-generation reproductive NOAEC (based on increased retinal folds, increased dead pups at birth, decreased viability and lactation indices, decreased pup body weight) (most sensitive reproduction NOAEC)	
Terrestrial Invertebrates	Acute Survival	Honey Bee (<i>Apis mellifera</i>) acute contact study (single study available)	µg abamectin /Animal
Freshwater fish ³	Acute Survival	Rainbow Trout (<i>Oncorhynchus mykiss</i>) 96-h LC ₅₀ (most sensitive 96-h fish acute LC ₅₀)	Surface water peak concentration (EEC) ⁴
	Survival, reproduction ⁵ and growth	Rainbow Trout (<i>Oncorhynchus mykiss</i>) Early Life-Stage NOAEC (wet weight) (single freshwater vertebrate early life-cycle study available)	Surface water 60-d average concentration (EEC) ⁴
Freshwater invertebrates	Acute Survival	Water Flea (<i>Daphnia magna</i>) 48-h EC ₅₀ (most sensitive freshwater invertebrate 48-h EC ₅₀ or 96-h LC ₅₀)	Surface water peak concentration (EEC) ⁴
	Survival, reproduction ⁵ and growth	Water Flea (<i>D. magna</i>) Life cycle NOAEC (reproduction) (single freshwater invertebrate life cycle study available)	Surface water 21-d average concentration (EEC) ⁴
Estuarine/ marine fish	Acute Survival	Sheepshead Minnow (<i>Cyprinodon variegatus</i>) 96-h LC ₅₀ (single estuarine/marine fish acute 96-h LC ₅₀ available)	Surface water peak concentration (EEC) ⁴
	Survival, reproduction ⁵ and growth	No data available; used acute to chronic ratio using rainbow trout data	Surface water 60-d average concentration (EEC) ⁴
Estuarine/ marine invertebrates	Acute Survival	Mysid Shrimp (<i>Americamysis bahia</i>) 96-h EC ₅₀ (most sensitive estuarine/marine acute 96-h LC ₅₀ or IC ₅₀ available)	Surface water peak concentration (EEC) ⁴
	Survival, reproduction and growth	Mysid Shrimp (<i>A. bahia</i>) Life cycle NOAEC (reproduction) (single estuarine/marine life cycle study available)	Surface water 21-d average concentration (EEC) ⁴
Aquatic plants	Biomass and Growth Rate	Vascular plant Duckweed (<i>Lemna gibba</i>) 14 day IC ₅₀ (single vascular aquatic plant study available)	Surface water peak concentration (EEC) ⁴
	Biomass and Growth Rate	Nonvascular plant Freshwater alga (<i>Selenastrum capricornutum</i>) 9 day EC ₅₀ (single alga study available)	

LD₅₀ = Lethal dose to 50% of the exposed test population; NOAEC = No observed adverse effect concentration; NOAEL = No observed adverse effect level; LC₅₀ = Lethal concentration to 50% of the exposed test population; EC₅₀ = Effect concentration to 50% of the test population; IC₅₀ = inhibition concentration resulting in a 50% inhibition in the test population response (e.g., growth rate, biomass)

¹Values listed in this table represent the most sensitive study result within the taxonomic group and for the measurement endpoint identified to evaluate attribute changes.

²Birds represent surrogates for amphibians (terrestrial-phase) and reptiles.

³Freshwater fish are used here as surrogates for amphibians (aquatic-phase).

⁴One in 10-year return frequency.

⁵Sensitive early-life stage embryo development, hatching success, and survival and growth of the young are used as a measure of reproduction success.

2.3.2 Incident Database Review

A review of the Ecological Incident Information System (EIIS, version 2.1), which is maintained by the Agency's Office of Pesticide Programs, and the Avian Monitoring Information System (AIMS), which is maintained by the American Bird Conservancy, indicates a total of seven reported ecological incidents associated with the use of abamectin, which are summarized below.

All of the abamectin reported incidents occurred between 1998 and 2003. Two of the abamectin incidents involved aquatic animals, one involved terrestrial animals, and four involved plants. The certainty categories on the likelihood that the use of abamectin caused the seven incidents ranged from possible (4 incidents) to probable (3 incidents). The incidents were considered registered uses at the time of the incident. The one incident with the bees was from the Section 18 use of abamectin for avocados in California. One of the incidents involved an additional chemical besides abamectin. Six reported incidents for abamectin involved uses that are currently Section 3 registrations (almonds, grapes, citrus, and fire ant control). In the report for the incident with the Section 18 for avocados in California, it was reported that the abamectin was not being applied in accordance with the label. The reported incidents associated with the six currently registered uses had certainty categories of possible and probable. A summary of the reported incidences are listed in Appendix A.

According to Office of Pesticides Program Ecological Incident Information System (EIIS), seven incident reports exist in EFED's database. Three of the incidents occurred in June 1998 from direct application of Agri-Mek to almonds in California (I007644-001, 002, 003). The type of injury to the almonds was not reported, but was reported to occur to all applied (34-106 acres). Agri-Mek was applied directly to 34 acres of grapes in June 2000 in California, with all 34 acres affected (I10837-019). They type of injury was not reported, and in the report, the inspector stated "Questionable" in regards to the question "Application within Label". There were two incidents involving freshwater fish. The first incident occurred in April 2000 in Texas, where 100 catfish died two days after 1/8 of a pound of both the pesticide Ascend Fire Ant Stopper (abamectin) and Award (fenoxycarb) were applied to areas around the pond (I010221-001) was reported. The next day one to one and a half inches of rain fell. No other fish species in the pond were observed to be affected. The second fish incident occurred in June 2003 in Florida where a citrus grove was treated with Agri-Mek less than 25 feet from a lake in the morning and

then it rained in the afternoon (I014237-001). One week after the application, the reported indicated that "tons" of dead small bait fish were observed around the pond edges. The last incident involved the spraying of abamectin (Agri-Mek) to avocados in California (I008611-001) under a Section 18 label in April 1999. Southern California beekeepers indicated that the abamectin was aerially sprayed during the daytime during full bloom which was not consistent with favored County instructions. They indicated that it is common to keep bee colonies in avocado fields. The report indicated that 100 colonies were affected.

In addition to the incidents recorded in EIIS and AIMS, additional incidents have been reported to the Agency in aggregated incident reports. Pesticide registrants report certain types of incidents to the Agency as aggregate counts of incidents occurring per product per quarter. Ecological incidents reported in aggregate reports include those categorized as 'minor fish and wildlife' (W-B), 'minor plant' (P-B), and 'other non-target' (ONT) incidents. 'Other non-target' incidents include reports of adverse effects to insects and other terrestrial invertebrates. For abamectin, registrants have reported one minor fish and wildlife incident and four other non-target incidents. Unless additional information on this aggregated incident becomes available, it will be assumed to be representative of registered uses of abamectin in the risk assessment.

A major incident report for abamectin has not been received by the Agency since 2003 and twelve incidents total (7 major and 5 minor) have been reported to the Agency. Incident reports for non-target organisms typically provide information only on mortality events and plant damage. Sublethal effects in organisms such as abnormal behavior, reduced growth and/or impaired reproduction are rarely reported, except for phytotoxic effects in terrestrial plants. EPA's changes in the registrant reporting requirements for incidents in 1998 may account for a reduced number of reported incidents. Registrants are now only required to submit detailed information on 'major' fish, wildlife, and plant incidents. Minor fish, wildlife, and plant incidents, as well as all other non-target incidents, are generally reported aggregately and are not included in EIIS. In addition, there have been changes in state monitoring efforts due to a lack of resources.

2.4 Ecosystems Potentially at Risk

The ecosystems at risk are often extensive in scope, and as a result it may not be possible to identify specific ecosystems during the development of a baseline risk assessment. However, in general terms, terrestrial ecosystems potentially at risk could include the treated field and areas immediately adjacent to the treated field that may receive drift or runoff. Areas adjacent to the treated field could include cultivated fields, fencerows and hedgerows, meadows, fallow fields or grasslands, woodlands, riparian habitats and other uncultivated areas.

Aquatic ecosystems potentially at risk include water bodies adjacent to, or down stream from, the treated field and might include impounded bodies such as ponds, lakes and reservoirs, or flowing waterways such as streams or rivers. For uses in coastal areas, aquatic habitat also includes marine ecosystems, including estuaries.

2.5 Conceptual Model

A conceptual model provides a written description and visual representation of the predicted relationships between abamectin, potential routes of exposure, and the predicted effects for the assessment endpoint. A conceptual model consists of two major components: risk hypothesis and a conceptual diagram (EPA, 1998).

2.5.1 Risk Hypothesis

For abamectin, the following ecological risk hypothesis is being employed for this baseline risk assessment:

Abamectin, when used in accordance with the label, results in potential adverse effects upon the survival, growth, and reproduction of non-target terrestrial and aquatic organisms.

2.5.2 Conceptual Diagram

For a pesticide to pose an ecological risk, it must reach ecological receptors in toxicologically significant concentrations. An exposure pathway is the means by which the pesticide moves in the environment from a source to reach the receptor. For an ecological exposure pathway to be complete, it must have a source, a release mechanism, an environmental transport medium, a point of exposure for ecological receptors, and a feasible route of exposure. The assessment of ecological exposure pathways, therefore, includes an examination of the source and potential fate and transport pathways for the pesticide, and the determination of potential exposure routes, (*e.g.*, ingestion, inhalation, and dermal contact).

Figure 3 depicts the potential exposure pathways associated with the proposed use of abamectin. The conceptual model generically depicts the potential source of abamectin, release mechanisms, abiotic and biotic receiving media, biological receptors, and attribute changes of potential concern and the measurement endpoints used to evaluate them.




Figure 3 Conceptual diagram for assessment of risks from abamectin use on various crops

Figure 3 depicts the potential exposure pathways associated with abamectin used as a foliar application to almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes. Based on the use pattern for abamectin, the main exposure pathways for terrestrial organisms are direct exposure to abamectin via consumption of food items. In the figure above, the dashed line represents the pathways of exposure that are unlikely to occur because of physical or chemical properties. Although abamectin has a log K_{ow} of 4.4, BCF in bluegill sunfish were in the range of 19-69 (whole fish) and 6.6-33 (fillet); indicating that bioconcentration in aquatic organisms is low. Volatilization is also not expected to be a concern based on the vapor pressure of abamectin (1.5×10^{-9} Torr).

2.6 Analysis Plan

This assessment focuses on adverse acute and chronic reproductive effects to terrestrial and aquatic wildlife associated with proposed abamectin foliar application use on almonds, walnuts, apples, avocados, celeriac, citrus, cotton, cucurbit, fruiting vegetables,

grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes. This analysis plan identifies the approach, methods, specific models, information, and data that will be used to estimate and evaluate risks from proposed labeled uses of abamectin based on the conceptual model and risk hypotheses.

This assessment focuses on adverse acute and chronic reproductive effects to terrestrial and aquatic wildlife associated with proposed abamectin use. This analysis plan identifies the approach, methods, specific models, information, and data that will be used to estimate and evaluate risks from proposed labeled uses of abamectin based on the conceptual model and risk hypotheses.

2.6.1 Conclusions from Previous Risk Assessments

An ecological risk assessment evaluating abamectin for foliar ground application on citrus (DP 210767) concluded that the abamectin may pose acute and chronic risks to birds and small herbivorous mammals. This assessment also concluded that ground applications of abamectin to citrus may pose acute and chronic risks to freshwater and estuarine/marine invertebrates.

2.6.2 Preliminary Identification of Data Gaps

This assessment is potentially underestimating risk to both terrestrial and aquatic organisms from exposure to abamectin. This potential underestimation is due to a lack of available toxicity data as well as technical issues with the data submitted for some species. Therefore, the following toxicity studies are requested:

- OPPTS 850.1400- Early Life-Stage Toxicity Test. There are no chronic toxicity data available for the Agency to assess chronic risk of abamectin to estuarine/marine fish.
- OPPTS 850.4225 – Seedling Emergence, Tier II and OPPTS 850.4250 – Vegetative Vigor, Tier II. Seedling emergence and vegetative vigor toxicity data are not available for terrestrial plants.
- OPPTS 850.2300 - Avian reproduction Study. A reproduction study with bobwhite quail is not available.
- OPPTS 850.2100 - Acute Oral Toxicity with a Passerine Bird. An acute oral toxicity study with a passerine bird is not available. No species recommended at this point. Protocol should be submitted prior to test initiation.
- Whole Sediment Toxicity Test: Chronic Invertebrates Freshwater and Marine. Based on the physiochemical properties, abamectin may sorb to organic materials in sediment and may be toxic to organisms that dwell in and ingest sediment as abamectin is very highly toxic to other aquatic invertebrates. Since abamectin is a

foliar application, spray drift to both freshwater and estuarine-marine environments is possible. The concentration of abamectin in water from spray drift from ground or aerial application is greater than the acute EC₅₀ value for the estuarine/marine mysid shrimp. 40 CFR Part 158.630 requires a chronic freshwater sediment study if the half-life is greater than or equal to 10 days and any of the following conditions exist: i. $K_d \geq 50$, ii. the $\log K_{ow} \geq 3$, or iii. the $K_{oc} \geq 1000$. Abamectin meets these criteria. A protocol should be submitted to the Agency for review prior to testing.

- OPPTS 850.1075 – Fish Acute Toxicity Test, freshwater and marine; 850.1010-Aquatic Invertebrate Acute Toxicity test with Daphnia; 850.1025 or 1055 – Oyster Acute Toxicity Test (shell deposition) or Bivalve Acute Toxicity Test (embryo-larvae). The registrant submitted test were conducted as static tests that were conducted above the reported water solubility, conducted using a potential photosensitizing solvent (acetone), and test concentrations were not measured. As a result, the actual test concentrations (dissolved bioavailable abamectin) are not known which may be underestimating risk. Therefore, a new acute toxicity study for a coldwater and warmwater freshwater fish, estuarine-marine fish and *Daphnia magna* is requested. An oyster shell deposition or a bivalve embryo-larvae toxicity study is also requested.
- OPPTS 850.1300 – Daphnia Chronic Toxicity Test. The registrant submitted chronic daphnia toxicity test did not measure growth for the surviving adults at test termination. The study indicates that the surviving daphnia in the two lowest concentrations tested were pale and smaller than the control. Measurement of growth is required under the current guidance. Therefore, a new study is requested.
- OPPTS 850. 5400 – Algal Toxicity and 850.4400 Aquatic Plant Toxicity Test using Lemna spp. There are limited studies (data on two of the five species available (duckweed and a green alga study)) addressing the toxicity of abamectin to aquatic plants; the studies conducted with duckweed and green algae were conducted above solubility, with a potential photosensitizing solvent (acetone), and test concentrations were not measured. Abamectin toxicity studies with a marine diatom, freshwater diatom and blue-green algae are requested as well as new studies for the green algae and duckweed.
- Submitted field dissipation studies are unacceptable; therefore, the behavior of abamectin in the field as compared to the laboratory cannot be demonstrated. In most cases we would expect dissipation in the field to be greater than that predicted by laboratory studies due to pesticide transport.

3.0 Analysis

3.1 Exposure Characterization

Abamectin is moderately persistent in the environment. The reported laboratory soil aerobic half-life was 115 days. Abamectin is relatively stable to hydrolysis but may undergo direct photolysis (photolysis half-life in surface soil = 21 hours). Abamectin has low vapor pressure (1.5×10^{-9} Torr), indicating that volatilization from dry soil surfaces will not be an important environmental fate process. An estimated Henry's Law constant of 2.6×10^{-8} atm-m³/mol was derived from the vapor pressure and water solubility values provided by the registrant. This value suggests that volatilization from moist soil is not expected to be an important fate process. Abamectin adsorbs strongly to soil surfaces (reported K_{oc} values range from 2,531-12,051), and according to the FAO classification, abamectin is slightly to hardly mobile in soil and that leaching to groundwater will not be an important route of dissipation.

If abamectin was to contaminate surface water, photolysis in sunlit surface waters would be an important environmental fate process based on an aqueous photolysis half-life of 12 hours. Volatilization from water is not expected to be an important fate process based on the estimated Henry's Law constant. The large K_{oc} values suggest that adsorption to suspended solids and sediment in the water column will occur. Bioconcentration factors (BCF) in bluegill sunfish were in the range of 19-69 (whole fish) and 6.6-33 (fillet); suggesting bioconcentration in aquatic organisms is low.

3.1.1 Measures of Aquatic Exposure

3.1.1.1 Aquatic Exposure Modeling

At the screening risk assessment level for aquatic organisms, such as plants, fish, aquatic-phase amphibians, and invertebrates, computer simulation models are used to estimate acute (annual instantaneous peak) and chronic (21 and 60 day weighted average annual peaks for aquatic invertebrates and fish, respectively) residue levels of the dissolved pesticide active ingredient in surface water and sediment pore water and in bulk sediment from runoff and spray drift. These models calculate EECs in surface water and sediment using environmental fate data for abamectin. Monitoring data, if available, may also be used to determine EECs or to support the model's exposure estimates. PRZM-EXAMS as documented at www.epa.gov/oppefed1/models/water/index.htm is the model used to simulate the fate and transport of abamectin from a treated field to and in a receiving water body adjacent to the treated field. Cropping patterns, soil structure, and weather input data for the simulation modeling has been standardized for a number of crops, referred to as crop scenarios, to provide high-end estimates of runoff and soil erosion representative of the primary growing area for a given crop. The quality control checked crop scenarios and associated meteorological files available for use in a risk analysis are also found at the same web address under the bullet "PRZM crop scenario metadata".

PRZM-EXAMS model inputs for abamectin and its major degradate (a mixture of 8- α -hydroxy and a ring opened aldehyde derivative fate parameters (e.g., aerobic metabolism, photolysis, etc.) are listed in Table 4. The scenarios modeled reflect differences in weather and cropping patterns, soil structure, and abamectin application dates in different major growing areas. A screening assessment of estimated environmental concentrations (EECs) for abamectin and its major soil degradate (a mixture of 8- α -hydroxy and a ring opened aldehyde derivative) in surface water resulting from the proposed label uses was performed.

PRZM/EXAMS modeling output files are listed in Appendix B. Tier II Surface Water 1-in10 Year EECs (ppb) of abamectin in surface water from its new proposed uses from PRZM/EXAMS modeling are shown in Table 6.

Table 4 Surface water exposure inputs for PRZM/EXAMS

MODEL INPUT VARIABLE	INPUT VALUE	SOURCE and COMMENTS
Application rate (kg ai/hectare) and application interval	See Table 6	Some crops were modeled at 0.023 and 0.0235 lb ai/A but 0.0235 lb ai/A used to determine risk quotients
K _d (mL/g)	82 (average)	MRID 40856301; no data for degradate; Input guideline, 2002
Aerobic Soil Metabolic Half-life (days)	150	Total toxic residue half-life for parent and degradate (a mixture of 8- α -hydroxy and a ring opened aldehyde derivative)
Is the pesticide wetted-in?	No	EPA Reg. No. 100-RGLR
Spray Drift Fraction	0.05	Input guideline, 2002
Application Efficiency	0.95	Input guideline, 2002
Solubility (μ g/L)	78	10x reported value (7.8 μ g/L) per guidance (Input guideline, 2002); as there is no data for degradate it was assumed that it was no more soluble than the parent.
Aerobic Aquatic Metabolic Half-life (days)	300	No acceptable aerobic aquatic metabolism data were available, therefore 2x the aerobic soil metabolism half-life (identified above) was used per guidance (Input guideline, 2002).
Hydrolysis (pH 7) half-life (days)	0	Stable. No MRID available. Review dated 4/18/83; no data for degradate.
Aquatic Photolysis Half-life (days)	0.5	Dark-control adjusted half-life. Ku and Jacob, 1983 (Public literature, EFED Review dated 3/28/84); no data for degradate.

Table 5. Tier II Surface Water 1-in10 Year EECs (ppb) of abamectin and its major soil degradate (a mixture of 8- α -hydroxy and a ring opened aldehyde derivative)

Crop	Application Rate (lb ai/acre); (# Applications/ Application interval)	PRZM Scenario; method of application	Peak EEC (ppb)	21-day avg EEC (ppb)	60-day avg EEC (ppb)
Almonds & Walnuts	0.0235; (2/21) ¹	CAalmond_WirrigSTD	0.075	0.059	0.048
Apples	0.0235; (2/21) ¹	PAApplesSTD	0.339	0.266	0.214
Avocados	0.0235; (2/30) ¹	FLAvocadoSTD	0.142	0.111	0.102
Celeriac	0.0187; (3/7) ²	FLCarrotSTD	0.429	0.351	0.298
Citrus	0.0235; (2/30)	FLCitrusSTD	0.394	0.318	0.278
Cotton	0.019; (2/21)	MScottonSTD	0.420	0.348	0.291
Cucurbit	0.0187; (3/7) ²	FLcucumberSTD	0.540	0.446	0.386
Fruiting Veg	0.0187; (3/7) ²	FLpepperSTD	0.493	0.410	0.373
Grapes	0.019; (2/21)	NYgrapesstd	0.466	0.404	0.361
Herb	0.0187; (3/7) ²	ORMintSTD	0.084	0.075	0.065
Hops	0.019; (2,21)	ORhopsSTD	0.158	0.136	0.130
Leafy Veg	0.0187; (3/7) ²	FLcabbageSTD	0.277	0.217	0.174
Mint	0.014; (3/7)	ORMintSTD	0.156	0.129	0.107
Pears	0.0235; (2/21) ¹	WAorchards	0.029	0.023	0.020
Plums & Prunes	0.0235; (2/21) ¹	WAorchards	0.040	0.031	0.023
Potatoes	0.0187; (3/7) ²	MEpotatoSTD	0.651	0.564	0.498

¹ These crops were modeled using the maximum seasonal application rate divided by 2 applications.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

3.1.1.2 Aquatic Exposure Monitoring and Field Data

Groundwater and surface water monitoring data are not available. Screening models were used to determine estimated concentrations for abamectin in groundwater and surface water for the proposed uses.

3.1.2 Measures of Terrestrial Exposure

Avian and Mammalian Dietary Exposure

The Terrestrial Exposure (T-REX) model (Version 1.4. 1), an EFED computer model that uses a first-order dissipation relationship to account for residue dissipation between applications, was used to estimate exposure concentrations of abamectin to terrestrial wildlife. The T-REX simulation model incorporates the nomogram (Fletcher *et al.*, 1994; Hoerger and Kenaga, 1972; Pfleeger *et al.*, 1996) relationship between the amount of pesticide applied and the amount of pesticide residue present on a given food item. In addition to exposure concentrations (dose and diet-based), the T-REX model calculates risk quotients based on food items for mammals and birds, including herbivores, insectivores, and granivores. For dose-based exposures, three weight classes of mammals (15, 35, and 1000 g) and birds (20, 100, and 1000 g) are considered (Appendix C).

A default foliar dissipation half-life of 35 days was used in this assessment, although, residue concentrations may be lower as a honey-bee foliar residue study on citrus, demonstrates that residues are toxic above background levels for approximately 48 hours.

Since the label does not specifically state the interval between the second sequential application and subsequent applications for a number of crops (celeriace, cucurbit, fruiting vegetable, leafy vegetable, mint, herbs and potatoes (for potato psyllid), three applications at seven day intervals using the maximum seasonal rate divided by three (which is slightly less than three applications at the maximum single application rate, 0.0187 vs. 0.019 lb ai/A) was modeled for environmental exposure. The dietary exposure model T-REX can not model different application intervals or application rates at the same time. In addition, the application rate for almonds, walnuts, apples, citrus, avocados, pears, plums and prunes was modeled using the maximum seasonal application rate divided by two applications (0.0235 lb ai/A).

Input parameters, such as application rate, interval, and number of applications, used in T-REX model are presented with corresponding EECs in Table 6, Table 7, and Table 8.

Table 6. Avian Dose-Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Application of Abamectin

Crop; (Application Rate (lb ai/A); # of Applications; Application Interval (days))	Size Class (g) ¹	Avian Dose-Based EECs (ppm)				
		Dietary Item				
		Short Grass	Tall Grass	Broadleaf plants/ sm insects	Fruits/pods/ seeds/ lg insects	Granivore
Celeriac, cucurbit, fruiting and leafy	20	13.43	6.16	7.56	0.84	0.19
	100	7.66	3.51	4.31	0.48	0.11

vegetables, herbs, potato; (0.0187;3;7) ²	1000	3.43	1.57	1.93	0.21	0.05
Cotton, grapes, hops; (0.019;2;21)	20	8.62	3.95	4.85	0.54	0.12
	100	4.92	2.25	2.76	0.31	0.07
	1000	2.20	1.01	1.24	0.14	0.03
Almonds, walnuts, apple, pears, plums, prunes ; (0.0235;2;21) ²	20	10.66	4.89	6.00	0.67	0.15
	100	6.08	2.79	3.42	0.38	0.08
	1000	2.72	1.25	1.53	0.17	0.04
Avocados, citrus; (0.0235;2;30) ²	20	9.97	4.57	5.61	0.62	0.14
	100	5.68	2.61	3.20	0.36	0.08
	1000	2.55	1.17	1.43	0.16	0.04
Mint; (0.014;3;7)	20	10.06	4.61	5.66	0.63	0.14
	100	5.74	2.63	3.23	0.36	0.08
	1000	2.57	1.18	1.44	0.16	0.04

¹ Adjusted LD₅₀ (mg/kg-bw) based on avian body weight: 20 g = 44.13, 100 g = 56.18, 1000 g = 79.36

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.
These crops were modeled using the maximum seasonal application rate divided by 2 applications.

Table 7. Mammalian Dose-Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Application of Abamectin

Crop; (Application Rate (lb ai/A); # of Applications; Application Interval (days))	Size Class (g) ¹	Mammalian Dose-Based EECs (ppm)				
		Dietary Item				
		Short Grass	Tall Grass	Broadleaf plants/ sm insects	Fruits/pods/ seeds/ lg insects	Granivore
Celeriac, cucurbit, fruiting and leafy vegetables, herbs, potato; (0.0187;3;7) ²	15	11.25	5.15	6.33	0.70	0.16
	35	7.77	3.56	4.37	0.49	0.11
	1000	1.80	0.83	1.01	0.11	0.03
Cotton, grapes, hops; (0.019;2;21)	15	7.22	3.31	4.06	0.45	0.10
	35	4.99	2.29	2.81	0.31	0.07
	1000	1.16	0.53	0.65	0.07	0.02

Almonds, walnuts, apple, pears, plums, prunes ; (0.0235;2;21) ³	15	8.93	4.09	5.02	0.56	0.12
	35	6.17	2.83	3.47	0.39	0.09
	1000	1.43	0.66	0.80	0.09	0.02
Avocados, citrus; (0.0235;2;30) ³	15	8.35	3.83	4.69	0.52	0.12
	35	5.77	2.64	3.24	0.36	0.08
	1000	1.34	0.61	0.75	0.08	0.02
Mint; (0.014;3;7)	15	8.42	3.86	4.74	0.53	0.12
	35	5.82	2.67	3.27	0.36	0.08
	1000	1.35	0.62	0.76	0.08	0.02
¹ Adjusted LD ₅₀ (mg/kg-bw) based on mammalian body weight: 15 g = 29.89, 35 g = 24.18, 1000 g = 10.46; Adjusted NOAEL: 15 g = 0.26, 35 g = 0.21, 1000 g = 0.09 ² These crops were modeled using the maximum seasonal application rate divided by 3 applications. ³ These crops were modeled using the maximum seasonal application rate divided by 2 applications.						

Table 8. Dietary Based Estimated Environmental Concentrations (EECs) for Terrestrial Dietary Items from Foliar Exposure to Abamectin

Crop; (Application Rate (lb ai/A); # of Applications; Application Interval (days))	Dietary-Based EECs (ppm)			
	Dietary Item			
	Short Grass	Tall Grass	Broadleaf plants/ sm insects	Fruits/pods/seeds/ lg insects
Celeriac, cucurbit, fruiting and leafy vegetables, herbs, potato; (0.0187;3;7) ¹	11.80	5.41	6.64	0.74
Cotton, grapes, hops; (0.019;2;21)	7.57	3.47	4.26	0.47
Almonds, walnuts, pears, apple, plums, prunes ; (0.0235;2;21) ²	9.36	4.29	5.27	0.59
Avocados, citrus; (0.0235;2;30) ²	8.75	4.01	4.92	0.55
Mint; (0.014;3;7)	8.83	4.05	4.97	0.55
¹ These crops were modeled using the maximum seasonal application rate divided by 3 applications. ² These crops were modeled using the maximum seasonal application rate divided by applications				

Terrestrial Plants

There are no data regarding the explicit toxicity of abamectin to terrestrial plants. Therefore, no modeling of exposure for soil or foliar residues for terrestrial and semi-aquatic plants was performed.

3.2 Ecological Effects Characterization

In screening-level ecological risk assessments, effects characterization describes the types of effects a pesticide can produce in an organism or plant. This characterization is based on registrant-submitted studies that describe acute and chronic effects toxicity information for various aquatic and terrestrial animals and plants. All acceptable or supplemental guideline study data for technical grade abamectin, formulations, and degradates are summarized in Appendix D.

3.2.1.1 Terrestrial Animals

The most sensitive avian and mammalian acute and chronic toxicity test results and terrestrial invertebrates toxicity data selected for use in assessing baseline risk from abamectin are summarized in Table 9.

Birds

In birds, the acute toxicity of abamectin technical varies, depending on the species tested. The acute oral LD₅₀ for bobwhite quail (*Colinus virginianus*) is >2,000 mg ai/kg-bw (MRID 00129879, practically nontoxic), whereas the acute oral LD₅₀ for mallard ducks (*Anas platyrhynchos*) is 85 mg ai/kg-bw (MRID 00097859, moderately toxic). Regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity. The LC₅₀ values obtained in acceptable sub-acute dietary toxicity tests with bobwhite quail and mallard duck are >3,102 (MRID 00129880, slightly toxic) and 383 mg ai/kg-diet, respectively (MRID 00129520, highly toxic). A reproduction toxicity study with the bobwhite quail was not available. There were no statistically significant effects on growth, survival or reproduction in the mallard duck reproduction study at the highest concentration tested, 12 mg ai/kg-diet, therefore, the no observed adverse effect concentration (NOAEC) is at least 12 mg ai/kg-diet for the mallard duck chronic reproduction study (MRID 40318601). During the pilot study for the mallard duck reproduction study, the average number of eggs laid was markedly less in the 64 mg ai/kg treatment group.

Mammals

Based on data for laboratory rats, abamectin technical has an acute toxicity LD₅₀ value of 13.6 mg/kg-bw when using sesame oil as a delivery vehicle but 214 – 232 mg/kg-bw using a methyl cellulose delivery vehicle (MRID 0006894, 45607202). There are three prenatal developmental studies, three 1-generation reproduction studies and a 2-generation study with laboratory rats (Appendix D). The most sensitive reproductive endpoint was the 2-generation reproduction toxicity NOAEL value of 0.12 mg/kg-bw/day

based on increased retinal folds, increased dead pups at birth, decreased viability and lactation indices, and decreased pup body weight (MRID 00265576).

Although data exists for other routes of exposure (Appendix D), given the proposed application and the physical properties of the chemical, the expected significant route of exposure is oral. Therefore the focus of the risk estimation is on this route of exposure.

Terrestrial Invertebrates

Based on the honey bee LD₅₀ value of 0.41 µg/bee toxicity value, abamectin is highly toxic to terrestrial invertebrates (MRID 00159162). There was 13% mortality at 48-hrs at the lowest concentration tested for the acute contact study. A honey bee foliar exposure study indicated that exposure to abamectin treated citrus foliage is toxic for approximately 48 hours after application to the foliage (MRID 00159161). The proposed label states not to apply Agri-Flex SC or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Table 9. Summary of Most Sensitive Acute and Chronic Toxicity Data for Birds, Mammals and Terrestrial Invertebrates Exposed to Abamectin

Assessment Endpoint	Measurement Endpoint	Selected Measurement Endpoint Value and Source				
		Species	Study Duration	Toxicity Value	Most Sensitive Endpoint	Source and Study Classification
Survival and Reproduction of Birds	Most sensitive avian acute oral toxicity, LD ₅₀ (single-dose)	Mallard duck (<i>A. platyrhynchos</i>)	Single Oral Dose, post 14 day	LD ₅₀ = 85 mg a.i./kg-bw ¹	Mortality	00097859 Supplemental
	Most sensitive acute avian dietary toxicity	Mallard duck (<i>A. platyrhynchos</i>)	8 d (5 d exposure, post 3 d)	LC ₅₀ = 383 (mg ai/kg-diet)	Mortality	00129520 Acceptable
	Most sensitive avian reproductive toxicity NOAEC	Mallard duck (<i>A. platyrhynchos</i>)	18 Weeks	NOAEL ≥ 12 (mg ai/kg-diet), highest conc. tested ²	No statistically significant effect at highest conc. tested.	40318601 Acceptable
Survival and Reproduction of Terrestrial Mammals	Most sensitive acute oral toxicity, LD ₅₀ (single-dose)	Rat	Single oral dose	LD ₅₀ 13.6 mg /kg-bw	Mortality	00006894
	Most sensitive reproduction NOAEL	Rat	2-gen reproduction	0.12 mg a.i./kg-bw/d	Reproduction ³	00265576
Survival of Terrestrial Invertebrates and beneficial insects	Most sensitive acute contact LD ₅₀ (µg/bee)	Honey bee (<i>Apis mellifera</i>)	96-hr	LD ₅₀ = 0.41 µg per bee	Mortality	00159162 Acceptable
¹ Regurgitation observed in all treatment groups, therefore actual LD50 may be lower. ² In pilot test, marked decrease in average number of eggs laid at 64 ppm. ³ increased retinal folds, increased dead pups at birth, decreased viability and lactation indices, and decreased pup body weight.						

3.2.1.2 Terrestrial Plants

Registrant submitted seedling emergence or vegetative vigor toxicity data are not available for avermectin components, abamectin, or major degradates.

3.2.2 Aquatic Effects Characterization

3.2.2.1 Aquatic Animals

Abamectin is very highly toxic to both freshwater and estuarine/marine fish (Table 10). The 96-hr LC₅₀ values for rainbow trout (*Oncorhynchus mykiss*) and bluegill sunfish (*Lepomis macrochirus*) are 3.2 and 9.6 µg ai/L (total form (dissolved and undissolved abamectin)), respectively (MRID 00088780 and 00088782). For the estuarine/marine fish, sheepshead minnow (*Cyprinodon variegatus*), the 96-hr LC₅₀ value is 15 µg ai/L (total form) (MRID 00150910). All three of these reported fish studies were conducted above the reported limit of solubility for abamectin (7.8 ppb in distilled water; <1 ppb in tap water); acetone was used to increase abamectin solubility in water, and acetone can be a potential photosensitizer and abamectin undergoes rapid photolysis. These studies were based on nominal concentrations, as test solutions were not measured in these studies. Therefore, the actual concentrations of abamectin these organisms were exposed to are not known. An early life-cycle toxicity study was conducted with rainbow trout, and the reported no observed adverse effect concentration (NOAEC) was 0.52 µg ai/L (MRID 40069609) based on growth (wet weight).

An early life stage value for estuarine/marine fish has not been submitted to the Agency. However, an ACR of 6.2⁷ was calculated from the rainbow trout (*O. mykiss*) acute and chronic toxicity data, and was used to extrapolate from an acute 96-h LC₅₀ value for the sheepshead minnow to an early-life stage NOAEC. An acute to chronic ratio is available for both rainbow trout and aquatic invertebrates, but since abamectin is an insecticide, the mode of action is expected to be different for fish and invertebrates. Therefore the rainbow trout toxicity values were used to calculate the ACR. The extrapolated sheepshead NOAEC is 2.4 µg/L⁸.

Aquatic invertebrates are the aquatic species most sensitive to abamectin. It is very highly acutely toxic to aquatic invertebrates, with a 48-hr EC₅₀ value of 0.34 µg ai/L in the freshwater waterflea, *Daphnia magna* (MRID 00088784), and a 96-hr LC₅₀ of 0.020 µg ai/L in the estuarine/marine mysid shrimp, *Americamysis bahia* (MRID 40856305). Abamectin is highly toxic to the embryo/larval stages of mollusks with a 48-h EC₅₀ of 430 µg ai/L (total form) in the Eastern oyster (*Crassostrea virginica*) (MRID 00159158). The oyster embryo/larvae study was conducted above the water solubility limit of abamectin (7.8 ppb in distilled water; <1 ppb in tap water); acetone was used to increase solubility in water. Again, the daphnia and oyster larvae studies were evaluated using

⁷ *O. mykiss* ACR = 96-h LC₅₀/early-life stage NOAEC = 3.2 ppb/0.52 ppb = 6.2

⁸ Sheepshead Minnow early life stage NOAEC = 96-h LC₅₀/fish ACR = 15 ppb/6.2 = 2.4 ppb.

nominal concentrations, therefore, the actual concentrations these organisms were exposed to are not known. The life-cycle toxicity test with the *Daphnia magna* resulted in a reproductive NOAEC of 0.030 µg ai/L which was the lowest concentration tested, but the adults in the two lowest treatment groups were observed to be pale and smaller compared to the controls and growth was not analyzed (MRID 00153570). Therefore, the reproductive NOAEC appears to underestimate the true no-effect concentration for *Daphnia* from chronic exposure to abamectin, as the NOAEC appears to be lower than 0.030 µg ai/L (30 parts-per-trillion). An acute to chronic ration using the mysid shrimp toxicity data was used to calculate a chronic no-effect concentration for the daphnia and is 0.006 µg ai/L (6 parts-per-trillion)⁹. The NOAEC value for the life-cycle toxicity test with the mysid shrimp (*Americamysis bahia*) was previously reported as 0.0035 µg ai/L based on reproduction when compared to the solvent control, but is 0.00035 µg ai/L (0.35 parts-per-trillion) based on reproduction when compared to the negative control as there was a difference between the negative and solvent control for reproduction. Current EFED policy is to compare treatment groups to the negative control, therefore, the NOAEC value of 0.00035 µg ai/L was used in the assessment.

Table 10. Summary of Selected Acute and Chronic Toxicity Data for Fish and Aquatic Invertebrates Exposed to Abamectin for use in Determining Risk

Assessment Endpoint	Measurement Endpoint	Selected Measurement Endpoint Value and Source				
		Species	Study Duration	Toxicity Value	Most Sensitive Endpoint	Source and Study Classification
Survival and reproduction of freshwater vertebrates (fishes, etc)	Most sensitive acute freshwater fish LC ₅₀	Rainbow trout (<i>Oncorhynchus mykiss</i>)	96 hr LC ₅₀ , Static	3.2 µg ai/L (total form) ¹	Mortality	00088780 Acceptable
	Most sensitive freshwater fish early life stage or life cycle NOAEC	Rainbow trout (<i>Oncorhynchus mykiss</i>)	60-day	NOAEC = 0.52 µg ai/L	Growth	40069609 Acceptable
Survival and reproduction of freshwater invertebrates	Most sensitive acute freshwater invertebrate LC ₅₀ (or EC ₅₀)	Water flea, (<i>Daphnia magna</i>)	48 hr EC ₅₀ , Static	0.34 µg ai/L	Immobilization and mortality	00088784 Acceptable
	Most sensitive aquatic invertebrate life cycle NOAEC	Water flea, (<i>Daphnia magna</i>)	21 day Flow-through	ACR = 0.006 µg ai/L ²	Reproduction and growth	00153570 Acceptable
Survival and reproduction of marine/ estuarine vertebrates (fishes, etc)	Most sensitive acute marine/ estuarine vertebrate LC ₅₀	Sheepshead minnow (<i>Cyprinodon variegatus</i>)	96 hr Static-renewal	15 µg ai/L (total form) ¹	Mortality	00150910 Supplemental
	Most sensitive marine/estuarine fish early life stage or life cycle NOAEC	Sheepshead minnow (<i>Cyprinodon variegatus</i>)	28 day	No data available; ACR used value = 2.4 µg ai/L	NA	NA

⁹ Mysid shrimp ACR = 96-h EC₅₀/reproduction NOAEC = 0.020 ppb/0.00035 ppb = 57
Daphnia chronic NOAEC = 48-hr EC₅₀/mysid ACR = 0.34 ppb/57 = 0.006 ppb

Assessment Endpoint	Measurement Endpoint	Selected Measurement Endpoint Value and Source				
		Species	Study Duration	Toxicity Value	Most Sensitive Endpoint	Source and Study Classification
Survival and reproduction of marine/estuarine invertebrates	Most sensitive marine/estuarine acute mollusk shell deposition or embryo larval EC ₅₀	Eastern oyster (<i>Crassostrea virginica</i>) embryo/larvae	96 hr EC ₅₀ Static	430 µg ai/L (total form) ¹	Embryo development	00159158 Supplemental
	Most sensitive marine/estuarine acute invertebrate EC ₅₀	Mysids (<i>Americamysis bahia</i>)	96 hour EC ₅₀ Flow-through	0.020 µg ai/L	Mortality	40856305 Acceptable
	Most sensitive marine/estuarine life cycle invertebrate NOAEC	Mysids (<i>Americamysis bahia</i>)	28 day Flow-through	NOAEC = 0.00035 µg ai/L	Reproduction	40856306 Supplemental
¹ Study conducted above limit of solubility for abamectin so value may contain both dissolved and undissolved abamectin. Studies used acetone to increase water solubility. ² Adult daphnia in two lowest treatment groups were reported as pale in coloration and small compared to controls (NOAEC may be less than 0.030 ppb) so an acute to chronic ratio was calculated using mysid shrimp toxicity data.						

3.2.2.2 Aquatic Plants

Abamectin has been tested for phytotoxicity with only two aquatic plant species of the five listed for testing under guideline testing. The IC₅₀ values based on biomass or growth rate measures obtained in these two studies are >100,000 ppb and 3,900 ppb for the green alga *Selenastrum capricornutum* and the vascular aquatic plant *Lemna gibba*, respectively (MRID 00088787 and 00088788) (Table 11). These studies were evaluated using nominal concentrations since test solutions were not measured. Also, the studies were conducted using acetone which is a potential photosensitizer and abamectin is subject to photolysis. Therefore, the actual test concentrations these organisms were exposed to are not known (Table 11).

Table 11. Summary of Acute Toxicity Data for Aquatic Plants Exposed to Abamectin

Assessment Endpoint	Measurement Endpoint	Selected Measurement Endpoint Value and Source				
		Species	Study Duration	Toxicity Value	Most Sensitive Endpoint	Source and Study Classification
Reduced biomass and growth rate of aquatic plants	Most sensitive vascular plant biomass and area under curve NOAEL and IC ₅₀	Duckweed (<i>Lemna gibba</i>)	14 day Static EC ₅₀	3,900 µg ai/L (total form) ¹	Frond number	00088787
				NOAEC 1,200 µg ai/L		
	Most sensitive nonvascular plant biomass and growth rate NOAEL and IC ₅₀	Green algae (<i>Selenastrum capricornutum</i>)	9 days static	>100,000 µg ai/L (total form) ^{1,2}	Biomass	00088788
				NOAEC = Not Available		

Assessment Endpoint	Measurement Endpoint	Selected Measurement Endpoint Value and Source				
		Species	Study Duration	Toxicity Value	Most Sensitive Endpoint	Source and Study Classification
¹ Concentrations tested were above the solubility in water (7.8 ppb in distilled) so test solutions may contain both dissolved and undissolved abamectin. Acetone was used to increase solubility in water.						
² Precipitate was observed at concentrations of 25,000 ppb and above.						

4.0 Risk Characterization

Risk characterization is the integration of exposure and effects characterization to determine the ecological risk from the use of abamectin and the likelihood of effects on aquatic life, wildlife, and plants based on varying pesticide-use scenarios. The risk characterization provides estimation and a description of the risk; articulates risk assessment assumptions, limitations, and uncertainties; synthesizes an overall conclusion; and provides the risk managers with information to make regulatory decisions.

4.1 Risk Estimation – Integration of Exposure and Effects Data

Results of the exposure and toxicity effects data are used to evaluate the likelihood of adverse ecological effects on non-target species. For the assessment of abamectin risks, the risk quotient (RQ) method is used to compare exposure and measured toxicity values. Estimated environmental concentrations (EECs) are divided by acute and chronic toxicity values. The RQ's are compared to the Agency's levels of concern (LOCs). These LOCs are the Agency's interpretive policy and are used to analyze potential risk to non-target organisms and the need to consider regulatory action. These criteria are used to indicate when a pesticide's use as directed on the label has the potential to cause adverse effects on non-target organisms. The LOC's are listed in Appendix E.

4.1.1 Non-target Aquatic Animals and Plants

4.1.1.1 *Non-target Aquatic Animals*

Surface water concentrations resulting from abamectin application were predicted with the PRZM-EXAMS model. These aquatic estimated environmental concentrations (EEC's) are listed in Table 6. Peak EECs were then compared to acute toxicity endpoints to derive acute RQ's. The 60- day EECs were compared to chronic toxicity endpoints (NOAEC values) to derive chronic RQ's for fish, and 21-day EECs were compared to chronic toxicity endpoints (NOAEC values) for aquatic invertebrates. Acute RQ's for freshwater and estuarine/marine organisms for different exposure scenarios are presented in Table 12 and chronic RQ's for these species are presented in Table 13.

Fish and Aquatic Invertebrates

Acute

Non-Listed Species

There were no acute non-listed LOC exceedances for either freshwater or estuarine/marine fish. RQ values did exceed the acute non-listed LOC of 0.5 for freshwater aquatic invertebrates from abamectin use on apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes and potatoes. The acute estuarine/marine invertebrates RQ values also exceeded the acute non-listed LOC for all crop scenarios.

Listed Species

The acute freshwater and estuarine/marine invertebrate RQ values exceed the Agency's acute listed LOC of 0.05 for all crop scenarios. The acute freshwater fish RQ values exceed the Agency's acute listed LOC for abamectin application to apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes, and potatoes. None of the crop scenario RQ values exceeded the listed LOC for estuarine/marine fish.

Chronic

Chronic freshwater and estuarine/marine invertebrate RQ's exceed the chronic LOC (1.0) for all crop scenarios. Freshwater fish and estuarine/marine fish chronic RQ values do exceed the chronic LOC for any crop scenario.

Table 12. Acute Risk Quotients for Fish and Aquatic Invertebrates from Abamectin Applied to Various Crops

Crop Scenario	Application Rate (lb ai/acre); (# Applications/ Application interval)	Calculated EECs	Freshwater Fish ^a	Freshwater Invertebrates ^b	Estuarine/ Marine Fish ^c	Estuarine/ Marine Invertebrates ^d
		Peak (µg/L)	LC ₅₀ = 3.2 µg/L	LC ₅₀ = 0.34 µg/L	LC ₅₀ = 15.0 µg/L	LC ₅₀ = 0.02 µg/L
Almonds & Walnuts	0.0235; (2/21) ¹	0.075	0.023	0.219	0.005	3.73*
Apples	0.0235; (2/21) ¹	0.339	0.106	0.997*	0.023	17.0*
Avocados	0.0235; (2/30) ¹	0.142	0.044	0.418	0.009	7.10*
Celeriac	0.0187; (3/7) ²	0.429	0.134	1.26*	0.029	21.5*
Citrus	0.0235; (2/30) ¹	0.394	0.123	1.16*	0.026	19.7*

Crop Scenario	Application Rate (lb ai/acre); (# Applications/ Application interval)	Calculated EECs	Freshwater Fish ^a	Freshwater Invertebrates ^b	Estuarine/ Marine Fish ^c	Estuarine/ Marine Invertebrates ^d
		Peak (µg/L)	LC ₅₀ = 3.2 µg/L	LC ₅₀ = 0.34 µg/L	LC ₅₀ = 15.0 µg/L	LC ₅₀ = 0.02 µg/L
Cotton	0.019; (2/21)	0.420	0.131	1.24*	0.028	21.0*
Cucurbit	0.0187; (3/7) ²	0.540	0.169	1.59*	0.036	27.0*
Fruiting Veg	0.0187; (3/7) ²	0.493	0.154	1.45*	0.033	24.7*
Grapes	0.019; (2/21)	0.466	0.146	1.37*	0.031	23.3*
Herb	0.0187; (3/7) ²	0.084	0.026	0.247	0.006	4.20*
Hops	0.019; (2/21)	0.158	0.049	0.465	0.011	7.90*
Leafy Veg	0.0187; (3/7) ²	0.277	0.087	0.815*	0.018	13.9*
Mint	0.014; (3/7)	0.156	0.049	0.459	0.010	7.80*
Pears	0.0235; (2/21) ¹	0.029	0.009	0.085	0.002	1.45*
Plums & Prunes	0.0235; (2/21) ¹	0.040	0.013	0.118	0.003	2.00*
Potatoes	0.0187; (3/7) ²	0.651	0.203	1.91*	0.043	32.6*

¹ These crops were modeled using the maximum seasonal application rate divided by 2 applications.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications
 Bolded RQ values exceed the Agency's acute listed LOC (0.05) for direct effects to listed species

* = RQ values exceed the Agency's non-listed acute LOC (0.5) for non-listed species

^a Based on Rainbow Trout (*Oncorhynchus mykiss*)

^b Based on Water Flea (*Daphnia magna*)

^c Based on Sheepshead Minnow (*Cyprinodon variegatus*)

^d Based on Mysid Shrimp (*Americamysis bahia*)

Table 13. Chronic Risk Quotients for Fish and Aquatic Invertebrates from Abamectin Applied to Various Crops

Crop Scenario	Application Rate (lb ai/acre); (# Applications/ Application interval)	Calculated EECs (µg/L)		Freshwater Fish ^a	Estuarine/ Marine Fish ^b	Freshwater Invertebrates ^c	Estuarine/ Marine Invertebrates ^d
		21-d ³	60-d ³	NOAEC = 0.52 µg/L	NOAEC = 2.4 µg/L	NOAEC = 0.006 µg/L	NOAEC = 0.00035 µg/L
Almonds & Walnuts	0.0235; (2/21) ¹	0.059	0.048	0.09	0.02	9.83	169
Apples	0.0235; (2/21) ¹	0.266	0.214	0.41	0.09	44.3	760

Avocados	0.0235; (2/30) ¹	0.111	0.102	0.20	0.04	18.5	317
Celeriac	0.0187; (3/7) ²	0.351	0.298	0.57	0.12	58.5	1003
Citrus	0.0235; (2/30) ¹	0.318	0.278	0.53	0.12	53.0	909
Cotton	0.019; (2/21)	0.348	0.291	0.56	0.12	58.0	994
Cucurbit	0.0187; (3/7) ²	0.446	0.386	0.74	0.16	74.3	1274
Fruiting Veg	0.0187; (3/7) ²	0.410	0.373	0.72	0.15	68.3	1171
Grapes	0.019; (2/21)	0.404	0.361	0.69	0.15	67.3	1154
Herb	0.0187; (3/7) ²	0.075	0.065	0.13	0.03	12.5	214
Hops	0.019; (2,21)	0.136	0.130	0.25	0.05	22.7	389
Leafy Veg	0.0187; (3/7) ²	0.217	0.174	0.33	0.07	36.2	620
Mint	0.014; (3/7)	0.129	0.107	0.21	0.04	21.5	369
Pears	0.0235; (2/21) ¹	0.023	0.020	0.04	0.01	3.83	65.7
Plums & Prunes	0.0235; (2/21) ¹	0.031	0.023	0.04	0.01	5.17	88.6
Potatoes	0.0187; (3/7) ²	0.564	0.498	0.96	0.21	94.0	1611

¹ These crops were modeled using the maximum seasonal application rate divided by 3 applications.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications

³ Freshwater and estuarine/marine invertebrates NOAEC values were compared to the 21-day EEC, and freshwater and estuarine/marine fish NOAEC values were compared to the 60-day EEC.

Bolded RQ values exceed the Agency's chronic LOC (1.0)

^a Based on Rainbow Trout (*Oncorhynchus mykiss*)

^b Estimated early life stage NOAEC using an ACR of 6.2

^c Estimated using an ACR of 57 (Based on Water Flea (*Daphnia magna*)) and mysid shrimp)

^d Based on Mysid Shrimp (*Americamysis bahia*)

4.1.1.2 Aquatic Plants

Calculated peak EECs were compared to IC₅₀ endpoints for to derive aquatic vascular and non-vascular plant RQ's for non-listed species, and the peak EECs were compared to the aquatic vascular NOAEC value to derive RQ's for listed species. Listed species RQ values were not calculated for the non-vascular species (*Selenastrum capricornutum*) as a NOAEC value was not available. Acute RQ's for aquatic vascular and nonvascular plants are summarized in Table 14. RQ values did not exceed the plant LOC of 1.0 for any crop. However, data for only two of the five species was available for review. In addition, submitted studies were conducted as nominal concentrations with the use of a potential photosensitizing solvent; therefore, risk may be underestimated.

Table 14. Risk quotients for Aquatic Plants Exposed to Foliar Applications of Abamectin

Crop Scenario	Application Rate (lb ai/acre); (# Applications/ Application interval)	Calculated EECs Peak (µg/L)	Vascular	Vascular	Non-Vascular
			Non-Listed ^a	Listed ^a	Non-Listed ^b
			IC ₅₀ = 3,900 ppb	NOAEC = 1,200 ppb	IC ₅₀ >100,000 ppb
Almonds & Walnuts	0.0235; (2/21) ¹	0.075	<0.01	<0.01	<0.01
Apples	0.0235; (2/21) ¹	0.339	<0.01	<0.01	<0.01
Avocados	0.0235; (2/30) ¹	0.142	<0.01	<0.01	<0.01
Celeriac	0.0187; (3/7) ²	0.429	<0.01	<0.01	<0.01
Citrus	0.0235; (2/30) ¹	0.394	<0.01	<0.01	<0.01
Cotton	0.019; (2/21)	0.420	<0.01	<0.01	<0.01
Cucurbit	0.0187; (3/7) ²	0.540	<0.01	<0.01	<0.01
Fruiting Veg	0.0187; (3/7) ²	0.493	<0.01	<0.01	<0.01
Grapes	0.019; (2/21)	0.466	<0.01	<0.01	<0.01
Herb	0.0187; (3/7) ²	0.084	<0.01	<0.01	<0.01
Hops	0.019; (2,21)	0.158	<0.01	<0.01	<0.01
Leafy Veg	0.0187; (3/7) ²	0.277	<0.01	<0.01	<0.01
Mint	0.014; (3/7)	0.156	<0.01	<0.01	<0.01
Pears	0.0235; (2/21) ¹	0.029	<0.01	<0.01	<0.01
Plums & Prunes	0.0235; (2/21) ¹	0.040	<0.01	<0.01	<0.01
Potatoes	0.0187; (3/7) ²	0.651	<0.01	<0.01	<0.01

¹ These crops were modeled using the maximum seasonal application rate divided by 2 applications.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

^a Based on Duckweed (*Lemna gibba*)

^b Based on (*Selenastrum capricornutum*)

4.1.1.3 Non-target Terrestrial Animals

The RQ's for avian species are summarized in Table 15 through Table 17, and mammalian RQ's are summarized in Table 18 through Table 20. EEC comparisons to terrestrial invertebrate toxicity are summarized in Table 21.

Acute Avian Risk

Non-Listed Species

The acute dose-based and dietary-based RQ values for birds did not exceed the non-listed LOC of 0.5 for any crop scenario (Table 15 and Table 16). However, regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity.

Listed Species

Acute avian dietary-based RQ values did not exceed the acute endangered LOC of 0.1 for any crop scenario. However, the acute avian dose-based RQ values exceeded the acute listed LOC for small birds feeding on small and tall grass, broadleaf plants and small insects for all crop scenarios, except for tall grasses for cotton, grapes and hops. Acute avian dose-based RQ values also exceed the acute listed LOC for medium birds consuming short grasses for all crops except cotton, grapes and hops (Table 15 and Table 16).

Chronic Avian Risk

For the mallard duck chronic reproduction toxicity study, the highest concentration tested (12 mg ai/kg) resulted in no statistically significant effect for survival, growth or reproduction, therefore, chronic RQ values were not calculated. This highest tested concentration, 12 mg ai/kg, was compared to the calculated EECs, and all EECs were lower than this tested concentration (Table 17).

Table 15. Upper bound acute dose-based RQ values for birds for foliar application of abamectin

Crop and Application Rate	Functional Feeding Group Dietary Item	20 g bird Acute ¹	100 g bird Acute ¹	1000 g bird Acute ¹
<u>Celeriac, cucurbit, fruiting and leafy veg., herbs, potato</u> ² 0.0187 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.30	0.14	0.04
	Tall Grass	0.14	0.06	0.02
	Broadleaf plants/sm insects	0.17	0.08	0.02
	Fruits/pods/lg insects	0.02	0.01	<0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Cotton, grapes, hops</u>	<i>Herbivores/Insectivores</i>			
	Short Grass	0.20	0.09	0.03
	Tall Grass	0.09	0.04	0.01

Crop and Application Rate	Functional Feeding Group Dietary Item	20 g bird Acute ¹	100 g bird Acute ¹	1000 g bird Acute ¹
0.019 lb ai/A/ 2 Apps/21-d interval	Broadleaf plants/sm insects	0.11	0.05	0.02
	Fruits/pods/lg insects	0.01	0.01	<0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Almonds, walnuts, apple, pears, plums, prunes</u> ³ 0.0235 lb ai/A/ 2 apps/21-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.24	0.11	0.03
	Tall Grass	0.11	0.05	0.02
	Broadleaf plants/sm insects	0.14	0.06	0.02
	Fruits/pods/lg insects	0.02	0.01	<0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Avocado, citrus</u> ³ 0.0235 lb ai/A/ 2 apps/30-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.23	0.10	0.03
	Tall Grass	0.10	0.05	0.01
	Broadleaf plants/sm insects	0.13	0.06	0.02
	Fruits/pods/lg insects	0.01	0.01	<0.01
	<i>Granivore</i>			
<u>Mint</u> 0.014 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.23	0.10	0.03
	Tall Grass	0.10	0.05	0.01
	Broadleaf plants/sm insects	0.13	0.06	0.02
	Fruits/pods/lg insects	0.01	0.01	<0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01

Bolded RQ values exceed the listed LOC of 0.1;

¹ Acute RQ = (upper bound dose-based EEC, mg/kg-bw) / (LD₅₀; mg/kg-bw). The upper bound EECs for a given body weight and LD₅₀ values adjusted for the given body weight are in Table 6.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

³ These crops were modeled using the maximum seasonal application rate divided by 3 applications

Table 16. Upper Bound Acute Avian Dietary-based RQ values from Foliar Application of Abamectin to Celeriac, Cucurbit, Fruiting and Leafy Vegetables, Herbs and Potato

Crop and Application Rate	Dietary Item	EEC (mg/kg-diet) ¹	Acute Dietary RQ ²
Celeriac, cucurbit, fruiting and leafy veg., herbs, potato 0.0187 lb ai/A/ 3 apps/7-d interval	Short Grass	11.80	0.03
	Tall Grass	5.41	0.01
	Broadleaf plants/sm Insects	6.64	0.02
	Fruits/pods/seeds/lg insects	0.74	<0.01

¹ Dietary-based residue levels for application from Table 8.

² Acute RQ = (EEC, mg/kg-diet) / acute dietary LC50, mg/kg-diet; where the acute dietary LC50 is 383 mg/kg-diet for the mallard duck from Table 9.

Table 17. Comparison of the Dietary EECs from Foliar Application of Abamectin to the Chronic Avian NOAEC

Crop and Application Rate	Dietary Item	EEC (mg/kg-diet) ¹	Chronic Avian NOAEC ² (mg ai/kg-diet)
Celeriac, cucurbit, fruiting and leafy veg., herbs, potato 0.0187 lb ai/A/ 3 apps/7-d interval	Short Grass	11.80	< 12
	Tall Grass	5.41	<12
	Broadleaf plants/sm Insects	6.64	<12
	Fruits/pods/seeds/lg insects	0.74	<12

¹ Dietary-based residue levels for applications from Table 8.

² the chronic NOAEC is 12 mg ai/kg-diet for the mallard duck, the highest dose tested Table 9.

Acute Mammalian Risk

Non-Listed Species

No acute dose-based RQ values exceeded the acute LOC (0.5) for non-listed mammalian species in any scenario tested (Table 18).

Listed Species

Acute dose-based RQ values exceed the Agency's listed LOC of 0.1 for small and medium mammals consuming short and tall grass, broadleaf plants and small insects for all crops except for medium mammals consuming tall grass for cotton, grapes and hops. The acute dose-based listed LOC was also exceeded for large mammals feeding on short grasses for all crop scenarios and broadleaf plants and small insects for abamectin application to celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes (Table 18).

Chronic Mammalian Risk

Chronic dose-based RQ values exceed the Agency's chronic LOC (1.0) for small, medium and large mammals feeding on short grass, tall grass, broadleaf plants, small

insects, fruits, pods or large insects for all crops, except for large mammals consuming fruits, pods and large insects in which only abamectin use on celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes exceeded the LOC for fruits, pods and large insects. No chronic dose-based RQ values exceeded the Agency's chronic LOC for mammals feeding on seeds (Table 19).

Chronic dietary-based RQ values exceeded the LOC for mammals consuming short and tall grass, broadleaf plants and small insects for all crops. No chronic dietary-based RQ values exceeded the chronic LOC for mammals consuming fruits, pods, seeds, or large insects (Table 20).

Table 18. Upper bound Mammalian Acute Dose-based RQ values for Foliar Application of Abamectin

Crop	Functional Feeding Group Dietary Item	15 g mammals Acute ¹	35 g mammals Acute ¹	1000 g mammals Acute ¹
<u>Celeriac, cucurbit, fruiting and leafy veg., herbs, potato</u> ² 0.0187 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.38	0.32	0.17
	Tall Grass	0.17	0.15	0.08
	Broadleaf plants/sm insects	0.21	0.18	0.10
	Fruits/pods/lg insects	0.02	0.02	0.01
	<i>Granivore</i>			
	Seeds	0.01	<0.01	<0.01
<u>Cotton, grapes, hops</u> 0.019 lb ai/A/ 2 Apps/21-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.24	0.21	0.11
	Tall Grass	0.11	0.09	0.05
	Broadleaf plants/sm insects	0.14	0.12	0.06
	Fruits/pods/lg insects	0.02	0.01	0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Almonds, walnuts, apple, pears, plums, prunes</u> ³ 0.0235 lb ai/A/ 2 apps/21-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.30	0.26	0.14
	Tall Grass	0.14	0.12	0.06
	Broadleaf plants/sm insects	0.17	0.14	0.08
	Fruits/pods/lg insects	0.02	0.02	0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Avocado, citrus</u> ³ 0.0235 lb ai/A/ 2	<i>Herbivores/Insectivores</i>			
	Short Grass	0.28	0.24	0.13
	Tall Grass	0.13	0.11	0.06
	Broadleaf plants/	0.16	0.13	0.07

Crop	Functional Feeding Group Dietary Item	15 g mammals Acute ¹	35 g mammals Acute ¹	1000 g mammals Acute ¹
apps/30-d interval	sm insects			
	Fruits/pods/lg insects	0.02	0.01	0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01
<u>Mint</u> 0.014 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	0.28	0.24	0.13
	Tall Grass	0.13	0.11	0.06
	Broadleaf plants/ sm insects	0.16	0.14	0.07
	Fruits/pods/lg insects	0.02	0.02	0.01
	<i>Granivore</i>			
	Seeds	<0.01	<0.01	<0.01

Bolded RQ values exceed the listed LOC of 0.1;

¹ Acute RQ = (upper bound dose-based EEC, mg/kg-bw) / (LD₅₀, mg/kg-bw). The upper bound EECs for a given body weight and LD₅₀ values adjusted for the given body weight are in Table 6.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

³ These crops were modeled using the maximum seasonal application rate divided by 2 applications.

Table 19. Upper bound Mammalian Chronic Dose-based RQ values for Foliar Application of Abamectin

Crop	Functional Feeding Group Dietary Item	15 g mammals Acute ¹	35 g mammals Acute ¹	1000 g mammals Acute ¹
<u>Celeriac,</u> <u>cucurbit</u> <u>fruiting and</u> <u>leafy veg.,</u> <u>herbs, potato</u> ² 0.0187 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	42.64	36.43	19.53
	Tall Grass	19.55	16.70	8.95
	Broadleaf plants/ sm insects	23.99	20.49	10.98
	Fruits/pods/lg insects	2.67	2.28	1.22
	<i>Granivore</i>			
	Seeds	0.59	0.51	0.27
<u>Cotton,</u> <u>grapes, hops</u> 0.019 lb ai/A/ 2 Apps/21-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	27.36	23.37	12.53
	Tall Grass	12.54	10.71	5.74
	Broadleaf plants/ sm insects	15.39	13.15	7.05
	Fruits/pods/lg insects	1.71	1.46	0.78
	<i>Granivore</i>			
	Seeds	0.38	0.32	0.17
<u>Almonds,</u> <u>walnuts, pears,</u>	<i>Herbivores/Insectivores</i>			
	Short Grass	33.84	28.91	15.49

Crop	Functional Feeding Group Dietary Item	15 g mammals Acute ¹	35 g mammals Acute ¹	1000 g mammals Acute ¹
<u>apple, plums, prunes</u> 0.0235 lb ai/A/ 2 apps/21-d interval	Tall Grass	15.51	13.25	7.10
	Broadleaf plants/sm insects	19.04	16.26	8.72
	Fruits/pods/lg insects	2.12	1.81	0.97
	<i>Granivore</i>			
	Seeds	0.47	0.40	0.22
<u>Avocado, citrus</u> 0.0235 lb ai/A/ 2 apps/30-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	31.64	27.03	14.49
	Tall Grass	14.50	12.39	6.64
	Broadleaf plants/sm insects	17.80	15.20	8.15
	Fruits/pods/lg insects	1.98	1.69	0.91
	<i>Granivore</i>			
	Seeds	0.44	0.38	0.20
<u>Mint</u> 0.014 lb ai/A/ 3 apps/7-d interval	<i>Herbivores/Insectivores</i>			
	Short Grass	31.93	27.27	14.62
	Tall Grass	14.63	12.50	6.70
	Broadleaf plants/sm insects	17.96	15.34	8.22
	Fruits/pods/lg insects	2.00	1.70	0.91
	<i>Granivore</i>			
	Seeds	0.44	0.38	0.20

Bolded RQ values exceed the listed LOC of 1

¹ Chronic RQ = (upper bound dose-based EEC, mg/kg-bw) / (NOAEL; mg/kg-bw). The upper bound EECs for a given body weight and NOAEL values adjusted for the given body weight are in Table 6.

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

³ These crops were modeled using the maximum seasonal application rate divided by 2 applications

Table 20. Upper bound Chronic Dietary-based RQ Values for Mammals for Foliar Application of Abamectin

Crop and Application Rate	Dietary Item	EEC (mg/kg-diet) ¹	Chronic Mammalian RQ Value ¹
<u>Celeriac, cucurbit, fruiting and leafy veg., herbs, potato</u> ² 0.0187 lb ai/A/ 3 apps/7-d interval	Short Grass	11.80	4.92
	Tall Grass	5.41	2.25
	Broadleaf plants/sm Insects	6.64	2.76
	Fruits/pods/seeds/lg insects	0.75	0.31
<u>Cotton, grapes, hops</u> 0.019 lb ai/A/ 2 Apps/21-d interval	Short Grass	7.57	3.15
	Tall Grass	3.47	1.45
	Broadleaf plants/sm Insects	4.26	1.77
	Fruits/pods/seeds/lg insects	0.47	0.20

<u>Almonds, walnuts, apple, pears, plums, prunes</u> ³ 0.0235 lb ai/A/ 2 apps/21-d interval	Short Grass	9.36	3.90
	Tall Grass	4.29	1.79
	Broadleaf plants/sm Insects	5.27	2.19
	Fruits/pods/seeds/lg insects	0.59	0.24
<u>Avocado, citrus</u> ² 0.0235 lb ai/A/ 2 apps/30-d interval	Short Grass	8.75	3.65
	Tall Grass	4.01	1.67
	Broadleaf plants/sm Insects	4.92	2.05
	Fruits/pods/seeds/lg insects	0.55	0.23
<u>Mint</u> 0.014 lb ai/A/ 3 apps/7-d interval	Short Grass	8.83	3.68
	Tall Grass	4.05	1.69
	Broadleaf plants/sm Insects	4.97	2.07
	Fruits/pods/seeds/lg insects	0.55	0.23

Bolded RQ values exceed the listed LOC of 1

¹ Chronic RQ = (upper bound dietary-based EEC, mg/kg-diet) / (NOAEL; mg/kg-diet). The upper bound EECs for a crop are in Table 8 and chronic dietary NOAEL value is 2.40 mg/kg-diet, calculated from dose-based NOAEL of 0.12 mg/kg-bw

² These crops were modeled using the maximum seasonal application rate divided by 3 applications.

³ These crops were modeled using the maximum seasonal application rate divided by 3 applications

Terrestrial Invertebrates

Currently, there is not a method to quantify risk to non-listed terrestrial invertebrates. Abamectin is registered for use to control terrestrial invertebrates such as leafminers, mites, beetles, and ants; therefore, abamectin exposure to non-target terrestrial invertebrates is expected to also impact these non-target species. The acute contact abamectin LD₅₀ value for the honeybee is 0.41 µg ai/bee. This acute contact LD₅₀ value was converted to a body weight value using 0.128 g as the body weight of a bee. The extrapolated acute contact toxicity value for terrestrial invertebrates is 3.20 ppm.¹⁰ For the acute contact honeybee study, there was 13% mortality at the lowest concentration tested. Risk to insects were evaluated by comparing abamectin toxicity, as determined in the submitted honeybee acute contact study, with the residue levels from abamectin use on small and large insects generated as dietary-based EECs for birds and mammals using T-REX. Comparisons of the EECs for abamectin uses and the extrapolated acute toxicity are presented in Table 21. The small insect EECs are greater than the extrapolated acute contact value for all crops. So while the large insect EECs are less than the extrapolated LD₅₀ value, abamectin may still have the potential to cause adverse effects to terrestrial invertebrates as the acute contact toxicity data indicates that abamectin is highly toxic to

$$^{10} \text{ Extrapolated } LD50_{\text{terrestrial insect}} = \frac{LD50_{\text{honey bee}}}{BW_{\text{honey bee}}} = \frac{0.41 \text{ } \mu\text{g}}{0.128 \text{ g}} = 3.20 \text{ ppm}$$

the honeybee. Also, a foliage toxicity study indicated that foliar residues of abamectin may remain toxic to bees for two days following application.

Table 21. Comparisons of Small and Large Insect EECs from Foliar Application of Abamectin to the Extrapolated Acute Contact Honeybee Concentration

Application Rate (Crop)	Dietary Item	EEC (mg/kg-diet)	Extrapolated Acute Contact Value 3.20 (mg/kg)
<u>Celery, cucurbit, fruiting and leafy veg., herbs, potato</u> ¹ 0.0187 lb ai/A/ 3 apps/7-d interval	Small insects	6.64	>3.20
	Large insects	0.74	<3.20
<u>Cotton, grapes, hops</u> 0.019 lb ai/A/ 2 Apps/21-d interval	Small insects	4.26	>3.20
	Large insects	0.47	<3.20
<u>Almonds, walnuts, apple, pears, plums, prunes</u> ² 0.0235 lb ai/A/ 2 apps/21-d interval	Small insects	5.27	>3.20
	Large insects	0.59	<3.20
<u>Avocado, citrus</u> ² 0.0235 lb ai/A/ 2 apps/30-d interval	Small insects	4.92	>3.20
	Large insects	0.55	<3.20
<u>Mint</u> 0.014 lb ai/A/ 3 apps/7-d interval	Small insects	4.97	>3.20
	Large insects	0.55	<3.20

Bold values indicate the EEC exceeds the extrapolated acute contact value.

¹These crops were modeled using the maximum seasonal application rate divided by 3 applications.

²These crops were modeled using the maximum seasonal application rate divided by 3 applications

4.1.1.4 Non-target Terrestrial and Semi-Aquatic Plants

There are no toxicity data available to calculate RQ values for terrestrial and semi-aquatic plants.

4.2 Risk Description

The results of this risk assessment indicate that there are potential effects to listed freshwater fish species, listed and non-listed freshwater and estuarine/marine invertebrates, listed bird species, listed and non-listed mammalian species and terrestrial invertebrates from proposed new end-use abamectin product.

4.2.1 Risks to Aquatic Organisms

The proposed label indicates that Agri-Mek SC can not be applied within 25 ft for ground application or 150 ft for aerial application of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, natural ponds, estuaries or commercial fish farm ponds. In addition, the label restricts cultivation within 25 ft of the aquatic area to allow growth of a vegetative filter strip.

4.2.1.1 Fish and Aquatic Invertebrates

Calculated estimated exposure concentrations EECs from run-off and spray drift, based on modeling, potentially pose acute and chronic risks to listed and non-listed freshwater and estuarine/marine invertebrates and potentially acute risks to listed freshwater fish.

Acute

Non-Listed Species

Acute risk to non-listed fish is not expected as there were no acute non-listed LOC exceedances for either freshwater or estuarine/marine fish. RQ values did exceed the acute non-listed LOC of 0.5 for estuarine/marine invertebrates for all crops (RQs 1.45-32.6) and for freshwater aquatic invertebrates from abamectin use on apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes and potatoes.

Listed Species

Acute risk to listed estuarine/marine fish is not expected, as none of the crop scenario RQ values exceeded the listed LOC. The acute freshwater and estuarine/marine invertebrate RQ values exceed the Agency's acute listed LOC of 0.05 for all crop scenarios (RQs 0.085-1.91 for freshwater and 1.45-32.6 for estuarine/marine). The acute freshwater fish RQ values exceed the Agency's acute listed LOC for abamectin application to apples, celeriac, citrus, cotton, cucurbit, fruiting and leafy vegetables, grapes, and potatoes (RQs 0.087-0.203). In addition, fish are used as surrogates for aquatic phase amphibians and since there is potential risk to freshwater fish, risk to these species is also assumed.

Based on the calculated RQ values and a default concentration-response slope of 4.5, the probability of an individual mortality was calculated using the model IEC v1.1 (EPA, 2004a). For freshwater fish RQ values, this corresponds to a probability of mortality of less than 1 in 1 million to 1 in 1090, and for freshwater invertebrates, the probability of mortality ranges from less than 1 in 1.4 million to 1 in 1. Based on the calculated RQ's for estuarine/marine invertebrates, the probability of mortality is 1 in 1.

Chronic

Chronic risk to fish from abamectin use is not expected because the chronic RQ values did not exceed the LOC for any crop scenario. Chronic freshwater and estuarine/marine invertebrate RQ's exceed the chronic LOC (1.0) for all crop scenarios, except freshwater invertebrates exposed from abamectin application to pears (RQs 3.83-94.0 for freshwater and 65.7 -1611 for estuarine/marine).

The life-cycle toxicity test with the *Daphnia magna* resulted in a reproductive NOAEC of 0.030 µg ai/L which was the lowest concentration tested, but the adults in the two lowest treatment groups were observed to be pale and smaller compared to the controls (MRID 00153570). Therefore, the reproductive NOAEC appears to underestimate the true no effect concentration for *Daphnia* from chronic exposure to abamectin, as the NOAEC appears to be lower than 0.030 µg ai/L which may be underestimated risk. An extrapolated NOAEC value was calculated using the mysid shrimp toxicity data, but there is uncertainty as this extrapolated value may underestimate or overestimate risk.

4.2.1.2 Aquatic Plants

The aquatic plant RQ values did not exceed the acute non-listed or listed LOCs, however this is based on only two of the five guideline studies. These studies were conducted without measuring test concentrations, so the actual toxicity concentrations are not known. In addition, submitted studies were conducted with the use of a potential photosensitizing solvent; therefore, risk may be underestimated. If the nominal concentrations tested in the duckweed and green algae were maintained throughout the study, these untested species would have to be about 1,800 times more sensitive than current data indicate in order to exceed listed LOC's.

4.2.2 Risks to Terrestrial Organisms

4.2.2.1 Terrestrial Animals

Birds and Mammals

Acute

Non-Listed Species

Acute risk to non-listed birds and mammals from abamectin use is not expected, as the acute dose-based and dietary-based RQ values for birds and dose-based RQ values for mammals did not exceed the non-listed LOC of 0.5 for any crop scenario. However, regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity.

Listed Species

Acute dietary risk for birds is not expected as the avian acute dietary-based RQ values did not exceed the acute endangered LOC of 0.1 for any crop scenario. However, the acute avian dose-based RQ values exceed the acute listed LOC for small birds feeding on small and tall grass, broadleaf plants and small insects for all crop scenarios, except for tall grasses for cotton, grapes and hops, and the LOC was exceeded for medium birds consuming short grasses for all crops except for cotton, grapes and hops (RQs 0.10-0.30). Since birds are surrogates for reptiles and land-phase amphibians, the potential for direct effects may exist for these taxa as well.

Acute dose-based RQ values exceeded the LOC for small and medium mammals consuming short and tall grass, broadleaf plants and small insects for all crops except for medium mammals consuming tall grass for cotton, grapes and hops (RQs 0.11-0.38). The acute dose-based listed LOC was also exceeded for large mammals feeding on short grasses for all crop scenarios and broadleaf plants and small insects for abamectin application to celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes (RQs 0.10-0.17).

Based on the calculated RQ values and a concentration-response slope of 7.3 for the acute oral bird study and default concentration-response slope of 4.5 for mammals, the probability of an individual mortality was calculated using the model IEC v1.1 (EPA, 2004a). For the bird RQ values, this corresponds to a probability of mortality of less than 1 in seven trillion to 1 in 14,800, and for mammals, the probability of mortality ranges from less than 1 in 294,000 to 1 in 34.

Chronic

Chronic dose-based and dietary-based RQ values exceed the Agency's chronic LOC (1.0) for mammals feeding on short and tall grass, broadleaf plants and small insects (RQs 5.74-42.64 for dose-based and 1.45-4.92 for dietary based). Chronic dose-based RQ values also exceeded the LOC for small and medium mammals consuming fruits, pods or large insects for all crops and for large mammals for celeriac, cucurbit, fruiting and leafy vegetables, herbs and potatoes (RQs 1.22-2.67). No chronic dietary-based RQ values exceeded the chronic LOC for mammals consuming fruits, pods, seeds, or large insects or for seeds on a chronic dose basis.

For the mallard duck chronic reproduction toxicity study, the highest concentration tested (12 mg ai/kg) resulted in no statistically significant effect for survival, growth or reproduction, therefore, chronic RQ values were not calculated. This highest tested concentration, 12 mg ai/kg, was compared to the EECs, and all EECs were lower than this tested concentration.

The label states not to make more than two sequential applications of Agri-Mek SC, but the maximum seasonal amount allowed for these crops is greater than two applications at the maximum single application rate. Also, the maximum amount allowed per season for these crops is slightly less (0.0187 lb ai/A) than the amount applied using three applications at the maximum single application rate of 0.19 lb ai/A. Since the label does not specifically state the interval between the second sequential application and subsequent applications, three applications at seven day intervals using the maximum seasonal rate divided by three (0.0187 lb ai/A) was modeled for environmental exposure as the dietary exposure model T-REX can not model different application intervals or application rates at the same time. In addition, the application rate for almonds, walnuts, apples, citrus, avocados, pears, plums and prunes was modeled using the maximum seasonal application rate, 0.047 lb ai/A, divided by two applications (0.0235 lb ai/A).

The label indicates that the maximum single application rate for these crops is 0.023 lb ai/A, and with a maximum number of 2 applications, calculates 0.046. The label also indicates that the maximum seasonal application rate is 8.5 fl oz/A which calculates to 0.04648 lb ai/A, therefore it is not known if the reported 0.047 lb ai/A is due to rounding. Whether abamectin was modeled at 0.0235 or 0.023 lb ai/A, it resulted in exactly the same LOC exceedances.

In an effort to compare avian and mammalian acute and chronic dietary RQ's for other application scenarios, applications were modeled using the maximum single rate of 0.019 lb ai/A and three applications applied seven days apart. In addition, EECs were calculated using the maximum single application rate applied twice seven days apart with the assumption that subsequent applications would be applied at a later date in which the residues from the previous applications would have dissipated. For both birds and mammals using these two alternative application scenarios, the acute RQ values exceeded the listed LOC for exactly the same dietary items and body classes as the maximum seasonal application rate divided by three applications, except for large mammals consuming broadleaf plants and small insects for the two application scenario. Also, the chronic RQ values for mammals using the two alternative application methods exceeded the LOC for the same dietary items and body classes, except for large mammals consuming fruits, pods and large insects for the two application scenario. Therefore, except for large mammals consuming broadleaf plants, small and large insects, fruits and pods, acute and chronic RQ values will exceed the LOC whether abamectin is applied two or three times at the maximum single application rate or whether it is applied at the maximum seasonal rate divided by three applications.

Only the short grass EEC modeled using the maximum single rate of 0.019 lb ai/A and three applications applied every seven days was equal to the highest concentration tested in the mallard reproduction study (EEC = 11.99 vs. 12 ppm), but this modeling scenario is very slightly more (0.001 lb ai/A) than the maximum seasonal rate allowed (0.057 vs. 0.056 lb ai/A). In addition, EECs were calculated using the maximum single application rate applied twice seven days, and these EECs were lower than the mallard study concentration. Moreover, the level in which an adverse effect will not occur is not known but is observed to be at least 12 mg ai/kg. During the pilot study for the mallard reproduction study, the average number of eggs laid was markedly less in the 64 mg ai/kg treatment group. Overall, if two sequential applications at the single maximum application rate are applied seven days apart, and any subsequent application, even at the single maximum application rate, is applied more than seven days after the last application, the calculated EECs will be less than the highest concentration tested in the mallard reproduction study. Therefore, the potential for chronic risk to birds is not anticipated.

Terrestrial Invertebrates

Abamectin is highly toxic to the honeybee. The calculated EECs for small insects were greater than the extrapolated acute contact value (LD50) for the honeybee. Additionally, an incident was reported in EFED's Ecological Incident Information System (EIIS)

database (Incident No. I008611-001), where thousands of bees were killed during a registered use of abamectin on avocados in San Diego County CA in 1999. A foliar residue study on citrus demonstrated that foliar residues of abamectin are toxic to honeybees for approximately 48 hours after application (Appendix D). In addition, abamectin is registered for use to control terrestrial invertebrates such as leafminers, mites, beetles, and ants; therefore, abamectin exposure to non-target terrestrial invertebrates is expected to also impact these non-target species. Therefore, the proposed abamectin use is expected to be toxic to terrestrial invertebrates and beneficial insects.

The proposed label has environmental hazard labeling regarding bees and indicates not to apply when weather conditions favor drift from target areas, and that the product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. It also indicates not to apply the product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

4.2.2.2 *Terrestrial Plants*

There are no data regarding the toxicity of abamectin to terrestrial plants, therefore RQ values were not calculated.

According to the EHS incidence database there were three incidents for almonds in June 1998 from direct application of Agri-Mek in California (I007644-001, 002, 003). The type of injury to the almonds was not reported, but was reported to occur to all applied (34-106 acres). In addition, Agri-Mek was applied directly to 34 acres of grapes in June 2000 in California, with all 34 acres affected (I10837-019). The type of injury was not reported, and in the report, the inspector stated "Questionable" in regards to the question "Application within Label". All of these incidences were classified as possible.

Since there is no submitted toxicity data to evaluate terrestrial plants, and there are reported possible incidences for almonds and grapes, adverse risk to terrestrial plants can not be precluded.

4.2.3 *Federally Threatened and Endangered (Listed) Species Concerns*

4.2.3.1 *Taxonomic Groups potentially at Risk*

The Agency's LOC is exceeded for Federally listed Endangered and Threatened birds, mammals, and freshwater and estuarine/marine invertebrates for this proposed new end-use abamectin product for all listed crops (almonds, walnuts, apples, avocados, celery, citrus, cotton, cucurbit, fruiting vegetables, grapes, herbs, hops, leafy vegetables, mint, pears, plums, prunes and potatoes). The acute listed LOC is also exceeded for freshwater fish for abamectin use on apples, celery, citrus, cotton, cucurbit, fruiting and leafy vegetable, grapes, and potatoes. Since there is no data for reptiles and land-phase amphibians, birds were used as surrogates for these species, and due to potential risk to birds, risk to these species are assumed. In addition, fish are used as surrogates for aquatic phase amphibians and since there is potential risk to freshwater fish, risk to these

species is also assumed. Abamectin is highly toxic to bees, and the potential for adverse risk may occur from abamectin use. In addition, because of the lack of submitted terrestrial plant toxicity data and reported possible incidences involving almonds and grapes, adverse risk to terrestrial and semi-aquatic plants can not be precluded. A list of endangered/threatened species at the state level for these taxonomic groups and crops is attached to this assessment (Appendix F).

4.2.3.2 Direct and Indirect Effects

Due to the potential for direct effects to listed birds, reptiles, amphibians, mammals, fish, aquatic and terrestrial invertebrates, the potential for indirect effects may exist. The indirect effects may be from loss of the above species due to impacts on survival, growth, and reproduction. This loss may result in structural and functional changes of both the aquatic and terrestrial ecosystems. Changes may be manifested in the form of disruption of food chain and reduced biodiversity.

4.3 Description of Assumptions, Limitations, Uncertainties and Data Gaps.

4.3.1 Related to Exposure for All Species

4.3.1.1 General Exposure Parameters

- This screening-level risk assessment relies on labeled statements of the maximum rate of abamectin application, the maximum number of applications, and the shortest interval between applications. Together, these assumptions constitute a maximum use scenario. The frequency at which actual uses approach these maximums is dependant on resistance to the insecticide, timing of applications, and market forces.
- The label states that for a number of crops (celeriac, cucurbit, fruiting vegetable, leafy vegetable, mint and potatoes (for potato psyllid) not to make more than two sequential applications of Agri-Mek SC or any other foliar applied abamectin containing product, but the maximum seasonal amount allowed for these crops is greater than two applications at the maximum single application rate. The application interval for these crops is 7 days, and the label does not state how long to wait between the second sequential application and subsequent applications. Also, the maximum amount allowed per season for these crops, except mint, is slightly less (0.001 lb ai/A) than the amount applied using three applications at the maximum single application rate. Since the label does not specifically state the interval between the second sequential application and subsequent applications, three applications at seven day intervals using the maximum seasonal rate divided by three was modeled for environmental exposure. In addition, alternative application scenarios were also modeled and described in the Risk Characterization section (section 4.0)

- For application to herbs, the label states not to make more than two applications of Agri-Mek SC per single cutting (harvest), but the maximum amount allowed per cropping season is greater than two applications at the maximum single application rate but slightly less than three applications at the maximum single application rate. Therefore, environmental exposure concentrations were modeled in the same manner as discussed above.
- For application to almonds, walnuts, apples, avocados, citrus, pears, plums and prunes, the label states that for the maximum amount per season, not to apply more than 8.5 fl oz/A (or 0.047 lb ai/A) of Agri-Mek SC or any other foliar applied abamectin containing product in a growing season. Based on the density of the formulation, 8.5 fl oz/A calculates to 0.04648 lb ai/A, therefore, it is not known if the reported 0.047 lb ai/A is a rounding issue or if another abamectin product can be applied at 0.001 lb ai/A. In addition, the single maximum application rate reported is 0.023 lb ai/A, and two applications would be 0.046 lb ai/A. For this assessment, abamectin was modeled at 0.0235 lb ai/A (0.047 divided by two applications). Abamectin was also modeled at 0.023 lb ai/A which resulted in the same LOC exceedances as the 0.0235 lb ai/A application.
- The maximum seasonal application rate for cotton, potatoes (for Colorado potato beetle) and grapes on the label is reported as 0.038 lb ai/A, but the label also indicates not to apply more than 6.75 fl oz/A of Agri-Mek SC per season which calculates to 0.0369 (0.037) lb ai/A. The maximum single application rate for cotton, potatoes and grapes is 0.019 lb ai/A, and if applied twice per season, the maximum seasonal application rate would be 0.038 lb ai/A. Therefore, a maximum seasonal application rate of 0.038 lb ai/A was used for determining environmental exposure concentrations.

4.3.2 Related to Exposure Assessment

4.3.2.1 Related to Exposure for Aquatic Species

For an acute risk assessment, there is no averaging time for exposure. An instantaneous peak concentration, with a 1 in 10 year return frequency, is assumed. The use of the instantaneous peak assumes that instantaneous exposure is of sufficient duration to elicit acute effects comparable to those observed over more protracted exposure periods tested in the laboratory, typically 48 to 96 hours. In the absence of data regarding time-to-toxic event analyses and latent responses to instantaneous exposure, the degree to which risk is overestimated cannot be quantified.

4.3.2.2 Related to Exposure for Terrestrial Species

Screening-level risk assessments for applications of pesticides consider dietary exposure alone. Other routes of exposure, not considered in this assessment, are discussed below:

Incidental soil ingestion exposure - This risk assessment does not consider incidental soil ingestion. Available data suggests that up to 15% of the diet can consist of incidentally ingested soil depending on the species and feeding strategy (Beyer et al., 1994). Being that the proposed new use is a granular formulation, significant exposure via this scenario is not expected.

Inhalation Exposure - The screening risk assessment does not consider inhalation exposure. Such exposure may occur through three potential sources: (1) spray material in droplet form at the time of application (2) vapor phase pesticide volatilizing from treated surfaces, and (3) airborne particulate (soil, vegetative material, and pesticide dusts). Being that the proposed new use is a granular formulation, significant inhalation exposure is not expected.

Dermal Exposure - The screening assessment does not consider dermal exposure, except as it is indirectly included in calculations of RQ's based on lethal doses per unit of pesticide treated area. Dermal exposure may occur through three potential sources: (1) direct application of spray to terrestrial wildlife in the treated area or within the drift footprint, (2) incidental contact with contaminated vegetation, or (3) contact with contaminated water or soil. Being that the proposed new use is a use is a granular formulation, significant exposure via these scenarios is not expected.

Drinking Water Exposure - Drinking water exposure to a pesticide active ingredient may be the result of consumption of surface water or consumption of the pesticide in dew or other water on the surfaces of treated vegetation. For pesticide active ingredients with a potential to dissolve in runoff, puddles on the treated field may contain the chemical.

4.3.3 Related to Effects Assessment

4.3.3.1 Age class and sensitivity of effects thresholds

It is generally recognized that test organism age may have a significant impact on the observed sensitivity to a toxicant. The screening risk assessment acute toxicity data for fish are collected on juvenile fish between 0.1 and 5 grams. Aquatic invertebrate acute testing is performed on recommended immature age classes (e.g., first instar for daphnids, second instar for amphipods, stoneflies and mayflies, and third instar for midges). Similarly, acute dietary testing with birds is also performed on juveniles, with mallard being 5-10 days old and quail 10-14 days old.

Testing of juveniles may overestimate toxicity at older age classes for active ingredients, such as abamectin, that act directly (without metabolic transformation) because younger age classes may not have the enzymatic systems associated with detoxifying xenobiotics. The screening risk assessment has no current provisions for a generally applied method that accounts for this uncertainty. Insofar as the available toxicity data may provide ranges of sensitivity information with respect to age class, the risk assessment uses the most sensitive life-stage information as the conservative screening endpoint.

4.3.3.2 *Aquatic Studies Conducted Above Water Solubility*

A number of the acute toxicity tests, primarily for fish, oyster and aquatic plants, were conducted as nominal and were above the known solubility limit for abamectin (<1.0 µg/L in tap water). Therefore, the dissolved bioavailable form in these toxicity tests is unknown. Risk quotients calculated from these values may underestimate risks.

4.3.3.3 *Lack of Effect Studies and Complete Review of Aquatic Plant Data*

There are no chronic toxicity data available for the Agency to assess chronic risk of abamectin to marine and estuarine fish. There is also no registered submitted data for vegetative vigor and seedling emergence toxicity data for terrestrial plants. An acute oral toxicity study with a passerine bird species and a chronic reproduction study with the bobwhite quail are also not available. Toxicity tests with sediment organisms are also not available, and the potential for abamectin to be present in the sediment exists. There are only two of the five studies addressing the acute toxicity of abamectin to aquatic plants available.

4.3.3.4 *Uncertainty in LD₅₀ for Mallards and NOAEC for Chronic Daphnia Study*

The acute oral LD₅₀ for mallard ducks (*Anas platyrhynchos*) is 85 mg ai/kg-bw (MRID 00097859, moderately toxic). However, regurgitation was observed in all the mallard duck acute oral treatment groups, therefore, the reported acute oral LD₅₀ might be underestimating toxicity.

The life-cycle toxicity test with the *Daphnia magna* resulted in a reproductive NOAEC of 0.030 µg ai/L which was the lowest concentration tested, but the adults in the two lowest treatment groups were observed to be pale and smaller compared to the controls (MRID 00153570). Therefore, the reproductive NOAEC appears to underestimate the true no effect concentration for *Daphnia* from chronic exposure to abamectin, as the NOAEC appears to be lower than 0.030 µg ai/L which may be underestimating risk.

4.3.3.5 *Use of the Most Sensitive Species Tested*

Although the screening risk assessment relies on a selected toxicity endpoint from the most sensitive species tested, it does not necessarily mean that the selected toxicity endpoints reflect sensitivity of the most sensitive species existing in a given environment. The relative position of the most sensitive species tested in the distribution of all possible species is a function of the overall variability among species to a particular chemical. In the case of listed species, there is uncertainty regarding the relationship of the listed species' sensitivity and the most sensitive species tested.

5.0 Literature Cited

- Fletcher, J., J. Nellessen and T Pfleeger. 1994. Literature review and evaluation of the EPA Food-Chain (Kenaga) Nomogram, an Instrument for Estimating Pesticide Residues on Plants. *Environ. Tox. Chem.* 13(9): 1383-1391.
- Hoerger, F. and E. E. Kenaga. 1972. Pesticide residues on plants: correlation of representative data as a basis for estimation of their magnitude in the environment. in: F. Coulston and F. Korte (editors), *Environmental Quality and Safety: Chemistry, Toxicology, and Technology*. Vol I. Georg Thieme Publishers, Stuttgart, West Germany, pp. 9-28.
- Pfleeger, T.G., A. Fong, R. Hayes, H. Ratsch and C. Wickliff. 1996. Field evaluation of the EPA (Kenaga) nomogram, a method for estimating wildlife exposure to pesticide residues on plants. *Env. Toxicol. Chem.* 15:535-543, 1996.
- Sherma, J. and Cairns, T. Comprehensive analytical profiles of important pesticides. 1993. CRC Press, Inc. Boca Raton, Fl. p 75.
- Tomlin, C.D.S. (ed.). 1994. *The Pesticide Manual - World Compendium*. 10th ed. Surrey, UK. The British Crop Protection Council, p. 4:
- U.S. Environmental Protection Agency. 1998. Guidelines for Ecological Risk Assessment. Risk Assessment Forum, Office of Research and Development, Washington, D.C. EPA/630/R-95/002F. April 1998.
- U.S. Environmental Protection Agency. 2004. Overview of the Ecological Risk Assessment Process in the Office of Pesticide Programs, U.S. Environmental Protection Agency. Endangered and Threatened Species Effects Determinations. Office of Prevention, Pesticides and Toxic Substances, Office of Pesticide Programs, Washington, D.C. January 23, 2004.
- U.S. Environmental Protection Agency. 2004a. Individual Effect Chance Model. Version 1.1, Developed by Ed Odenkirchen, Environmental Effects and Fate Division, Office of Pesticides. June 22, 2004.

Appendix A. EIIS Incident Reports

Incident No.	Year	State	Organism Affected	No. Acres/Animal Affected	Mixture – if mixture; abamectin plus names of others chemicals	Certainty index	Comments
I007644-001	June 1998	CA	Almonds	All 65	Agri-Mek (EPA# 100-898) abamectin	possible	Almond field treated directly w/Agri-Mek. Type of injury not reported.
I007644-002	June 1998	CA	Almonds	All 34	Agri-Mek (EPA# 100-898) abamectin	possible	Almond field treated directly w/Agri-Mek. Type of injury not reported.
I007644-003	June 1998	CA	Almonds	All 106	Agri-Mek (EPA# 100-898) abamectin	possible	Almond field treated directly w/Agri-Mek. Type of injury not reported.
I008611-001	April 1999	CA	Bees	100 colonies	Agri-Mek (EPA# 100-898) abamectin	probable	<p>Section 18 exemption for avocados for thrip problem. Southern California beekeepers reported bee kills where beehives kept in avocado groves.</p> <p>Report indicates that contrary to recommendation helicopters have been spraying during the day instead of at night as County instructions favored; also the labels warn of drift if bees are visiting crops. Report indicated that thousands of dead bees littered the bee yard. The County sent a representative to take samples.</p>
I010221-001	April 2000	TX	Catfish	100 dead (1/8 acre pond)	PT 370 Ascend Fire Ant Stopper (EPA# 499-370) abamectin; Award (EPA#100-722) fenoxycarb	probable	1/8 lb of both Ascend and Award to applied to areas around pond. 1 to 1 ½ in. of rain fell the next day. 100 catfish of varying sizes and age died 2 days after application. No other species in pond observed dead. Pond located in woods w/little to no runoff or stream flow, and is filled w/well water.
I-10837-019	June 2000	CA	Grapes	All 34	Agri-Mek (EPA# 100-898) abamectin	possible	Applied at 10 gal/A directly to foliar crop by airblast (broadcast). Type of injury not reported.

							Registrant inspector in responding to question "Application within Label" stated "Questionable".
I014237-001	June 2003	FL	Bait Fish (small)	"tons"	Agri-Mek 0.15 (EPA# 100-898) abamectin	probable	Agri-Mek applied to citrus grove less than 25 ft from lake at a reported rate of 10 oz. Application made in morning and rain fell in afternoon. One week after application, "tons" of dead small bait fish observed around edges of lake.

Appendix B. PRZM/EXAMS Output Files

Almonds & Walnuts

stored as A49nd.out

Chemical: Abamectin

PRZM environment:

CAalmond_WirrigSTD.txt

modified Tuesday, 26 August 2008 at 05:16:36

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w23232.dvf

modified Tuesday, 26 August 2008 at 05:15:38

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.04042	0.03762	0.02937	0.01716	0.01382	0.007031
1962	0.1863	0.1721	0.1305	0.09103	0.0762	0.0358
1963	0.0782	0.07558	0.06845	0.06027	0.05784	0.04969
1964	0.06302	0.06128	0.05414	0.04567	0.0432	0.03916
1965	0.05042	0.04893	0.04575	0.03988	0.03742	0.03374
1966	0.04619	0.04444	0.04009	0.03475	0.03128	0.02794
1967	0.05962	0.05763	0.05198	0.04225	0.03895	0.03318
1968	0.04064	0.03904	0.03608	0.03243	0.03128	0.0257
1969	0.04048	0.03924	0.03552	0.03225	0.0311	0.02682
1970	0.07429	0.0704	0.05852	0.04059	0.03467	0.02976
1971	0.04454	0.04288	0.03889	0.03565	0.03428	0.0292
1972	0.04066	0.03907	0.03449	0.03069	0.02964	0.02529
1973	0.04234	0.04112	0.03759	0.03409	0.03264	0.02961
1974	0.04055	0.03859	0.036	0.03547	0.03502	0.02774
1975	0.03886	0.03724	0.03331	0.03013	0.02918	0.02381
1976	0.03948	0.03754	0.03411	0.02954	0.02858	0.02099
1977	0.03813	0.03658	0.03265	0.02888	0.02769	0.02172
1978	0.05851	0.05568	0.04864	0.04252	0.03918	0.03234
1979	0.04474	0.04354	0.03824	0.035	0.03424	0.02998
1980	0.04284	0.04167	0.03884	0.03696	0.03562	0.03011
1981	0.06692	0.06301	0.05304	0.04275	0.03634	0.02852
1982	0.07453	0.071	0.0596	0.04866	0.04527	0.041
1983	0.05544	0.05408	0.04511	0.04347	0.0423	0.03868
1984	0.04931	0.04785	0.04341	0.0384	0.0362	0.03178
1985	0.04294	0.04086	0.03631	0.03017	0.02883	0.02434
1986	0.05697	0.05398	0.04572	0.0369	0.03431	0.02965
1987	0.03928	0.03727	0.03335	0.02981	0.02853	0.0232
1988	0.03674	0.03521	0.03225	0.02849	0.02733	0.02119
1989	0.04267	0.0405	0.03414	0.02889	0.02756	0.02245
1990	0.04835	0.04679	0.03988	0.03336	0.03189	0.02694

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258065	0.1863	0.1721	0.1305	0.09103	0.0762	0.04969
0.064516129	0.0782	0.07558	0.06845	0.06027	0.05784	0.041

0.096774194	0.07453	0.071	0.0596	0.04866	0.04527	0.03916
0.129032258	0.07429	0.0704	0.05852	0.04567	0.0432	0.03868
0.161290323	0.06692	0.06301	0.05414	0.04347	0.0423	0.0358
0.193548387	0.06302	0.06128	0.05304	0.04275	0.03918	0.03374
0.225806452	0.05962	0.05763	0.05198	0.04252	0.03895	0.03318
0.258064516	0.05851	0.05568	0.04864	0.04225	0.03742	0.03234
0.290322581	0.05697	0.05408	0.04575	0.04059	0.03634	0.03178
0.322580645	0.05544	0.05398	0.04572	0.03988	0.0362	0.03011
0.35483871	0.05042	0.04893	0.04511	0.0384	0.03562	0.02998
0.387096774	0.04931	0.04785	0.04341	0.03696	0.03502	0.02976
0.419354839	0.04835	0.04679	0.04009	0.0369	0.03467	0.02965
0.451612903	0.04619	0.04444	0.03988	0.03565	0.03431	0.02961
0.483870968	0.04474	0.04354	0.03889	0.03547	0.03428	0.0292
0.516129032	0.04454	0.04288	0.03884	0.035	0.03424	0.02852
0.548387097	0.04294	0.04167	0.03824	0.03475	0.03264	0.02794
0.580645161	0.04284	0.04112	0.03759	0.03409	0.03189	0.02774
0.612903226	0.04267	0.04086	0.03631	0.03336	0.03128	0.02694
0.64516129	0.04234	0.0405	0.03608	0.03243	0.03128	0.02682
0.677419355	0.04066	0.03924	0.036	0.03225	0.0311	0.0257
0.709677419	0.04064	0.03907	0.03552	0.03069	0.02964	0.02529
0.741935484	0.04055	0.03904	0.03449	0.03017	0.02918	0.02434
0.774193548	0.04048	0.03859	0.03414	0.03013	0.02883	0.02381
0.806451613	0.04042	0.03762	0.03411	0.02981	0.02858	0.0232
0.838709677	0.03948	0.03754	0.03335	0.02954	0.02853	0.02245
0.870967742	0.03928	0.03727	0.03331	0.02889	0.02769	0.02172
0.903225806	0.03886	0.03724	0.03265	0.02888	0.02756	0.02119
0.935483871	0.03813	0.03658	0.03225	0.02849	0.02733	0.02099
0.967741935	0.03674	0.03521	0.02937	0.01716	0.01382	0.007031

0.1	0.074506	0.07094	0.059492	0.048361	0.045063	0.039112
Average of yearly averages:						0.028912

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: CAAlmond

Metfile: w23232.dvf

PRZM scenario: CAAlmond_WirrigSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m^3/mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	

Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0263	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	
Application Date	Date	6-May	dd/mm or dd/mmm or dd-mm or dd-mmm	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Apples

stored as PAAppl.es.out

Chemical: Abamectin

PRZM environment:

PAappleSTD.txt

modified Tuesday, 26 August 2008 at 05:16:42

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w14751.dvf

modified Tuesday, 26 August 2008 at 05:15:00

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.1297	0.1202	0.09809	0.08633	0.07887	0.03257
1962	0.1091	0.1048	0.09207	0.08378	0.08291	0.06465
1963	0.08413	0.08276	0.08112	0.08014	0.07843	0.06935
1964	0.1102	0.1058	0.09275	0.08206	0.07893	0.07167
1965	0.08485	0.08388	0.08044	0.07455	0.07202	0.06443
1966	0.2341	0.22	0.1795	0.1396	0.1277	0.07623
1967	0.1997	0.1925	0.1709	0.1447	0.1349	0.109
1968	0.2175	0.2059	0.1717	0.1402	0.132	0.1061
1969	0.4276	0.4026	0.3431	0.2618	0.2348	0.1472
1970	0.2222	0.2152	0.1944	0.1863	0.1818	0.1615
1971	0.283	0.2684	0.2253	0.1945	0.1802	0.14
1972	0.6103	0.5716	0.4606	0.3474	0.3116	0.1998
1973	0.2601	0.2502	0.2208	0.2035	0.2009	0.1769
1974	0.212	0.2058	0.1902	0.1769	0.1688	0.1446
1975	0.3447	0.3255	0.27	0.2154	0.197	0.1434
1976	0.2086	0.2012	0.181	0.1611	0.155	0.1432

1977	0.1525	0.1522	0.1494	0.1485	0.1466	0.1193
1978	0.1635	0.1575	0.1447	0.1267	0.1195	0.1039
1979	0.1498	0.1443	0.133	0.1203	0.114	0.09969
1980	0.0978	0.09458	0.09239	0.09027	0.08997	0.07791
1981	0.1141	0.1102	0.09913	0.091	0.08601	0.07523
1982	0.1074	0.1034	0.09186	0.08491	0.08043	0.07089
1983	0.09096	0.08812	0.07998	0.06195	0.0573	0.05515
1984	0.1063	0.1024	0.09483	0.08547	0.07987	0.06588
1985	0.09061	0.08677	0.07597	0.06727	0.0667	0.05932
1986	0.1731	0.1646	0.1424	0.1172	0.1091	0.07807
1987	0.1499	0.1441	0.1288	0.1115	0.1074	0.0899
1988	0.1514	0.1451	0.1263	0.108	0.1015	0.09059
1989	0.1653	0.1591	0.1406	0.1269	0.1208	0.0931
1990	0.1436	0.1387	0.1241	0.11	0.1081	0.09543

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.6103	0.5716	0.4606	0.3474	0.3116	0.1998
0.064516	0.4276	0.4026	0.3431	0.2618	0.2348	0.1769
0.096774	0.3447	0.3255	0.27	0.2154	0.2009	0.1615
0.129032	0.283	0.2684	0.2253	0.2035	0.197	0.1472
0.16129	0.2601	0.2502	0.2208	0.1945	0.1818	0.1446
0.193548	0.2341	0.22	0.1944	0.1863	0.1802	0.1434
0.225806	0.2222	0.2152	0.1902	0.1769	0.1688	0.1432
0.258065	0.2175	0.2059	0.181	0.1611	0.155	0.14
0.290323	0.212	0.2058	0.1795	0.1485	0.1466	0.1193
0.322581	0.2086	0.2012	0.1717	0.1447	0.1349	0.109
0.354839	0.1997	0.1925	0.1709	0.1402	0.132	0.1061
0.387097	0.1731	0.1646	0.1494	0.1396	0.1277	0.1039
0.419355	0.1653	0.1591	0.1447	0.1269	0.1208	0.09969
0.451613	0.1635	0.1575	0.1424	0.1267	0.1195	0.09543
0.483871	0.1525	0.1522	0.1406	0.1203	0.114	0.0931
0.516129	0.1514	0.1451	0.133	0.1172	0.1091	0.09059
0.548387	0.1499	0.1443	0.1288	0.1115	0.1081	0.0899
0.580645	0.1498	0.1441	0.1263	0.11	0.1074	0.07807
0.612903	0.1436	0.1387	0.1241	0.108	0.1015	0.07791
0.645161	0.1297	0.1202	0.09913	0.091	0.08997	0.07623
0.677419	0.1141	0.1102	0.09809	0.09027	0.08601	0.07523
0.709677	0.1102	0.1058	0.09483	0.08633	0.08291	0.07167
0.741935	0.1091	0.1048	0.09275	0.08547	0.08043	0.07089
0.774194	0.1074	0.1034	0.09239	0.08491	0.07987	0.06935
0.806452	0.1063	0.1024	0.09207	0.08378	0.07893	0.06588
0.83871	0.0978	0.09458	0.09186	0.08206	0.07887	0.06465
0.870968	0.09096	0.08812	0.08112	0.08014	0.07843	0.06443
0.903226	0.09061	0.08677	0.08044	0.07455	0.07202	0.05932
0.935484	0.08485	0.08388	0.07998	0.06727	0.0667	0.05515
0.967742	0.08413	0.08276	0.07597	0.06195	0.0573	0.03257

0.1	0.33853	0.31979	0.26553	0.21421	0.20051	0.16007
					Average of yearly averages:	0.100832

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: PAAples

Metfile: w14751.dvf

PRZM scenario: PAApleSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0263	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	
Application Date	Date	15-06	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Avocado

stored as FLAvocado.out

Chemical: Abamectin

PRZM environment:

modified Tuesday, 26 August 2008 at 05:16:38

FLavocadoSTD.txt
EXAMS environment:
pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w12839.dvf

modified Tuesday, 26 August 2008 at 05:14:20

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.08559	0.07854	0.05815	0.04895	0.04471	0.02267
1962	0.0994	0.09215	0.07112	0.05737	0.05277	0.03224
1963	0.1122	0.1042	0.08108	0.06411	0.05753	0.03613
1964	0.1057	0.0986	0.08067	0.06656	0.06339	0.04367
1965	0.1107	0.1035	0.08255	0.06902	0.06265	0.04452
1966	0.1126	0.1055	0.08778	0.08032	0.07547	0.0513
1967	0.1333	0.1254	0.1052	0.0844	0.07945	0.05455
1968	0.119	0.1144	0.1038	0.08812	0.08123	0.05369
1969	0.1151	0.108	0.08723	0.07354	0.06721	0.0449
1970	0.1236	0.1161	0.09566	0.08259	0.07511	0.04773
1971	0.1095	0.1023	0.08199	0.06817	0.06359	0.04208
1972	0.1252	0.1173	0.09597	0.08014	0.07283	0.04585
1973	0.1104	0.1033	0.08259	0.06883	0.06229	0.04094
1974	0.1059	0.09874	0.07793	0.06439	0.06025	0.03785
1975	0.1044	0.09731	0.07669	0.06283	0.05643	0.03503
1976	0.103	0.09601	0.07565	0.06181	0.05588	0.03494
1977	0.1895	0.175	0.1344	0.1144	0.09966	0.05722
1978	0.1186	0.1115	0.09066	0.07714	0.07016	0.04766
1979	0.3626	0.3324	0.2721	0.1868	0.1573	0.08354
1980	0.1429	0.1351	0.1121	0.1031	0.09322	0.06416
1981	0.117	0.1096	0.08832	0.07509	0.06799	0.04728
1982	0.1315	0.1227	0.09684	0.08019	0.07318	0.04522
1983	0.1084	0.1012	0.08001	0.0666	0.06126	0.03958
1984	0.1108	0.1043	0.08523	0.07678	0.07005	0.04372
1985	0.1087	0.1014	0.08023	0.06697	0.06062	0.04117
1986	0.1066	0.09954	0.07983	0.06589	0.05971	0.03763
1987	0.1042	0.09704	0.07626	0.06255	0.05713	0.037
1988	0.1049	0.09785	0.0772	0.06339	0.05933	0.03724
1989	0.1037	0.09634	0.07504	0.06166	0.05524	0.03438
1990	0.109	0.1016	0.07992	0.06982	0.06428	0.03898

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.3626	0.3324	0.2721	0.1868	0.1573	0.08354
0.064516	0.1895	0.175	0.1344	0.1144	0.09966	0.06416
0.096774	0.1429	0.1351	0.1121	0.1031	0.09322	0.05722
0.129032	0.1333	0.1254	0.1052	0.08812	0.08123	0.05455
0.16129	0.1315	0.1227	0.1038	0.0844	0.07945	0.05369
0.193548	0.1252	0.1173	0.09684	0.08259	0.07547	0.0513
0.225806	0.1236	0.1161	0.09597	0.08032	0.07511	0.04773
0.258065	0.119	0.1144	0.09566	0.08019	0.07318	0.04766
0.290323	0.1186	0.1115	0.09066	0.08014	0.07283	0.04728
0.322581	0.117	0.1096	0.08832	0.07714	0.07016	0.04585
0.354839	0.1151	0.108	0.08778	0.07678	0.07005	0.04522

0.387097	0.1126	0.1055	0.08723	0.07509	0.06799	0.0449
0.419355	0.1122	0.1043	0.08523	0.07354	0.06721	0.04452
0.451613	0.1108	0.1042	0.08259	0.06982	0.06428	0.04372
0.483871	0.1107	0.1035	0.08255	0.06902	0.06359	0.04367
0.516129	0.1104	0.1033	0.08199	0.06883	0.06339	0.04208
0.548387	0.1095	0.1023	0.08108	0.06817	0.06265	0.04117
0.580645	0.109	0.1016	0.08067	0.06697	0.06229	0.04094
0.612903	0.1087	0.1014	0.08023	0.0666	0.06126	0.03958
0.645161	0.1084	0.1012	0.08001	0.06656	0.06062	0.03898
0.677419	0.1066	0.09954	0.07992	0.06589	0.06025	0.03785
0.709677	0.1059	0.09874	0.07983	0.06439	0.05971	0.03763
0.741935	0.1057	0.0986	0.07793	0.06411	0.05933	0.03724
0.774194	0.1049	0.09785	0.0772	0.06339	0.05753	0.037
0.806452	0.1044	0.09731	0.07669	0.06283	0.05713	0.03613
0.83871	0.1042	0.09704	0.07626	0.06255	0.05643	0.03503
0.870968	0.1037	0.09634	0.07565	0.06181	0.05588	0.03494
0.903226	0.103	0.09601	0.07504	0.06166	0.05524	0.03438
0.935484	0.0994	0.09215	0.07112	0.05737	0.05277	0.03224
0.967742	0.08559	0.07854	0.05815	0.04895	0.04471	0.02267
0.1	0.14194	0.13413	0.11141	0.101602	0.092021	0.056953
					Average of yearly averages:	0.044096

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: FLAvocado

Metfile: w12839.dvf

PRZM scenario: FLavocadoSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
Henry's Law Const.	henry	2.60E-08	atm-m ³ /mol	
Vapor Pressure	vapr	1.50E-09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0263	kg/ha	

Application Efficiency:	APPEFF	0.95	fraction
Spray Drift	DRFT	0.05	fraction of application rate applied to pond
Application Date	Date	4-May	dd/mm or dd/mm/mm or dd-mm or dd-mmm
Interval 1	interval	30	days Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha
Record 17:	FILTRA		
	IPSCND	1	
	UPTKF		
Record 18:	PLVKRT		
	PLDKRT		
	FEXTRC	0.5	
Flag for Index Res. Run	IR	EPA Pond	
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)

Celeriac

stored as FLCeleriac.out

Chemical: Abamectin

PRZM environment:

FLcarrotSTD.txt

modified Tuesday, 26 August 2008 at 05:16:38

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w12844.dvf

modified Tuesday, 26 August 2008 at 05:14:22

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.05137	0.04779	0.03901	0.02904	0.02987	0.01335
1962	0.2173	0.2056	0.1831	0.1507	0.1453	0.07778
1963	0.2796	0.267	0.231	0.1774	0.1589	0.1002
1964	0.3291	0.3171	0.2823	0.2533	0.2411	0.1621
1965	0.379	0.3599	0.3211	0.2751	0.2479	0.1966
1966	0.5594	0.5311	0.4475	0.3731	0.3554	0.2479
1967	0.3088	0.2977	0.2767	0.2504	0.243	0.2108
1968	0.4022	0.388	0.3507	0.2999	0.2884	0.2167
1969	0.5095	0.4777	0.4064	0.3401	0.3185	0.2442
1970	0.2504	0.243	0.2289	0.2168	0.2126	0.1928
1971	0.2942	0.2778	0.2504	0.1961	0.1837	0.147
1972	0.431	0.4132	0.3511	0.2761	0.2499	0.174
1973	0.2746	0.2622	0.2373	0.2214	0.2161	0.1698
1974	0.2463	0.239	0.2186	0.1977	0.1893	0.154
1975	0.2522	0.239	0.216	0.178	0.164	0.1359
1976	0.2291	0.2194	0.1902	0.176	0.1687	0.129
1977	0.3443	0.3279	0.2792	0.2161	0.1933	0.1369
1978	0.3669	0.344	0.2902	0.2391	0.2163	0.161
1979	0.3784	0.356	0.3187	0.2627	0.2376	0.172
1980	0.223	0.2137	0.2023	0.1816	0.1783	0.1543
1981	0.284	0.2673	0.2337	0.2159	0.2008	0.1391
1982	0.3897	0.3722	0.3228	0.2527	0.235	0.1805

1983	0.2748	0.2626	0.2301	0.2105	0.2049	0.1786
1984	0.2796	0.2673	0.2306	0.1921	0.1876	0.1649
1985	0.3228	0.3076	0.2669	0.2248	0.2173	0.16
1986	0.2509	0.2393	0.2255	0.2119	0.1982	0.1514
1987	0.2639	0.2498	0.2208	0.2046	0.192	0.1498
1988	0.4108	0.3877	0.3198	0.2826	0.2695	0.183
1989	0.1981	0.1907	0.1692	0.1531	0.1466	0.1343
1990	0.3108	0.2958	0.2648	0.2325	0.2216	0.1443

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.5594	0.5311	0.4475	0.3731	0.3554	0.2479
0.064516	0.5095	0.4777	0.4064	0.3401	0.3185	0.2442
0.096774	0.431	0.4132	0.3511	0.2999	0.2884	0.2167
0.129032	0.4108	0.388	0.3507	0.2826	0.2695	0.2108
0.16129	0.4022	0.3877	0.3228	0.2761	0.2499	0.1966
0.193548	0.3897	0.3722	0.3211	0.2751	0.2479	0.1928
0.225806	0.379	0.3599	0.3198	0.2627	0.243	0.183
0.258065	0.3784	0.356	0.3187	0.2533	0.2411	0.1805
0.290323	0.3669	0.344	0.2902	0.2527	0.2376	0.1786
0.322581	0.3443	0.3279	0.2823	0.2504	0.235	0.174
0.354839	0.3291	0.3171	0.2792	0.2391	0.2216	0.172
0.387097	0.3228	0.3076	0.2767	0.2325	0.2173	0.1698
0.419355	0.3108	0.2977	0.2669	0.2248	0.2163	0.1649
0.451613	0.3088	0.2958	0.2648	0.2214	0.2161	0.1621
0.483871	0.2942	0.2778	0.2504	0.2168	0.2126	0.161
0.516129	0.284	0.2673	0.2373	0.2161	0.2049	0.16
0.548387	0.2796	0.2673	0.2337	0.2159	0.2008	0.1543
0.580645	0.2796	0.267	0.231	0.2119	0.1982	0.154
0.612903	0.2748	0.2626	0.2306	0.2105	0.1933	0.1514
0.645161	0.2746	0.2622	0.2301	0.2046	0.192	0.1498
0.677419	0.2639	0.2498	0.2289	0.1977	0.1893	0.147
0.709677	0.2522	0.243	0.2255	0.1961	0.1876	0.1443
0.741935	0.2509	0.2393	0.2208	0.1921	0.1837	0.1391
0.774194	0.2504	0.239	0.2186	0.1816	0.1783	0.1369
0.806452	0.2463	0.239	0.216	0.178	0.1687	0.1359
0.83871	0.2291	0.2194	0.2023	0.1774	0.164	0.1343
0.870968	0.223	0.2137	0.1902	0.176	0.1589	0.129
0.903226	0.2173	0.2056	0.1831	0.1531	0.1466	0.1002
0.935484	0.1981	0.1907	0.1692	0.1507	0.1453	0.07778
0.967742	0.05137	0.04779	0.03901	0.02904	0.02987	0.01335

0.1	0.42898	0.41068	0.35106	0.29817	0.28651	0.21611
Average of yearly averages:						0.159408

Inputs generated by pe5.pl - November 2006

Data used for this run:

Output File: FLCeleriac

Metfile: w12844.dvf

PRZM scenario:	FLcarrotSTD.txt			
EXAMS environment file:	pond298.exv			
Chemical Name:	Abamectin			
	Variable			
Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	
Application Date	Date	6-May	dd/mm or dd/mmm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Citrus

stored as FLCitrustets.out

Chemical: Abamectin

PRZM environment:

FLcitrusSTD.txt

modified Tuesday, 26 August 2008 at 05:16:38

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w12844.dvf

modified Tuesday, 26 August 2008 at 05:14:22

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.09614	0.08833	0.07048	0.051	0.04599	0.02746
1962	0.1561	0.1466	0.136	0.1139	0.1088	0.07214
1963	0.2106	0.2017	0.1756	0.1406	0.1292	0.09892
1964	0.2915	0.279	0.233	0.2026	0.1897	0.1495
1965	0.2907	0.2755	0.2414	0.2213	0.2066	0.1671
1966	0.4028	0.3871	0.3319	0.2902	0.2724	0.1956
1967	0.2709	0.2588	0.2354	0.2076	0.1988	0.1662
1968	0.436	0.4091	0.3843	0.3202	0.2935	0.1994
1969	0.3951	0.3735	0.3187	0.2656	0.2516	0.2008
1970	0.333	0.3157	0.2746	0.238	0.2183	0.1696
1971	0.2409	0.2317	0.211	0.1846	0.1755	0.1375
1972	0.3593	0.3399	0.3078	0.2787	0.2689	0.1836
1973	0.2359	0.229	0.2131	0.2041	0.198	0.1624
1974	0.1917	0.1851	0.1699	0.158	0.1546	0.133
1975	0.2129	0.201	0.1778	0.1626	0.1491	0.1204
1976	0.2815	0.2641	0.2171	0.1739	0.1635	0.127
1977	0.3038	0.2878	0.2526	0.2279	0.205	0.1525
1978	0.3213	0.304	0.2595	0.2298	0.2119	0.1587
1979	0.297	0.2804	0.2368	0.2014	0.1863	0.1568
1980	0.2446	0.2322	0.2149	0.1865	0.1759	0.1436
1981	0.2212	0.2108	0.1883	0.1745	0.1641	0.1304
1982	0.3421	0.3247	0.2861	0.2271	0.2088	0.1571
1983	0.28	0.269	0.2576	0.2125	0.1976	0.1562
1984	0.2755	0.2598	0.2214	0.1999	0.1883	0.1488
1985	0.2327	0.2248	0.2001	0.1783	0.1697	0.1373
1986	0.1996	0.1914	0.1726	0.1678	0.1591	0.1242
1987	0.197	0.1882	0.1671	0.1546	0.1502	0.1258
1988	0.3873	0.3627	0.2953	0.2684	0.2542	0.1708
1989	0.1844	0.1766	0.1597	0.1498	0.147	0.1264
1990	0.249	0.2351	0.2165	0.1896	0.1801	0.1298

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.436	0.4091	0.3843	0.3202	0.2935	0.2008
0.064516	0.4028	0.3871	0.3319	0.2902	0.2724	0.1994
0.096774	0.3951	0.3735	0.3187	0.2787	0.2689	0.1956
0.129032	0.3873	0.3627	0.3078	0.2684	0.2542	0.1836
0.16129	0.3593	0.3399	0.2953	0.2656	0.2516	0.1708
0.193548	0.3421	0.3247	0.2861	0.238	0.2183	0.1696
0.225806	0.333	0.3157	0.2746	0.2298	0.2119	0.1671
0.258065	0.3213	0.304	0.2595	0.2279	0.2088	0.1662
0.290323	0.3038	0.2878	0.2576	0.2271	0.2066	0.1624
0.322581	0.297	0.2804	0.2526	0.2213	0.205	0.1587
0.354839	0.2915	0.279	0.2414	0.2125	0.1988	0.1571
0.387097	0.2907	0.2755	0.2368	0.2076	0.198	0.1568
0.419355	0.2815	0.269	0.2354	0.2041	0.1976	0.1562
0.451613	0.28	0.2641	0.233	0.2026	0.1897	0.1525
0.483871	0.2755	0.2598	0.2214	0.2014	0.1883	0.1495

0.516129	0.2709	0.2588	0.2171	0.1999	0.1863	0.1488
0.548387	0.249	0.2351	0.2165	0.1896	0.1801	0.1436
0.580645	0.2446	0.2322	0.2149	0.1865	0.1759	0.1375
0.612903	0.2409	0.2317	0.2131	0.1846	0.1755	0.1373
0.645161	0.2359	0.229	0.211	0.1783	0.1697	0.133
0.677419	0.2327	0.2248	0.2001	0.1745	0.1641	0.1304
0.709677	0.2212	0.2108	0.1883	0.1739	0.1635	0.1298
0.741935	0.2129	0.2017	0.1778	0.1678	0.1591	0.127
0.774194	0.2106	0.201	0.1756	0.1626	0.1546	0.1264
0.806452	0.1996	0.1914	0.1726	0.158	0.1502	0.1258
0.83871	0.197	0.1882	0.1699	0.1546	0.1491	0.1242
0.870968	0.1917	0.1851	0.1671	0.1498	0.147	0.1204
0.903226	0.1844	0.1766	0.1597	0.1406	0.1292	0.09892
0.935484	0.1561	0.1466	0.136	0.1139	0.1088	0.07214
0.967742	0.09614	0.08833	0.07048	0.051	0.04599	0.02746
0.1	0.39432	0.37242	0.31761	0.27767	0.26743	0.1944
					Average of yearly averages:	0.144301

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: FLCitrustets

Metfile: w12844.dvf

PRZM scenario: FLCitrusSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0263	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	30-04	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	30	days	Set to 0 or delete line for single app.

app. rate 1	apprate	kg/ha
Record 17:	FILTRA	
	IPSCND	1
	UPTKF	
Record 18:	PLVKRT	
	PLDKRT	
	FEXTRC	0.5
Flag for Index Res. Run	IR	EPA Pond
Flag for runoff calc.	RUNOFF	none none, monthly or total(average of entire run)

Cotton

stored as MSCotton.out

Chemical: Abamectin

PRZM environment:

MSCottonSTD.txt

modified Tuesday, 26 August 2008 at 05:16:40

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w03940.dvf

modified Tuesday, 26 August 2008 at 05:14:14

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.1998	0.19	0.1671	0.1376	0.1086	0.03978
1962	0.1758	0.1692	0.1497	0.1387	0.134	0.1156
1963	0.1838	0.1753	0.1578	0.1239	0.1183	0.1054
1964	0.4204	0.3953	0.3215	0.2588	0.2533	0.1659
1965	0.5373	0.5064	0.4192	0.3295	0.2991	0.2168
1966	0.3168	0.3085	0.2859	0.2529	0.2399	0.2125
1967	0.2961	0.2832	0.252	0.2162	0.2059	0.1755
1968	0.2596	0.2484	0.223	0.1882	0.1801	0.1636
1969	0.3521	0.3316	0.2718	0.2139	0.1964	0.1538
1970	0.3415	0.3281	0.3046	0.2734	0.2619	0.1885
1971	0.3651	0.3478	0.2969	0.2444	0.241	0.2033
1972	0.2499	0.2455	0.2385	0.2236	0.2159	0.1823
1973	0.2144	0.2072	0.1965	0.1834	0.1758	0.1614
1974	0.2794	0.2676	0.2429	0.2254	0.2132	0.1777
1975	0.4144	0.3952	0.3396	0.2907	0.275	0.2099
1976	0.3367	0.3272	0.3099	0.2722	0.2561	0.2161
1977	0.298	0.2886	0.2598	0.232	0.2236	0.1935
1978	0.2634	0.2538	0.2254	0.2079	0.2009	0.1776
1979	0.4249	0.4077	0.356	0.3253	0.3096	0.2515
1980	0.3067	0.3021	0.2947	0.2807	0.2744	0.23
1981	0.2521	0.2435	0.2176	0.1957	0.1949	0.1686
1982	0.4053	0.3878	0.349	0.2909	0.2865	0.2018
1983	0.3336	0.3247	0.3006	0.2845	0.2729	0.2413
1984	0.3439	0.3343	0.291	0.2662	0.2441	0.2085
1985	0.3483	0.3306	0.2867	0.24	0.236	0.1986
1986	0.2496	0.2444	0.2276	0.2153	0.2057	0.1691
1987	0.2262	0.2174	0.1938	0.1906	0.1889	0.1625

1988	0.263	0.2499	0.2142	0.2037	0.1976	0.1559
1989	0.258	0.2491	0.2296	0.2078	0.2035	0.1682
1990	0.2558	0.2463	0.2288	0.218	0.209	0.1818

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.5373	0.5064	0.4192	0.3295	0.3096	0.2515
0.064516	0.4249	0.4077	0.356	0.3253	0.2991	0.2413
0.096774	0.4204	0.3953	0.349	0.2909	0.2865	0.23
0.129032	0.4144	0.3952	0.3396	0.2907	0.275	0.2168
0.16129	0.4053	0.3878	0.3215	0.2845	0.2744	0.2161
0.193548	0.3651	0.3478	0.3099	0.2807	0.2729	0.2125
0.225806	0.3521	0.3343	0.3046	0.2734	0.2619	0.2099
0.258065	0.3483	0.3316	0.3006	0.2722	0.2561	0.2085
0.290323	0.3439	0.3306	0.2969	0.2662	0.2533	0.2033
0.322581	0.3415	0.3281	0.2947	0.2588	0.2441	0.2018
0.354839	0.3367	0.3272	0.291	0.2529	0.241	0.1986
0.387097	0.3336	0.3247	0.2867	0.2444	0.2399	0.1935
0.419355	0.3168	0.3085	0.2859	0.24	0.236	0.1885
0.451613	0.3067	0.3021	0.2718	0.232	0.2236	0.1823
0.483871	0.298	0.2886	0.2598	0.2254	0.2159	0.1818
0.516129	0.2961	0.2832	0.252	0.2236	0.2132	0.1777
0.548387	0.2794	0.2676	0.2429	0.218	0.209	0.1776
0.580645	0.2634	0.2538	0.2385	0.2162	0.2059	0.1755
0.612903	0.263	0.2499	0.2296	0.2153	0.2057	0.1691
0.645161	0.2596	0.2491	0.2288	0.2139	0.2035	0.1686
0.677419	0.258	0.2484	0.2276	0.2079	0.2009	0.1682
0.709677	0.2558	0.2463	0.2254	0.2078	0.1976	0.1659
0.741935	0.2521	0.2455	0.223	0.2037	0.1964	0.1636
0.774194	0.2499	0.2444	0.2176	0.1957	0.1949	0.1625
0.806452	0.2496	0.2435	0.2142	0.1906	0.1889	0.1614
0.83871	0.2262	0.2174	0.1965	0.1882	0.1801	0.1559
0.870968	0.2144	0.2072	0.1938	0.1834	0.1758	0.1538
0.903226	0.1998	0.19	0.1671	0.1387	0.134	0.1156
0.935484	0.1838	0.1753	0.1578	0.1376	0.1183	0.1054
0.967742	0.1758	0.1692	0.1497	0.1239	0.1086	0.03978
0.1	0.4198	0.39529	0.34806	0.29088	0.28535	0.22868
Average of yearly averages:						0.179899

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: MSCotton

Metfile: w03940.dvf

PRZM scenario: MSCottonSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Variable

Description	Name	Value	Units	Comments
-------------	------	-------	-------	----------

Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0213	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	28-07	dd/mm or dd/mmm or dd-mm or dd-mmm	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Cucurbit

stored as FLCucumber.out

Chemical: Abamectin

PRZM environment:

FLCucumberSTD.txt

modified Tuesday, 26 August 2008 at 05:16:38

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w12844.dvf

modified Tuesday, 26 August 2008 at 05:14:22

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.1287	0.1185	0.09315	0.07863		0.06947 0.02037
1962	0.3477	0.329	0.2647	0.1881		0.1633 0.07786
1963	0.5249	0.4996	0.423	0.3204		0.2823 0.1396
1964	0.5414	0.5172	0.4485	0.3885		0.3656 0.2242
1965	0.6089	0.5683	0.5133	0.421		0.367 0.2428
1966	0.3609	0.347	0.3143	0.3011		0.2827 0.2519

1967	0.4399	0.4188	0.3478	0.2995	0.2718	0.1988
1968	0.6154	0.5842	0.5041	0.4006	0.3538	0.2182
1969	0.4415	0.4238	0.3885	0.3595	0.3333	0.24
1970	0.3299	0.3143	0.2985	0.256	0.2434	0.2137
1971	0.3794	0.3574	0.2967	0.2324	0.2236	0.1591
1972	0.3461	0.3276	0.2731	0.2171	0.2084	0.1736
1973	0.2903	0.2791	0.2475	0.2277	0.2072	0.1589
1974	0.3951	0.3702	0.3061	0.2441	0.2273	0.1625
1975	0.3965	0.3696	0.3308	0.2553	0.2279	0.1531
1976	0.3115	0.2967	0.27	0.2345	0.2184	0.1557
1977	0.2938	0.2784	0.2433	0.2064	0.1941	0.1594
1978	0.3434	0.3296	0.2884	0.2424	0.2305	0.1647
1979	0.4792	0.4592	0.4258	0.3488	0.3098	0.1948
1980	0.2773	0.2634	0.2329	0.2078	0.2038	0.1785
1981	0.3819	0.3611	0.2897	0.2378	0.223	0.1478
1982	0.4169	0.399	0.3547	0.2952	0.2807	0.2001
1983	0.4808	0.4501	0.3689	0.3176	0.29	0.2221
1984	0.4765	0.4508	0.3758	0.2917	0.2865	0.2132
1985	0.3875	0.3645	0.3208	0.2589	0.2329	0.1919
1986	0.2638	0.2527	0.2219	0.2009	0.1958	0.1658
1987	0.4465	0.4169	0.3594	0.321	0.2962	0.1858
1988	0.2678	0.256	0.2218	0.2111	0.1993	0.1798
1989	0.2255	0.2126	0.1841	0.1718	0.1577	0.1277
1990	0.4437	0.4113	0.3601	0.2707	0.2364	0.1378

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.6154	0.5842	0.5133	0.421	0.367	0.2519
0.064516	0.6089	0.5683	0.5041	0.4006	0.3656	0.2428
0.096774	0.5414	0.5172	0.4485	0.3885	0.3538	0.24
0.129032	0.5249	0.4996	0.4258	0.3595	0.3333	0.2242
0.16129	0.4808	0.4592	0.423	0.3488	0.3098	0.2221
0.193548	0.4792	0.4508	0.3885	0.321	0.2962	0.2182
0.225806	0.4765	0.4501	0.3758	0.3204	0.29	0.2137
0.258065	0.4465	0.4238	0.3689	0.3176	0.2865	0.2132
0.290323	0.4437	0.4188	0.3601	0.3011	0.2827	0.2001
0.322581	0.4415	0.4169	0.3594	0.2995	0.2823	0.1988
0.354839	0.4399	0.4113	0.3547	0.2952	0.2807	0.1948
0.387097	0.4169	0.399	0.3478	0.2917	0.2718	0.1919
0.419355	0.3965	0.3702	0.3308	0.2707	0.2434	0.1858
0.451613	0.3951	0.3696	0.3208	0.2589	0.2364	0.1798
0.483871	0.3875	0.3645	0.3143	0.256	0.2329	0.1785
0.516129	0.3819	0.3611	0.3061	0.2553	0.2305	0.1736
0.548387	0.3794	0.3574	0.2985	0.2441	0.2279	0.1658
0.580645	0.3609	0.347	0.2967	0.2424	0.2273	0.1647
0.612903	0.3477	0.3296	0.2897	0.2378	0.2236	0.1625
0.645161	0.3461	0.329	0.2884	0.2345	0.223	0.1594
0.677419	0.3434	0.3276	0.2731	0.2324	0.2184	0.1591
0.709677	0.3299	0.3143	0.27	0.2277	0.2084	0.1589
0.741935	0.3115	0.2967	0.2647	0.2171	0.2072	0.1557

0.774194	0.2938	0.2791	0.2475	0.2111	0.2038	0.1531
0.806452	0.2903	0.2784	0.2433	0.2078	0.1993	0.1478
0.83871	0.2773	0.2634	0.2329	0.2064	0.1958	0.1396
0.870968	0.2678	0.256	0.2219	0.2009	0.1941	0.1378
0.903226	0.2638	0.2527	0.2218	0.1881	0.1633	0.1277
0.935484	0.2255	0.2126	0.1841	0.1718	0.1577	0.07786
0.967742	0.1287	0.1185	0.09315	0.07863	0.06947	0.02037
0.1	0.53975	0.51544	0.44623	0.3856	0.35175	0.23842
					Average of yearly averages:	0.175324

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: FLCucumber

Metfile: w12844.dvf

PRZM scenario: FLCucumberSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	9-May	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			

Flag for Index Res. Run	FEXTRC	0.5
Flag for runoff calc.	IR	EPA Pond
	RUNOFF	none none, monthly or total(average of entire run)

Fruiting Vegetables

stored as FLPepper.out

Chemical: Abamectin

PRZM environment:

FLpeppersSTD.txt

modified Tuesday, 26 August 2008 at 05:16:38

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w12844.dvf

modified Tuesday, 26 August 2008 at 05:14:22

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.1199	0.1119	0.09474	0.08143	0.07086	0.03974
1962	0.2091	0.2004	0.1846	0.1583	0.1515	0.1002
1963	0.248	0.2353	0.2118	0.1673	0.1496	0.1206
1964	0.3968	0.379	0.3203	0.2981	0.2758	0.2006
1965	0.3597	0.3453	0.3023	0.2792	0.2615	0.2129
1966	0.4966	0.476	0.4102	0.3624	0.3435	0.2481
1967	0.3309	0.319	0.2936	0.2583	0.2495	0.2107
1968	0.5813	0.5485	0.5146	0.4479	0.408	0.2691
1969	0.4705	0.4469	0.3911	0.337	0.3244	0.2634
1970	0.4022	0.3843	0.3454	0.3087	0.2885	0.2229
1971	0.3851	0.3697	0.3004	0.2716	0.2547	0.1896
1972	0.474	0.4461	0.409	0.3802	0.3749	0.2467
1973	0.3125	0.2984	0.2788	0.2708	0.2689	0.2177
1974	0.2444	0.2382	0.2213	0.2068	0.1997	0.1745
1975	0.2607	0.2477	0.2236	0.2137	0.2052	0.1574
1976	0.4753	0.4447	0.3632	0.291	0.2618	0.1846
1977	0.4953	0.477	0.4203	0.3742	0.3349	0.2324
1978	0.3953	0.3765	0.3298	0.2933	0.2759	0.2177
1979	0.3406	0.3238	0.2774	0.2535	0.2394	0.198
1980	0.3891	0.3672	0.3205	0.2731	0.2572	0.1931
1981	0.2786	0.2666	0.2418	0.2198	0.2062	0.1738
1982	0.429	0.4064	0.3632	0.3003	0.2787	0.2053
1983	0.36	0.3459	0.3247	0.2766	0.2619	0.2075
1984	0.445	0.4227	0.3508	0.3044	0.2837	0.212
1985	0.3131	0.3018	0.2711	0.2404	0.2317	0.1889
1986	0.2414	0.2354	0.2235	0.212	0.2025	0.1638
1987	0.3257	0.306	0.2538	0.2188	0.2108	0.1651
1988	0.4691	0.4402	0.3627	0.3348	0.3189	0.2158
1989	0.2498	0.2392	0.2159	0.1874	0.1826	0.1554
1990	0.3257	0.3061	0.277	0.2528	0.2459	0.1705

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.5813	0.5485	0.5146	0.4479	0.408	0.2691
0.064516	0.4966	0.477	0.4203	0.3802	0.3749	0.2634
0.096774	0.4953	0.476	0.4102	0.3742	0.3435	0.2481
0.129032	0.4753	0.4469	0.409	0.3624	0.3349	0.2467
0.16129	0.474	0.4461	0.3911	0.337	0.3244	0.2324
0.193548	0.4705	0.4447	0.3632	0.3348	0.3189	0.2229
0.225806	0.4691	0.4402	0.3632	0.3087	0.2885	0.2177
0.258065	0.445	0.4227	0.3627	0.3044	0.2837	0.2177
0.290323	0.429	0.4064	0.3508	0.3003	0.2787	0.2158
0.322581	0.4022	0.3843	0.3454	0.2981	0.2759	0.2129
0.354839	0.3968	0.379	0.3298	0.2933	0.2758	0.212
0.387097	0.3953	0.3765	0.3247	0.291	0.2689	0.2107
0.419355	0.3891	0.3697	0.3205	0.2792	0.2619	0.2075
0.451613	0.3851	0.3672	0.3203	0.2766	0.2618	0.2053
0.483871	0.36	0.3459	0.3023	0.2731	0.2615	0.2006
0.516129	0.3597	0.3453	0.3004	0.2716	0.2572	0.198
0.548387	0.3406	0.3238	0.2936	0.2708	0.2547	0.1931
0.580645	0.3309	0.319	0.2788	0.2583	0.2495	0.1896
0.612903	0.3257	0.3061	0.2774	0.2535	0.2459	0.1889
0.645161	0.3257	0.306	0.277	0.2528	0.2394	0.1846
0.677419	0.3131	0.3018	0.2711	0.2404	0.2317	0.1745
0.709677	0.3125	0.2984	0.2538	0.2198	0.2108	0.1738
0.741935	0.2786	0.2666	0.2418	0.2188	0.2062	0.1705
0.774194	0.2607	0.2477	0.2236	0.2137	0.2052	0.1651
0.806452	0.2498	0.2392	0.2235	0.212	0.2025	0.1638
0.83871	0.248	0.2382	0.2213	0.2068	0.1997	0.1574
0.870968	0.2444	0.2354	0.2159	0.1874	0.1826	0.1554
0.903226	0.2414	0.2353	0.2118	0.1673	0.1515	0.1206
0.935484	0.2091	0.2004	0.1846	0.1583	0.1496	0.1002
0.967742	0.1199	0.1119	0.09474	0.08143	0.07086	0.03974
0.1	0.4933	0.47309	0.41008	0.37302	0.34264	0.24796
Average of yearly averages:						0.191935

Inputs generated by pe5.pl - November 2006

Data used for this run:

Output File: FLPepper

Metfile: w12844.dvf

PRZM scenario: FLpeppersSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
Henry's Law Const.	henry	2.60E-08	atm-m ³ /mol	
Vapor Pressure	vapr	1.50E-09	torr	

Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	28-04	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Grapes

stored as NYGrapes.out

Chemical: Abamectin

PRZM environment:

NYGrapesSTD.txt

modified Tuesday, 26 August 2008 at 05:16:42

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w14860.dvf

modified Tuesday, 26 August 2008 at 05:15:12

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.0802	0.07444	0.06301	0.04638		0.04103 0.01936
1962	0.3028	0.2853	0.2356	0.1892		0.1765 0.09046
1963	0.1978	0.19	0.1793	0.1638		0.1552 0.1381
1964	0.3999	0.381	0.3249	0.2673		0.2471 0.1727
1965	0.318	0.3059	0.2816	0.2484		0.2384 0.2059
1966	0.3005	0.29	0.2583	0.2263		0.2151 0.1949
1967	0.3468	0.3319	0.2976	0.2575		0.2433 0.1938
1968	0.2259	0.2192	0.2062	0.2019		0.1977 0.1755
1969	0.2175	0.21	0.1887	0.1743		0.1684 0.1529
1970	0.3218	0.3077	0.2791	0.24		0.2364 0.1755

1971	0.3432	0.33	0.2901	0.2535	0.2423	0.209
1972	0.2991	0.289	0.2652	0.2445	0.2337	0.2066
1973	0.2306	0.2246	0.2065	0.2019	0.198	0.1804
1974	0.3054	0.293	0.2644	0.242	0.2353	0.1882
1975	0.3639	0.3549	0.3274	0.3008	0.2907	0.2304
1976	0.3091	0.3003	0.2857	0.2713	0.2584	0.2312
1977	0.4075	0.3929	0.3721	0.3352	0.3212	0.2481
1978	0.4711	0.4531	0.4074	0.3628	0.3456	0.2901
1979	0.4767	0.4586	0.4067	0.386	0.3757	0.3103
1980	0.5089	0.4926	0.4434	0.4036	0.3896	0.3311
1981	0.3877	0.3767	0.3541	0.3454	0.3381	0.3054
1982	0.3551	0.3438	0.3117	0.2954	0.2889	0.2598
1983	0.4178	0.403	0.379	0.3358	0.321	0.2567
1984	0.3358	0.3263	0.2984	0.2763	0.2722	0.2466
1985	0.3208	0.3093	0.2773	0.2453	0.2335	0.2209
1986	0.3276	0.3155	0.2933	0.2545	0.2416	0.2137
1987	0.3822	0.3675	0.3249	0.3067	0.2936	0.2306
1988	0.3432	0.3307	0.2935	0.2713	0.263	0.2347
1989	0.2624	0.2555	0.2348	0.2278	0.2238	0.2076
1990	0.3417	0.3286	0.3055	0.2694	0.2563	0.1976

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.5089	0.4926	0.4434	0.4036	0.3896	0.3311
0.064516	0.4767	0.4586	0.4074	0.386	0.3757	0.3103
0.096774	0.4711	0.4531	0.4067	0.3628	0.3456	0.3054
0.129032	0.4178	0.403	0.379	0.3454	0.3381	0.2901
0.16129	0.4075	0.3929	0.3721	0.3358	0.3212	0.2598
0.193548	0.3999	0.381	0.3541	0.3352	0.321	0.2567
0.225806	0.3877	0.3767	0.3274	0.3067	0.2936	0.2481
0.258065	0.3822	0.3675	0.3249	0.3008	0.2907	0.2466
0.290323	0.3639	0.3549	0.3249	0.2954	0.2889	0.2347
0.322581	0.3551	0.3438	0.3117	0.2763	0.2722	0.2312
0.354839	0.3468	0.3319	0.3055	0.2713	0.263	0.2306
0.387097	0.3432	0.3307	0.2984	0.2713	0.2584	0.2304
0.419355	0.3432	0.33	0.2976	0.2694	0.2563	0.2209
0.451613	0.3417	0.3286	0.2935	0.2673	0.2471	0.2137
0.483871	0.3358	0.3263	0.2933	0.2575	0.2433	0.209
0.516129	0.3276	0.3155	0.2901	0.2545	0.2423	0.2076
0.548387	0.3218	0.3093	0.2857	0.2535	0.2416	0.2066
0.580645	0.3208	0.3077	0.2816	0.2484	0.2384	0.2059
0.612903	0.318	0.3059	0.2791	0.2453	0.2364	0.1976
0.645161	0.3091	0.3003	0.2773	0.2445	0.2353	0.1949
0.677419	0.3054	0.293	0.2652	0.242	0.2337	0.1938
0.709677	0.3028	0.29	0.2644	0.24	0.2335	0.1882
0.741935	0.3005	0.289	0.2583	0.2278	0.2238	0.1804
0.774194	0.2991	0.2853	0.2356	0.2263	0.2151	0.1755
0.806452	0.2624	0.2555	0.2348	0.2019	0.198	0.1755
0.83871	0.2306	0.2246	0.2065	0.2019	0.1977	0.1727
0.870968	0.2259	0.2192	0.2062	0.1892	0.1765	0.1529

0.903226	0.2175	0.21	0.1887	0.1743	0.1684	0.1381
0.935484	0.1978	0.19	0.1793	0.1638	0.1552	0.09046
0.967742	0.0802	0.07444	0.06301	0.04638	0.04103	0.01936

0.1	0.46577	0.44809	0.40393	0.36106	0.34485	0.30387
-----	---------	---------	---------	---------	---------	---------

Average of yearly
averages: 0.210604

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: NYGrapes

Metfile: w14860.dvf

PRZM scenario: NYGrapesSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0213	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	25-06	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Herb

stored as ORHerb.out

Chemical: Abamectin

PRZM environment:

ORMintSTD.txt

modified Tuesday, 26 August 2008 at 05:16:42

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w24232.dvf

modified Tuesday, 26 August 2008 at 05:15:54

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.02616	0.02416	0.01956	0.01466	0.01249	0.007133
1962	0.04646	0.04367	0.0399	0.03379	0.03005	0.02157
1963	0.08679	0.08177	0.07761	0.06896	0.0629	0.04317
1964	0.07416	0.07171	0.05398	0.04802	0.04544	0.0395
1965	0.06217	0.06019	0.05607	0.05099	0.04892	0.0419
1966	0.06179	0.05965	0.05498	0.04862	0.04605	0.03986
1967	0.05849	0.05644	0.05176	0.04571	0.04351	0.03636
1968	0.05648	0.05446	0.05081	0.04498	0.0438	0.03883
1969	0.06019	0.05823	0.05358	0.04767	0.04532	0.04243
1970	0.06596	0.06449	0.05943	0.0531	0.05137	0.04336
1971	0.1289	0.1223	0.1006	0.07842	0.07104	0.05501
1972	0.08285	0.07988	0.07128	0.06399	0.06082	0.05053
1973	0.07338	0.07045	0.06405	0.05633	0.04822	0.04311
1974	0.08226	0.07931	0.07484	0.065	0.06035	0.05027
1975	0.06429	0.06221	0.05752	0.05136	0.04863	0.04175
1976	0.05914	0.05701	0.05232	0.04605	0.04343	0.03463
1977	0.05038	0.04813	0.04255	0.03352	0.03102	0.02657
1978	0.05133	0.04935	0.04462	0.04146	0.03956	0.03253
1979	0.05077	0.0488	0.04416	0.03866	0.03584	0.03213
1980	0.07322	0.06961	0.05993	0.04969	0.04597	0.03747
1981	0.07239	0.06939	0.06132	0.05322	0.05027	0.04474
1982	0.07457	0.07225	0.06599	0.05812	0.05425	0.04513
1983	0.08109	0.07764	0.07287	0.06112	0.0563	0.04597
1984	0.08368	0.08	0.06892	0.05906	0.05513	0.04519
1985	0.06204	0.05984	0.05513	0.04881	0.04789	0.04017
1986	0.05576	0.05372	0.04905	0.04349	0.04117	0.03387
1987	0.0662	0.063	0.05666	0.04031	0.03736	0.03371
1988	0.05995	0.05775	0.05299	0.04661	0.04351	0.03645
1989	0.05612	0.05355	0.04638	0.0404	0.03762	0.03223
1990	0.05599	0.05395	0.04928	0.04421	0.04153	0.03625

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.1289	0.1223	0.1006	0.07842	0.07104	0.05501
0.064516	0.08679	0.08177	0.07761	0.06896	0.0629	0.05053
0.096774	0.08368	0.08	0.07484	0.065	0.06082	0.05027
0.129032	0.08285	0.07988	0.07287	0.06399	0.06035	0.04597
0.16129	0.08226	0.07931	0.07128	0.06112	0.0563	0.04519
0.193548	0.08109	0.07764	0.06892	0.05906	0.05513	0.04513
0.225806	0.07457	0.07225	0.06599	0.05812	0.05425	0.04474

0.258065	0.07416	0.07171	0.06405	0.05633	0.05137	0.04336
0.290323	0.07338	0.07045	0.06132	0.05322	0.05027	0.04317
0.322581	0.07322	0.06961	0.05993	0.0531	0.04892	0.04311
0.354839	0.07239	0.06939	0.05943	0.05136	0.04863	0.04243
0.387097	0.0662	0.06449	0.05752	0.05099	0.04822	0.0419
0.419355	0.06596	0.063	0.05666	0.04969	0.04789	0.04175
0.451613	0.06429	0.06221	0.05607	0.04881	0.04605	0.04017
0.483871	0.06217	0.06019	0.05513	0.04862	0.04597	0.03986
0.516129	0.06204	0.05984	0.05498	0.04802	0.04544	0.0395
0.548387	0.06179	0.05965	0.05398	0.04767	0.04532	0.03883
0.580645	0.06019	0.05823	0.05358	0.04661	0.0438	0.03747
0.612903	0.05995	0.05775	0.05299	0.04605	0.04351	0.03645
0.645161	0.05914	0.05701	0.05232	0.04571	0.04351	0.03636
0.677419	0.05849	0.05644	0.05176	0.04498	0.04343	0.03625
0.709677	0.05648	0.05446	0.05081	0.04421	0.04153	0.03463
0.741935	0.05612	0.05395	0.04928	0.04349	0.04117	0.03387
0.774194	0.05599	0.05372	0.04905	0.04146	0.03956	0.03371
0.806452	0.05576	0.05355	0.04638	0.0404	0.03762	0.03253
0.83871	0.05133	0.04935	0.04462	0.04031	0.03736	0.03223
0.870968	0.05077	0.0488	0.04416	0.03866	0.03584	0.03213
0.903226	0.05038	0.04813	0.04255	0.03379	0.03102	0.02657
0.935484	0.04646	0.04367	0.0399	0.03352	0.03005	0.02157
0.967742	0.02616	0.02416	0.01956	0.01466	0.01249	0.007133

0.1	0.083597	0.079988	0.074643	0.064899	0.060773	0.04984
					Average of yearly averages:	0.038394

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: ORHerb

Metfile: w24232.dvf

PRZM scenario: ORmintSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
Henry's Law Const.	henry	2.60E-08	atm-m ³ /mol	
Vapor Pressure	vapr	1.50E-09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life

Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	
Application Date	Date	25-03	dd/mm or dd/mmm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Hops

stored as ORHops.out

Chemical: Abamectin

PRZM environment:

ORhopsSTD.txt

modified Tuesday, 26 August 2008 at 05:16:42

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w24232.dvf

modified Tuesday, 26 August 2008 at 05:15:54

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.07536	0.06927	0.05153	0.03893	0.03338	0.01445
1962	0.09738	0.09131	0.07827	0.06806	0.06613	0.04389
1963	0.122	0.1157	0.09715	0.0877	0.08129	0.06832
1964	0.138	0.1343	0.1072	0.09401	0.08675	0.07821
1965	0.1368	0.1306	0.1121	0.1018	0.09604	0.087
1966	0.1394	0.1329	0.1138	0.1016	0.09761	0.08948
1967	0.136	0.1294	0.11	0.09803	0.09202	0.0866
1968	0.1385	0.1324	0.1247	0.1161	0.1151	0.09585
1969	0.1547	0.1497	0.1364	0.1304	0.1286	0.1081
1970	0.1585	0.152	0.1344	0.13	0.1254	0.1119
1971	0.1612	0.1562	0.1387	0.1276	0.124	0.1089
1972	0.1533	0.1468	0.1294	0.1208	0.1171	0.1043
1973	0.1484	0.1439	0.1337	0.1211	0.1084	0.0956
1974	0.1512	0.1447	0.1254	0.1164	0.1121	0.1025
1975	0.146	0.1398	0.1212	0.1085	0.1021	0.09596
1976	0.141	0.1348	0.1164	0.1038	0.0961	0.08499
1977	0.1215	0.1152	0.1012	0.08607	0.07887	0.0685

1978	0.1305	0.1243	0.1059	0.09427	0.08846	0.07715
1979	0.1328	0.1277	0.1127	0.1074	0.1037	0.08465
1980	0.1497	0.1448	0.1259	0.1063	0.09729	0.09261
1981	0.1542	0.151	0.1423	0.1356	0.1341	0.1058
1982	0.159	0.1526	0.1334	0.1258	0.1221	0.1089
1983	0.152	0.1455	0.1261	0.1163	0.1089	0.1009
1984	0.1405	0.1342	0.1245	0.1182	0.1085	0.09448
1985	0.1398	0.1335	0.1149	0.1026	0.0955	0.08863
1986	0.1326	0.126	0.1066	0.0947	0.08829	0.0788
1987	0.1468	0.1413	0.1308	0.1002	0.09143	0.07967
1988	0.1397	0.1332	0.114	0.1018	0.09708	0.08868
1989	0.1353	0.1298	0.1165	0.09926	0.09201	0.08371
1990	0.1425	0.1362	0.1174	0.105	0.09959	0.09224

Sorted results
Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.1612	0.1562	0.1423	0.1356	0.1341	0.1119
0.064516	0.159	0.1526	0.1387	0.1304	0.1286	0.1089
0.096774	0.1585	0.152	0.1364	0.13	0.1254	0.1089
0.129032	0.1547	0.151	0.1344	0.1276	0.124	0.1081
0.16129	0.1542	0.1497	0.1337	0.1258	0.1221	0.1058
0.193548	0.1533	0.1468	0.1334	0.1211	0.1171	0.1043
0.225806	0.152	0.1455	0.1308	0.1208	0.1151	0.1025
0.258065	0.1512	0.1448	0.1294	0.1182	0.1121	0.1009
0.290323	0.1497	0.1447	0.1261	0.1164	0.1089	0.09596
0.322581	0.1484	0.1439	0.1259	0.1163	0.1085	0.09585
0.354839	0.1468	0.1413	0.1254	0.1161	0.1084	0.0956
0.387097	0.146	0.1398	0.1247	0.1085	0.1037	0.09448
0.419355	0.1425	0.1362	0.1245	0.1074	0.1021	0.09261
0.451613	0.141	0.1348	0.1212	0.1063	0.09959	0.09224
0.483871	0.1405	0.1343	0.1174	0.105	0.09761	0.08948
0.516129	0.1398	0.1342	0.1165	0.1038	0.09729	0.08868
0.548387	0.1397	0.1335	0.1164	0.1026	0.09708	0.08863
0.580645	0.1394	0.1332	0.1149	0.1018	0.0961	0.087
0.612903	0.1385	0.1329	0.114	0.1018	0.09604	0.0866
0.645161	0.138	0.1324	0.1138	0.1016	0.0955	0.08499
0.677419	0.1368	0.1306	0.1127	0.1002	0.09202	0.08465
0.709677	0.136	0.1298	0.1121	0.09926	0.09201	0.08371
0.741935	0.1353	0.1294	0.11	0.09803	0.09143	0.07967
0.774194	0.1328	0.1277	0.1072	0.0947	0.08846	0.0788
0.806452	0.1326	0.126	0.1066	0.09427	0.08829	0.07821
0.83871	0.1305	0.1243	0.1059	0.09401	0.08675	0.07715
0.870968	0.122	0.1157	0.1012	0.0877	0.08129	0.0685
0.903226	0.1215	0.1152	0.09715	0.08607	0.07887	0.06832
0.935484	0.09738	0.09131	0.07827	0.06806	0.06613	0.04389
0.967742	0.07536	0.06927	0.05153	0.03893	0.03338	0.01445
0.1	0.15812	0.1519	0.1362	0.12976	0.12526	0.10882
Average of yearly averages:						0.087359

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: ORHops

Metfile: w24232.dvf

PRZM scenario: ORhopsSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0213	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	17-07	dd/mm or dd/mm/yy or dd-mm or dd-mm/yy	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Leafy Vegetables

stored as FLCabbage.out

Chemical: Abamectin

PRZM environment:

FLCabbageSTD.txt

EXAMS environment:

modified Tuesday, 26 August 2008 at 05:16:38

modified Tuesday, 26 August 2008 at 05:14:08

pond298.exv

Metfile: w12842.dvf

modified Tuesday, 26 August 2008 at 05:14:20

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.1176	0.1076	0.08497	0.06842		0.04618 0.01139
1962	0.1584	0.1481	0.1249	0.09757		0.08099 0.05529
1963	0.2818	0.2625	0.2268	0.1653		0.1298 0.08723
1964	0.215	0.2046	0.1812	0.1622		0.1441 0.1327
1965	0.2001	0.1896	0.1662	0.141		0.125 0.1128
1966	0.1886	0.1781	0.1547	0.127		0.1112 0.1028
1967	0.1864	0.1759	0.1524	0.127		0.1112 0.09251
1968	0.2111	0.1986	0.1723	0.1367		0.1172 0.09265
1969	0.2372	0.2283	0.1931	0.1746		0.1483 0.1169
1970	0.1989	0.1885	0.1651	0.1474		0.1462 0.1224
1971	0.2044	0.1938	0.1703	0.1415		0.1277 0.1102
1972	0.198	0.1875	0.1655	0.1419		0.1227 0.1054
1973	0.1902	0.1792	0.1574	0.1342		0.1152 0.1002
1974	0.1873	0.1764	0.1527	0.1302		0.1121 0.09744
1975	0.1893	0.1786	0.1552	0.1266		0.1121 0.09294
1976	0.1942	0.184	0.1607	0.1326		0.1156 0.09789
1977	0.173	0.1628	0.1429	0.1183		0.09916 0.08415
1978	0.1752	0.1647	0.1413	0.1152		0.09833 0.08759
1979	0.3299	0.3124	0.2526	0.1892		0.17 0.139
1980	0.2094	0.1985	0.174	0.1449		0.1249 0.1149
1981	0.1933	0.1826	0.1592	0.1363		0.12 0.1097
1982	0.2097	0.199	0.1755	0.1482		0.1337 0.1201
1983	0.2044	0.194	0.1757	0.1598		0.1386 0.1257
1984	0.1873	0.1769	0.1535	0.1277		0.1235 0.1057
1985	0.1943	0.1834	0.1599	0.1309		0.1142 0.09152
1986	0.1804	0.1698	0.1464	0.1233		0.1052 0.09182
1987	0.2162	0.2044	0.1719	0.1423		0.1281 0.1103
1988	0.2888	0.2703	0.2197	0.1787		0.148 0.1087
1989	0.1915	0.1809	0.1588	0.1391		0.1204 0.1064
1990	0.1805	0.1699	0.1465	0.1185		0.1057 0.09499

Sorted results

Prob.

	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.3299	0.3124	0.2526	0.1892		0.17 0.139
0.064516	0.2888	0.2703	0.2268	0.1787		0.1483 0.1327
0.096774	0.2818	0.2625	0.2197	0.1746		0.148 0.1257
0.129032	0.2372	0.2283	0.1931	0.1653		0.1462 0.1224
0.16129	0.2162	0.2046	0.1812	0.1622		0.1441 0.1201
0.193548	0.215	0.2044	0.1757	0.1598		0.1386 0.1169
0.225806	0.2111	0.199	0.1755	0.1482		0.1337 0.1149
0.258065	0.2097	0.1986	0.174	0.1474		0.1298 0.1128
0.290323	0.2094	0.1985	0.1723	0.1449		0.1281 0.1103
0.322581	0.2044	0.194	0.1719	0.1423		0.1277 0.1102
0.354839	0.2044	0.1938	0.1703	0.1419		0.125 0.1097
0.387097	0.2001	0.1896	0.1662	0.1415		0.1249 0.1087

0.419355	0.1989	0.1885	0.1655	0.141	0.1235	0.1064
0.451613	0.198	0.1875	0.1651	0.1391	0.1227	0.1057
0.483871	0.1943	0.184	0.1607	0.1367	0.1204	0.1054
0.516129	0.1942	0.1834	0.1599	0.1363	0.12	0.1028
0.548387	0.1933	0.1826	0.1592	0.1342	0.1172	0.1002
0.580645	0.1915	0.1809	0.1588	0.1326	0.1156	0.09789
0.612903	0.1902	0.1792	0.1574	0.1309	0.1152	0.09744
0.645161	0.1893	0.1786	0.1552	0.1302	0.1142	0.09499
0.677419	0.1886	0.1781	0.1547	0.1277	0.1121	0.09294
0.709677	0.1873	0.1769	0.1535	0.127	0.1121	0.09265
0.741935	0.1873	0.1764	0.1527	0.127	0.1112	0.09251
0.774194	0.1864	0.1759	0.1524	0.1266	0.1112	0.09182
0.806452	0.1805	0.1699	0.1465	0.1233	0.1057	0.09152
0.83871	0.1804	0.1698	0.1464	0.1185	0.1052	0.08759
0.870968	0.1752	0.1647	0.1429	0.1183	0.09916	0.08723
0.903226	0.173	0.1628	0.1413	0.1152	0.09833	0.08415
0.935484	0.1584	0.1481	0.1249	0.09757	0.08099	0.05529
0.967742	0.1176	0.1076	0.08497	0.06842	0.04618	0.01139
0.1	0.27734	0.25908	0.21704	0.17367	0.14782	0.12537
					Average of yearly averages:	0.10071

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: FLCabbage

Metfile: w12842.dvf

PRZM scenario: FLCabbageSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	

Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	11-Jan	dd/mm or dd/mmm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Mint

stored as ORMint.out

Chemical: Abamectin

PRZM environment:

ORMintSTD.txt

modified Tuesday, 26 August 2008 at 05:16:42

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w24232.dvf

modified Tuesday, 26 August 2008 at 05:15:54

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.09218	0.08504	0.06781	0.04882	0.04129	0.02091
1962	0.1176	0.1098	0.09242	0.07316	0.06431	0.04089
1963	0.1528	0.1438	0.126	0.1043	0.09359	0.06201
1964	0.1374	0.1301	0.1128	0.09251	0.08395	0.06203
1965	0.1396	0.1324	0.1155	0.09595	0.08714	0.06472
1966	0.1401	0.1326	0.1152	0.09438	0.08537	0.06353
1967	0.1383	0.131	0.1136	0.09312	0.08418	0.06103
1968	0.1361	0.1287	0.1114	0.09106	0.08252	0.06302
1969	0.1395	0.1324	0.115	0.09478	0.08597	0.06631
1970	0.1435	0.1372	0.1196	0.09894	0.08976	0.06678
1971	0.1866	0.1764	0.1461	0.116	0.105	0.07566
1972	0.1569	0.149	0.1285	0.1074	0.0983	0.07234
1973	0.1413	0.1339	0.1165	0.09582	0.08698	0.06664
1974	0.156	0.1482	0.1307	0.108	0.09799	0.072
1975	0.1427	0.1354	0.118	0.09748	0.08848	0.06574
1976	0.1389	0.1315	0.1142	0.09355	0.08466	0.06042
1977	0.1286	0.1213	0.1039	0.08391	0.07564	0.0543
1978	0.1335	0.1263	0.109	0.09042	0.0827	0.05931
1979	0.1341	0.1269	0.1096	0.08955	0.0807	0.05968
1980	0.1465	0.1403	0.1189	0.09656	0.08747	0.06294
1981	0.1403	0.1355	0.1199	0.099	0.0904	0.06799

1982	0.1442	0.1373	0.123	0.1019	0.09246	0.06791
1983	0.1539	0.1463	0.1286	0.1041	0.09402	0.06877
1984	0.1511	0.1434	0.1242	0.1033	0.09409	0.0685
1985	0.1408	0.1334	0.116	0.09542	0.08817	0.0642
1986	0.1358	0.1285	0.1112	0.0912	0.08253	0.05945
1987	0.1332	0.1259	0.1088	0.0886	0.07975	0.05886
1988	0.1385	0.1311	0.1137	0.09327	0.08455	0.06098
1989	0.1342	0.1268	0.1094	0.08893	0.08022	0.05834
1990	0.1362	0.1289	0.1116	0.09202	0.08367	0.0617

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.1866	0.1764	0.1461	0.116	0.105	0.07566
0.064516	0.1569	0.149	0.1307	0.108	0.0983	0.07234
0.096774	0.156	0.1482	0.1286	0.1074	0.09799	0.072
0.129032	0.1539	0.1463	0.1285	0.1043	0.09409	0.06877
0.16129	0.1528	0.1438	0.126	0.1041	0.09402	0.0685
0.193548	0.1511	0.1434	0.1242	0.1033	0.09359	0.06799
0.225806	0.1465	0.1403	0.123	0.1019	0.09246	0.06791
0.258065	0.1442	0.1373	0.1199	0.099	0.0904	0.06678
0.290323	0.1435	0.1372	0.1196	0.09894	0.08976	0.06664
0.322581	0.1427	0.1355	0.1189	0.09748	0.08848	0.06631
0.354839	0.1413	0.1354	0.118	0.09656	0.08817	0.06574
0.387097	0.1408	0.1339	0.1165	0.09595	0.08747	0.06472
0.419355	0.1403	0.1334	0.116	0.09582	0.08714	0.0642
0.451613	0.1401	0.1326	0.1155	0.09542	0.08698	0.06353
0.483871	0.1396	0.1324	0.1152	0.09478	0.08597	0.06302
0.516129	0.1395	0.1324	0.115	0.09438	0.08537	0.06294
0.548387	0.1389	0.1315	0.1142	0.09355	0.08466	0.06203
0.580645	0.1385	0.1311	0.1137	0.09327	0.08455	0.06201
0.612903	0.1383	0.131	0.1136	0.09312	0.08418	0.0617
0.645161	0.1374	0.1301	0.1128	0.09251	0.08395	0.06103
0.677419	0.1362	0.1289	0.1116	0.09202	0.08367	0.06098
0.709677	0.1361	0.1287	0.1114	0.0912	0.0827	0.06042
0.741935	0.1358	0.1285	0.1112	0.09106	0.08253	0.05968
0.774194	0.1342	0.1269	0.1096	0.09042	0.08252	0.05945
0.806452	0.1341	0.1268	0.1094	0.08955	0.0807	0.05931
0.83871	0.1335	0.1263	0.109	0.08893	0.08022	0.05886
0.870968	0.1332	0.1259	0.1088	0.0886	0.07975	0.05834
0.903226	0.1286	0.1213	0.1039	0.08391	0.07564	0.0543
0.935484	0.1176	0.1098	0.09242	0.07316	0.06431	0.04089
0.967742	0.09218	0.08504	0.06781	0.04882	0.04129	0.02091
0.1	0.15579	0.14801	0.12859	0.10709	0.0976	0.071677
Average of yearly averages:						0.061899

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: ORMint

Metfile:	w24232.dvf			
PRZM scenario:	ORMintSTD.txt			
EXAMS environment file:	pond298.exv			
Chemical Name:	Abamectin			
	Variable			
Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0.	cm	
Application Rate:	TAPP	0.0158	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	25-03	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Pears

stored as WAPears.out

Chemical: Abamectin

PRZM environment:

WAorchardsNMC.txt

EXAMS environment:

pond298.exv

modified Thuday, 14 June 2007 at 10:19:00

modified Tuesday, 26 August 2008 at 05:14:08

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.01885	0.01738	0.01308	0.009906	0.00844	0.004075
1962	0.02178	0.02029	0.01593	0.01281	0.01127	0.006875
1963	0.02382	0.02236	0.01797	0.01471	0.01307	0.008392
1964	0.02417	0.02267	0.01825	0.0151	0.01352	0.009127
1965	0.02713	0.02564	0.02116	0.01798	0.01624	0.01167
1966	0.02644	0.02493	0.02044	0.0173	0.01562	0.01089
1967	0.02573	0.02425	0.01983	0.01663	0.01493	0.009979
1968	0.02595	0.02445	0.01999	0.01684	0.01516	0.01068
1969	0.02664	0.02515	0.0207	0.01751	0.01578	0.01089
1970	0.02855	0.02705	0.02252	0.01934	0.01753	0.01273
1971	0.0277	0.02621	0.02171	0.01854	0.01681	0.01194
1972	0.02662	0.02512	0.02064	0.01749	0.0158	0.01103
1973	0.02617	0.02465	0.02016	0.01706	0.01539	0.0111
1974	0.03401	0.03219	0.02758	0.02252	0.02056	0.01468
1975	0.02903	0.02752	0.02301	0.01984	0.01807	0.0141
1976	0.02882	0.02733	0.02286	0.01967	0.01791	0.01288
1977	0.02723	0.0257	0.02125	0.01821	0.01652	0.01214
1978	0.0292	0.02772	0.02325	0.02005	0.01824	0.01336
1979	0.02772	0.02623	0.02175	0.01856	0.01678	0.01185
1980	0.02672	0.02521	0.02077	0.01765	0.016	0.01142
1981	0.02702	0.02554	0.02111	0.01792	0.01624	0.01142
1982	0.02686	0.02536	0.02088	0.01775	0.01605	0.01253
1983	0.02869	0.02716	0.02258	0.01947	0.01768	0.01325
1984	0.02866	0.02717	0.02272	0.01956	0.01785	0.0131
1985	0.02852	0.027	0.02247	0.01935	0.01756	0.01282
1986	0.02802	0.02652	0.02211	0.01897	0.01722	0.01282
1987	0.02826	0.02675	0.02223	0.01912	0.01732	0.01295
1988	0.03297	0.03124	0.02611	0.02204	0.0202	0.01481
1989	0.02885	0.02733	0.02281	0.01971	0.01792	0.01282
1990	0.02842	0.02694	0.02251	0.01829	0.01662	0.01391

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.03401	0.03219	0.02758	0.02252	0.02056	0.01481
0.064516	0.03297	0.03124	0.02611	0.02204	0.0202	0.01468
0.096774	0.0292	0.02772	0.02325	0.02005	0.01824	0.0141
0.129032	0.02903	0.02752	0.02301	0.01984	0.01807	0.01391
0.16129	0.02885	0.02733	0.02286	0.01971	0.01792	0.01336
0.193548	0.02882	0.02733	0.02281	0.01967	0.01791	0.01325
0.225806	0.02869	0.02717	0.02272	0.01956	0.01785	0.0131
0.258065	0.02866	0.02716	0.02258	0.01947	0.01768	0.01295
0.290323	0.02855	0.02705	0.02252	0.01935	0.01756	0.01288
0.322581	0.02852	0.027	0.02251	0.01934	0.01753	0.01282
0.354839	0.02842	0.02694	0.02247	0.01912	0.01732	0.01282
0.387097	0.02826	0.02675	0.02223	0.01897	0.01722	0.01282
0.419355	0.02802	0.02652	0.02211	0.01856	0.01681	0.01273

0.451613	0.02772	0.02623	0.02175	0.01854	0.01678	0.01253
0.483871	0.0277	0.02621	0.02171	0.01829	0.01662	0.01214
0.516129	0.02723	0.0257	0.02125	0.01821	0.01652	0.01194
0.548387	0.02713	0.02564	0.02116	0.01798	0.01624	0.01185
0.580645	0.02702	0.02554	0.02111	0.01792	0.01624	0.01167
0.612903	0.02686	0.02536	0.02088	0.01775	0.01605	0.01142
0.645161	0.02672	0.02521	0.02077	0.01765	0.016	0.01142
0.677419	0.02664	0.02515	0.0207	0.01751	0.0158	0.0111
0.709677	0.02662	0.02512	0.02064	0.01749	0.01578	0.01103
0.741935	0.02644	0.02493	0.02044	0.0173	0.01562	0.01089
0.774194	0.02617	0.02465	0.02016	0.01706	0.01539	0.01089
0.806452	0.02595	0.02445	0.01999	0.01684	0.01516	0.01068
0.83871	0.02573	0.02425	0.01983	0.01663	0.01493	0.009979
0.870968	0.02417	0.02267	0.01825	0.0151	0.01352	0.009127
0.903226	0.02382	0.02236	0.01797	0.01471	0.01307	0.008392
0.935484	0.02178	0.02029	0.01593	0.01281	0.01127	0.006875
0.967742	0.01885	0.01738	0.01308	0.009906	0.00844	0.004075

0.1	0.029183	0.0277	0.023226	0.020029	0.018223	0.014081
					Average of yearly averages:	0.011675

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: WAPears

Metfile: w24243.dvf

PRZM scenario: WAorchardsNMC.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0263	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	

Application Date	Date	31-03	dd/mm or dd/mmm or dd-mm or dd-mmm
Interval 1	interval	21	days Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha
Record 17:	FILTRA		
	IPSCND	1	
	UPTKF		
Record 18:	PLVKRT		
	PLDKRT		
	FEXTRC	0.5	
Flag for Index Res. Run	IR	EPA Pond	
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)

Plums & Prunes

stored as WAPrunestest.out

Chemical: Abamectin

PRZM environment:

WAorchardsNMC.txt

modified Thuday, 14 June 2007 at 10:19:00

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w24243.dvf

modified Tuesday, 26 August 2008 at 05:15:56

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.01818	0.01669	0.01235	0.009318	0.007918	0.002808
1962	0.02107	0.01959	0.01525	0.01219	0.01104	0.006381
1963	0.02295	0.02143	0.01701	0.01401	0.01248	0.008281
1964	0.02858	0.0273	0.01764	0.01459	0.01304	0.009267
1965	0.0275	0.02599	0.02154	0.01852	0.01688	0.0141
1966	0.02627	0.02473	0.02022	0.01726	0.01564	0.01226
1967	0.02484	0.02329	0.01877	0.01582	0.01425	0.01062
1968	0.02557	0.0241	0.02051	0.01721	0.01564	0.01177
1969	0.02574	0.0242	0.0197	0.01679	0.01525	0.01173
1970	0.03202	0.03064	0.02725	0.02123	0.01909	0.01564
1971	0.02749	0.02593	0.02136	0.01847	0.01684	0.01391
1972	0.02611	0.02457	0.02006	0.01714	0.01556	0.01211
1973	0.02538	0.02385	0.01937	0.01643	0.01489	0.01213
1974	0.02967	0.02814	0.02551	0.02074	0.019	0.01635
1975	0.03907	0.03673	0.03027	0.02365	0.02171	0.01627
1976	0.02847	0.02697	0.02254	0.01944	0.01771	0.01441
1977	0.03614	0.0341	0.02695	0.01763	0.01607	0.01345
1978	0.02962	0.02812	0.02369	0.02068	0.01903	0.01622
1979	0.02734	0.02582	0.02135	0.01833	0.01668	0.01343
1980	0.02627	0.02475	0.02028	0.0173	0.01569	0.01275
1981	0.02674	0.02519	0.02063	0.01772	0.01645	0.01305
1982	0.02689	0.02535	0.02085	0.01852	0.01685	0.0139
1983	0.02872	0.02719	0.02267	0.01966	0.01794	0.01575
1984	0.02856	0.02701	0.02247	0.01956	0.01795	0.01556
1985	0.02844	0.02692	0.02245	0.01947	0.01788	0.01514

1986	0.02769	0.02614	0.02159	0.01996	0.01916	0.01488
1987	0.04104	0.03912	0.03334	0.0203	0.01791	0.01524
1988	0.03128	0.02971	0.0251	0.02257	0.02116	0.01833
1989	0.02861	0.0271	0.02264	0.01958	0.01788	0.01476
1990	0.05318	0.04967	0.03929	0.02948	0.02635	0.01689

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.05318	0.04967	0.03929	0.02948	0.02635	0.01833
0.064516	0.04104	0.03912	0.03334	0.02365	0.02171	0.01689
0.096774	0.03907	0.03673	0.03027	0.02257	0.02116	0.01635
0.129032	0.03614	0.0341	0.02725	0.02123	0.01916	0.01627
0.16129	0.03202	0.03064	0.02695	0.02074	0.01909	0.01622
0.193548	0.03128	0.02971	0.02551	0.02068	0.01903	0.01575
0.225806	0.02967	0.02814	0.0251	0.0203	0.019	0.01564
0.258065	0.02962	0.02812	0.02369	0.01996	0.01795	0.01556
0.290323	0.02872	0.0273	0.02267	0.01966	0.01794	0.01524
0.322581	0.02861	0.02719	0.02264	0.01958	0.01791	0.01514
0.354839	0.02858	0.0271	0.02254	0.01956	0.01788	0.01488
0.387097	0.02856	0.02701	0.02247	0.01947	0.01788	0.01476
0.419355	0.02847	0.02697	0.02245	0.01944	0.01771	0.01441
0.451613	0.02844	0.02692	0.02159	0.01852	0.01688	0.0141
0.483871	0.02769	0.02614	0.02154	0.01852	0.01685	0.01391
0.516129	0.0275	0.02599	0.02136	0.01847	0.01684	0.0139
0.548387	0.02749	0.02593	0.02135	0.01833	0.01668	0.01345
0.580645	0.02734	0.02582	0.02085	0.01772	0.01645	0.01343
0.612903	0.02689	0.02535	0.02063	0.01763	0.01607	0.01305
0.645161	0.02674	0.02519	0.02051	0.0173	0.01569	0.01275
0.677419	0.02627	0.02475	0.02028	0.01726	0.01564	0.01226
0.709677	0.02627	0.02473	0.02022	0.01721	0.01564	0.01213
0.741935	0.02611	0.02457	0.02006	0.01714	0.01556	0.01211
0.774194	0.02574	0.0242	0.0197	0.01679	0.01525	0.01177
0.806452	0.02557	0.0241	0.01937	0.01643	0.01489	0.01173
0.83871	0.02538	0.02385	0.01877	0.01582	0.01425	0.01062
0.870968	0.02484	0.02329	0.01764	0.01459	0.01304	0.009267
0.903226	0.02295	0.02143	0.01701	0.01401	0.01248	0.008281
0.935484	0.02107	0.01959	0.01525	0.01219	0.01104	0.006381
0.967742	0.01818	0.01669	0.01235	0.009318	0.007918	0.002808
0.1	0.038777	0.036467	0.029968	0.022436	0.02096	0.016342
Average of yearly averages:						0.013246

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: WAPrunestest

Metfile: w24243.dvf

PRZM scenario: WAorchardsNMC.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin

Description	Variable Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
Henry's Law Const.	henry	2.60E-08	atm-m ³ /mol	
Vapor Pressure	vapr	1.50E-09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	Half-life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.0258	kg/ha	
Application Efficiency:	APPEFF	0.99	fraction	
Spray Drift	DRFT	0.01	fraction of application rate applied to pond	
Application Date	Date	25-07	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	21	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		
	UPTKF			
Record 18:	PLVKRT			
	PLDKRT			
	FEXTRC	0.5		
Flag for Index Res. Run	IR	EPA Pond		
Flag for runoff calc.	RUNOFF	none	none, monthly or total(average of entire run)	

Potato

stored as MEPotato.out

Chemical: Abamectin

PRZM environment:

MEpotatoSTD.txt

modified Tuesday, 26 August 2008 at 05:16:40

EXAMS environment:

pond298.exv

modified Tuesday, 26 August 2008 at 05:14:08

Metfile: w14607.dvf

modified Tuesday, 26 August 2008 at 05:14:52

Water segment concentrations (ppb)

Year	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
1961	0.4108	0.3921	0.3188	0.243		0.2184
1962	0.2887	0.2794	0.2595	0.2297		0.2201
1963	0.3106	0.2996	0.2768	0.2466		0.2371
1964	0.3865	0.3714	0.3337	0.2859		0.2776
1965	0.327	0.3168	0.2948	0.2652		0.2532
						0.1197
						0.1962
						0.217
						0.2371
						0.2261

1966	0.2852	0.275	0.2517	0.2372	0.2277	0.2009
1967	0.354	0.3419	0.3123	0.2762	0.2689	0.2307
1968	0.3324	0.3219	0.2985	0.2682	0.2592	0.2464
1969	0.4322	0.4176	0.3732	0.3389	0.3225	0.2822
1970	0.4817	0.4649	0.4359	0.3951	0.3742	0.318
1971	0.4112	0.3995	0.376	0.3397	0.3295	0.2922
1972	0.4887	0.474	0.4431	0.4049	0.3893	0.3233
1973	0.6691	0.6505	0.5972	0.5377	0.5105	0.4125
1974	0.5244	0.5148	0.4885	0.4542	0.4398	0.3952
1975	0.5885	0.5749	0.5191	0.4616	0.4413	0.3736
1976	0.5814	0.5632	0.51	0.4702	0.4563	0.3945
1977	0.5343	0.5257	0.51	0.4634	0.4533	0.4022
1978	0.4601	0.4484	0.4257	0.4008	0.4008	0.3715
1979	0.6269	0.6055	0.5445	0.4765	0.45	0.3822
1980	0.4589	0.4473	0.4204	0.3859	0.3809	0.3499
1981	0.5447	0.5252	0.4679	0.4122	0.3958	0.3677
1982	0.4516	0.4399	0.4159	0.3905	0.3831	0.3629
1983	0.6535	0.6358	0.5667	0.5005	0.4822	0.4003
1984	0.658	0.638	0.5788	0.5317	0.5128	0.4285
1985	0.5236	0.5093	0.4679	0.4292	0.4156	0.374
1986	0.4764	0.4636	0.4282	0.3967	0.3804	0.3505
1987	0.4108	0.4011	0.3784	0.3692	0.3606	0.3284
1988	0.3936	0.3823	0.3585	0.3281	0.3227	0.2891
1989	0.4743	0.4614	0.413	0.3686	0.3459	0.2946
1990	0.5784	0.5598	0.4952	0.454	0.434	0.3608

Sorted results

Prob.	Peak	96 hr	21 Day	60 Day	90 Day	Yearly
0.032258	0.6691	0.6505	0.5972	0.5377	0.5128	0.4285
0.064516	0.658	0.638	0.5788	0.5317	0.5105	0.4125
0.096774	0.6535	0.6358	0.5667	0.5005	0.4822	0.4022
0.129032	0.6269	0.6055	0.5445	0.4765	0.4563	0.4003
0.16129	0.5885	0.5749	0.5191	0.4702	0.4533	0.3952
0.193548	0.5814	0.5632	0.51	0.4634	0.45	0.3945
0.225806	0.5784	0.5598	0.51	0.4616	0.4413	0.3822
0.258065	0.5447	0.5257	0.4952	0.4542	0.4398	0.374
0.290323	0.5343	0.5252	0.4885	0.454	0.434	0.3736
0.322581	0.5244	0.5148	0.4679	0.4292	0.4156	0.3715
0.354839	0.5236	0.5093	0.4679	0.4122	0.4008	0.3677
0.387097	0.4887	0.474	0.4431	0.4049	0.3958	0.3629
0.419355	0.4817	0.4649	0.4359	0.4008	0.3893	0.3608
0.451613	0.4764	0.4636	0.4282	0.3967	0.3831	0.3505
0.483871	0.4743	0.4614	0.4257	0.3951	0.3809	0.3499
0.516129	0.4601	0.4484	0.4204	0.3905	0.3804	0.3284
0.548387	0.4589	0.4473	0.4159	0.3859	0.3742	0.3233
0.580645	0.4516	0.4399	0.413	0.3692	0.3606	0.318
0.612903	0.4322	0.4176	0.3784	0.3686	0.3459	0.2946
0.645161	0.4112	0.4011	0.376	0.3397	0.3295	0.2922
0.677419	0.4108	0.3995	0.3732	0.3389	0.3227	0.2891
0.709677	0.4108	0.3921	0.3585	0.3281	0.3225	0.2822

0.741935	0.3936	0.3823	0.3337	0.2859	0.2776	0.2464
0.774194	0.3865	0.3714	0.3188	0.2762	0.2689	0.2371
0.806452	0.354	0.3419	0.3123	0.2682	0.2592	0.2307
0.83871	0.3324	0.3219	0.2985	0.2652	0.2532	0.2261
0.870968	0.327	0.3168	0.2948	0.2466	0.2371	0.217
0.903226	0.3106	0.2996	0.2768	0.243	0.2277	0.2009
0.935484	0.2887	0.2794	0.2595	0.2372	0.2201	0.1962
0.967742	0.2852	0.275	0.2517	0.2297	0.2184	0.1197
0.1	0.65084	0.63277	0.56448	0.4981	0.47961	0.40201
Average of yearly averages:						0.317607

Inputs generated by pe5.pl - Novemeber 2006

Data used for this run:

Output File: MEPotato

Metfile: w14607.dvf

PRZM scenario: MEpotatoSTD.txt

EXAMS environment file: pond298.exv

Chemical Name: Abamectin
Variable

Description	Name	Value	Units	Comments
Molecular weight	mwt	873.11	g/mol	
		2.60E-		
Henry's Law Const.	henry	08	atm-m ³ /mol	
		1.50E-		
Vapor Pressure	vapr	09	torr	
Solubility	sol	78	mg/L	
Kd	Kd	82	mg/L	
Koc	Koc		mg/L	
Photolysis half-life	kdp	0.5	days	Half-life
Aerobic Aquatic Metabolism	kbacw	300	days	Halfife
Anaerobic Aquatic Metabolism	kbacs	0	days	Halfife
Aerobic Soil Metabolism	asm	150	days	Halfife
Hydrolysis:	pH 7	0	days	life
Method:	CAM	2	integer	See PRZM manual
Incorporation Depth:	DEPI	0	cm	
Application Rate:	TAPP	0.021	kg/ha	
Application Efficiency:	APPEFF	0.95	fraction	
Spray Drift	DRFT	0.05	fraction of application rate applied to pond	
Application Date	Date	28-04	dd/mm or dd/mm or dd-mm or dd-mmm	
Interval 1	interval	7	days	Set to 0 or delete line for single app.
app. rate 1	apprate		kg/ha	
Interval 2	interval	7	days	Set to 0 or delete line for single app.
app. rate 2	apprate		kg/ha	
Record 17:	FILTRA			
	IPSCND	1		

Record 18:

UPTKF
PLVKRT
PLDKRT
FEXTRC

0.5

Flag for Index Res. Run
Flag for runoff calc.

IR
RUNOFF

EPA Pond

none none, monthly or total(average of entire run)

Appendix C. T-REX Outputs

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use:	potato, cucumber, fruit veg, herb, leafy veg, potato
Formulation:	agri-mek SC
Application Rate:	0.0187 lbs a.i./acre
Half-life:	35 days
Application Interval:	7 days
Maximum # Apps./Year:	3
Length of Simulation:	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is used in both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables below <0.01 in your assessment. This is due to rounding issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	11.80
Tall Grass	5.41
Broadleaf plants/sm Insects	6.64
Fruits/pods/seeds/lg Insects	0.74

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores (grams)		
Short Grass	13.43	7.66	3.43	20	100	1000
Tall Grass	6.16	3.51	1.57			
Broadleaf plants/sm Insects	7.56	4.31	1.93			
Fruits/pods/seeds/lg Insects	0.84	0.48	0.21	0.19	0.11	0.05

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.30	0.14	0.04
Tall Grass	0.14	0.06	0.02
Broadleaf plants/sm Insects	0.17	0.08	0.02
Fruits/pods/seeds/lg Insects	0.02	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.03	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.02	#DIV/0!
Fruits/pods/seeds/lg Insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	11.25	7.77	1.80			
Tall Grass	5.15	3.56	0.83			
Broadleaf plants/sm insects	6.33	4.37	1.01			
Fruits/pods/seeds/lg insects	0.70	0.49	0.11	0.16	0.11	0.03

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
	0.38	42.64	0.32	36.43	0.17	19.53
Short Grass	0.17	19.55	0.15	16.70	0.08	8.95
Tall Grass	0.21	23.99	0.18	20.49	0.10	10.98
Broadleaf plants/sm insects	0.02	2.67	0.02	2.28	0.01	1.22
Fruits/pods/lg insects	0.01	0.59	0.00	0.51	0.00	0.27
Seeds (granivore)						

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
	#DIV/0!	4.92
Short Grass	#DIV/0!	2.25
Tall Grass	#DIV/0!	2.76
Broadleaf plants/sm insects	#DIV/0!	0.31
Fruits/pods/seeds/lg insects		

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	almonds, walnuts, pears, plums, prunes, apples
Formulation	agri-mek SC
Application Rate	0.0235 lbs a.i./acre
Half-life	35 days
Application Interval	21 days
Maximum # Apps./Year	2
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables belo <0.01 in your assessment. This is due to rou figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck)	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40
Dietary-based EECs (ppm)			
		Kenaga Values	
Short Grass		9.36	
Tall Grass		4.29	
Broadleaf plants/sm insects		5.27	
Fruits/pods/seeds/lg insects		0.59	

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores(grams)		
				20	100	1000
Short Grass	10.66	6.08	2.72			
Tall Grass	4.89	2.79	1.25			
Broadleaf plants/sm insects	6.00	3.42	1.53			
Fruits/pods/seeds/lg insects	0.67	0.38	0.17	0.15	0.08	0.04

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.24	0.11	0.03
Tall Grass	0.11	0.05	0.02
Broadleaf plants/sm insects	0.14	0.06	0.02
Fruits/pods/seeds/lg insects	0.02	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	8.93	6.17	1.43			
Tall Grass	4.09	2.83	0.66			
Broadleaf plants/sm insects	5.02	3.47	0.80			
Fruits/pods/seeds/lg insects	0.56	0.39	0.09	0.12	0.09	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.30	33.84	0.26	28.91	0.14	15.49
Tall Grass	0.14	15.51	0.12	13.25	0.06	7.10
Broadleaf plants/sm insects	0.17	19.04	0.14	16.26	0.08	8.72
Fruits/pods/lg insects	0.02	2.12	0.02	1.81	0.01	0.97
Seeds (granivore)	0.00	0.47	0.00	0.40	0.00	0.22

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.90
Tall Grass	#DIV/0!	1.79
Broadleaf plants/sm insects	#DIV/0!	2.19
Fruits/pods/seeds/lg insects	#DIV/0!	0.24

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	almonds, walnuts, pears, plums, prunes
Formulation	agri-mek SC
Application Rate	0.023 lbs a.i./acre
Half-life	35 days
Application Interval	21 days
Maximum # Apps./Year	2
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Bound Kenaga Residues.

The maximum single day residue estimation is used both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables below <0.01 in your assessment. This is due to rounding issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40
Dietary-based EECs (ppm)		Kenaga Values	
Short Grass		9.16	
Tall Grass		4.20	
Broadleaf plants/sm Insects		5.15	
Fruits/pods/seeds/lg insects		0.57	

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body-wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores (grams)		
Short Grass	10.43	5.95	2.66			
Tall Grass	4.78	2.73	1.22			
Broadleaf plants/sm Insects	5.87	3.35	1.50			
Fruits/pods/seeds/lg insects	0.65	0.37	0.17	0.14	0.08	0.04

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.24	0.11	0.03
Tall Grass	0.11	0.05	0.02
Broadleaf plants/sm insects	0.13	0.06	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bwt/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Grainvores (grams)		
	15	35	1000	15	35	1000
Short Grass	8.74	6.04	1.40			
Tall Grass	4.00	2.77	0.64			
Broadleaf plants/sm insects	4.91	3.40	0.79			
Fruits/pods/seeds/lg insects	0.55	0.38	0.09	0.12	0.08	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
	0.29	33.12	0.25	28.29	0.13	15.17
Short Grass	0.13	15.18	0.11	12.97	0.06	6.95
Tall Grass	0.16	18.63	0.14	15.91	0.08	8.53
Broadleaf plants/sm insects	0.02	2.07	0.02	1.77	0.01	0.95
Fruits/pods/lg insects	0.00	0.46	0.00	0.39	0.00	0.21
Seeds (granivore)						

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
	#DIV/0!	3.82
Short, Grass	#DIV/0!	1.75
Tall Grass	#DIV/0!	2.15
Broadleaf plants/sm insects	#DIV/0!	0.24
Fruits/pods/seeds/lg insects		

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	avocados,citrus
Formulation	agri-mek SC
Application Rate	0.0235 lbs a.i./acre
Half-life	35 days
Application Interval	30 days
Maximum # Apps./Year	2
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables belo <0.01 in your assessment. This is due to rou figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck)	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL(mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	8.75
Tall Grass	4.01
Broadleaf plants/sm Insects	4.92
Fruits/pods/seeds/lg insects	0.55

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores(grams)		
Short Grass	9.97	5.68	2.55			
Tall Grass	4.57	2.61	1.17			
Broadleaf plants/sm Insects	5.61	3.20	1.43			
Fruits/pods/seeds/lg insects	0.62	0.36	0.16	0.14	0.08	0.04

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.23	0.10	0.03
Tall Grass	0.10	0.05	0.01
Broadleaf plants/sm insects	0.13	0.06	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	8.35	5.77	1.34			
Tall Grass	3.83	2.64	0.61			
Broadleaf plants/sm insects	4.69	3.24	0.75			
Fruits/pods/seeds/lg insects	0.52	0.36	0.08	0.12	0.08	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.28	31.64	0.24	27.03	0.13	14.49
Tall Grass	0.13	14.50	0.11	12.39	0.06	6.64
Broadleaf plants/sm insects	0.16	17.80	0.13	15.20	0.07	8.15
Fruits/pods/lg insects	0.02	1.98	0.01	1.69	0.01	0.91
Seeds (granivore)	0.00	0.44	0.00	0.38	0.00	0.20

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.65
Tall Grass	#DIV/0!	1.67
Broadleaf plants/sm insects	#DIV/0!	2.05
Fruits/pods/seeds/lg insects	#DIV/0!	0.23

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	avocados, citrus
Formulation	agri-mek SC
Application Rate	0.023 lbs a.i./acre
Half-life	35 days
Application Interval	30 days
Maximum # Apps./Year	2
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables belo <0.01 in your assessment. This is due to rou figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck)	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	8.57
Tall Grass	3.93
Broadleaf plants/sm Insects	4.82
Fruits/pods/seeds/lg insects	0.54

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores (grams)		
Short Grass	9.76	5.56	2.49			
Tall Grass	4.47	2.55	1.14			
Broadleaf plants/sm Insects	5.49	3.13	1.40			
Fruits/pods/seeds/lg insects	0.61	0.35	0.16	0.14	0.08	0.03

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.22	0.10	0.03
Tall Grass	0.10	0.05	0.01
Broadleaf plants/sm insects	0.12	0.06	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bwt/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores(grams)		
	15	35	1000	15	35	1000
Short Grass	8.17	5.65	1.31			
Tall Grass	3.74	2.59	0.60			
Broadleaf plants/sm insects	4.59	3.18	0.74			
Fruits/pods/seeds/lg insects	0.51	0.35	0.08	0.11	0.08	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.27	30.97	0.23	26.46	0.13	14.18
Tall Grass	0.13	14.19	0.11	12.13	0.06	6.50
Broadleaf plants/sm insects	0.15	17.42	0.13	14.88	0.07	7.98
Fruits/pods/lg insects	0.02	1.94	0.01	1.65	0.01	0.89
Seeds (granivore)	0.00	0.43	0.00	0.37	0.00	0.20

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.57
Tall Grass	#DIV/0!	1.64
Broadleaf plants/sm insects	#DIV/0!	2.01
Fruits/pods/seeds/lg insects	#DIV/0!	0.22

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	cotton,grapes,hops
Formulation	agri-mek SC
Application Rate	0.019 lbs a.i./acre
Half-life	35 days
Application Interval	21 days
Maximum # Apps./Year	2
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables below <0.01 in your assessment. This is due to rounding issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL(mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	7.57
Tall Grass	3.47
Broadleaf plants/sm Insects	4.26
Fruits/pods/seeds/lg insects	0.47

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores(grams)		
Short Grass	8.62	4.92	2.20			
Tall Grass	3.95	2.25	1.01			
Broadleaf plants/sm Insects	4.85	2.76	1.24			
Fruits/pods/seeds/lg insects	0.54	0.31	0.14	0.12	0.07	0.03

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.20	0.09	0.03
Tall Grass	0.09	0.04	0.01
Broadleaf plants/sm insects	0.11	0.05	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bwt/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	7.22	4.99	1.16			
Tall Grass	3.31	2.29	0.53			
Broadleaf plants/sm insects	4.06	2.81	0.65			
Fruits/pods/seeds/lg insects	0.45	0.31	0.07	0.10	0.07	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.24	27.36	0.21	23.37	0.11	12.53
Tall Grass	0.11	12.54	0.09	10.71	0.05	5.74
Broadleaf plants/sm insects	0.14	15.39	0.12	13.15	0.06	7.05
Fruits/pods/lg insects	0.02	1.71	0.01	1.46	0.01	0.78
Seeds (granivore)	0.00	0.38	0.00	0.32	0.00	0.17

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.15
Tall Grass	#DIV/0!	1.45
Broadleaf plants/sm insects	#DIV/0!	1.77
Fruits/pods/seeds/lg insects	#DIV/0!	0.20

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use	mint
Formulation	agri-mek SC
Application Rate	0.014 lbs a.i./acre
Half-life	35 days
Application Interval	7 days
Maximum # Apps./Year	3
Length of Simulation	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables belo <0.01 in your assessment. This is due to rou figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	8.83
Tall Grass	4.05
Broadleaf plants/sm Insects	4.97
Fruits/pods/seeds/lg insects	0.55

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores(grams)		
				20	100	1000
Short Grass	10.06	5.74	2.57			
Tall Grass	4.61	2.63	1.18			
Broadleaf plants/sm Insects	5.66	3.23	1.44			
Fruits/pods/seeds/lg insects	0.63	0.36	0.16	0.14	0.08	0.04

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.23	0.10	0.03
Tall Grass	0.10	0.05	0.01
Broadleaf plants/sm insects	0.13	0.06	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/Insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/Insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ Insectivores (grams)			Granivores(grams)		
	15	35	1000	15	35	1000
Short Grass	8.42	5.82	1.35			
Tall Grass	3.86	2.67	0.62			
Broadleaf plants/sm insects	4.74	3.27	0.76			
Fruits/pods/seeds/lg insects	0.53	0.36	0.08	0.12	0.08	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.28	31.93	0.24	27.27	0.13	14.62
Tall Grass	0.13	14.63	0.11	12.50	0.06	6.70
Broadleaf plants/sm insects	0.16	17.96	0.14	15.34	0.07	8.22
Fruits/pods/lg insects	0.02	2.00	0.02	1.70	0.01	0.91
Seeds (granivore)	0.00	0.44	0.00	0.38	0.00	0.20

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.68
Tall Grass	#DIV/0!	1.69
Broadleaf plants/sm insects	#DIV/0!	2.07
Fruits/pods/seeds/lg insects	#DIV/0!	0.23

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use:	oleriac, cucurbit, fruit veg, herb, leafy veg, potato
Formulation:	agri-mek SC
Application Rate:	0.019 lbs a.i./acre
Half-life:	35 days
Application Interval:	7 days
Maximum # Apps./Year:	3
Length of Simulation:	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables below <0.01 in your assessment. This is due to rounding figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40
Dietary-based EECs (ppm)		Kenaga Values	
Short Grass		11.99	
Tall Grass		5.49	
Broadleaf plants/sm Insects		6.74	
Fruits/pods/seeds/lg insects		0.75	

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	56.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores (grams)		
Short Grass	13.65	7.78	3.48			
Tall Grass	6.26	3.57	1.60			
Broadleaf plants/sm Insects	7.68	4.38	1.96			
Fruits/pods/seeds/lg insects	0.85	0.49	0.22	0.19	0.11	0.05

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.31	0.14	0.04
Tall Grass	0.14	0.06	0.02
Broadleaf plants/sm insects	0.17	0.08	0.02
Fruits/pods/seeds/lg insects	0.02	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.03	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.02	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bwt/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/Insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/Insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	11.43	7.90	1.83			
Tall Grass	5.24	3.62	0.84			
Broadleaf plants/sm insects	6.43	4.44	1.03			
Fruits/pods/seeds/g insects	0.71	0.49	0.11	0.16	0.11	0.03

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.38	43.33	0.33	37.01	0.18	19.84
Tall Grass	0.18	19.86	0.15	16.96	0.08	9.09
Broadleaf plants/sm insects	0.22	24.37	0.18	20.82	0.10	11.16
Fruits/pods/g insects	0.02	2.71	0.02	2.31	0.01	1.24
Seeds (granivore)	0.01	0.60	0.00	0.51	0.00	0.28

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	4.99
Tall Grass	#DIV/0!	2.29
Broadleaf plants/sm insects	#DIV/0!	2.81
Fruits/pods/seeds/g insects	#DIV/0!	0.31

Upper Bound Kenaga Residues For RQ Calculation

Chemical Name:	abamectin
Use:	eleriac, cucurbit, fruit veg, herb, leafy veg, pota
Formulation:	agri-mek SC
Application Rate:	0.019 lbs a.i./acre
Half-life:	35 days
Application Interval:	7 days
Maximum # Apps./Year:	2
Length of Simulation:	1 year

Acute and Chronic RQs are based on the Upper Kenaga Residues.

The maximum single day residue estimation is u both the acute and reproduction RQs.

RQs reported as "0.00" in the RQ tables belo <0.01 in your assessment. This is due to rou figure issues in Excel.

Endpoints			
Avian	Mallard duck	LD50 (mg/kg-bw)	85.00
	Mallard duck)	LC50 (mg/kg-diet)	383.00
	Mallard duck	NOAEL (mg/kg-bw)	0.00
	Mallard duck	NOAEC (mg/kg-diet)	0.00
Mammals		LD50 (mg/kg-bw)	13.60
		LC50 (mg/kg-diet)	0.00
		NOAEL (mg/kg-bw)	0.12
		NOAEC (mg/kg-diet)	2.40

Dietary-based EECs (ppm)	Kenaga Values
Short Grass	8.53
Tall Grass	3.91
Broadleaf plants/sm Insects	4.80
Fruits/pods/seeds/lg insects	0.53

Avian Results

Avian Class	Body Weight (g)	Ingestion (Fdry) (g bw/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Small	20	5	23	114	2.28E-02
Mid	100	13	65	65	6.49E-02
Large	1000	58	291	29	2.91E-01
Granivores	20	5	5	25	5.06E-03
	100	13	14	14	1.44E-02
	1000	58	65	6	6.46E-02

Avian Body Weight (g)	Adjusted LD50 (mg/kg-bw)
20	44.13
100	58.18
1000	79.36

Dose-based EECs (mg/kg-bw)	Avian Classes and Body Weights (grams)					
	small 20	mid 100	large 1000	Granivores (grams)		
Short Grass	9.71	5.54	2.48	20	100	1000
Tall Grass	4.45	2.54	1.14			
Broadleaf plants/sm Insects	5.46	3.12	1.40			
Fruits/pods/seeds/lg insects	0.61	0.35	0.16	0.13	0.08	0.03

Dose-based RQs (Dose-based EEC/adjusted LD50)	Avian Acute RQs Size Class (grams)		
	20	100	1000
Short Grass	0.22	0.10	0.03
Tall Grass	0.10	0.05	0.01
Broadleaf plants/sm insects	0.12	0.06	0.02
Fruits/pods/seeds/lg insects	0.01	0.01	0.00
Seeds (granivore)	0.00	0.00	0.00

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	RQs	
	Acute	Chronic
Short Grass	0.02	#DIV/0!
Tall Grass	0.01	#DIV/0!
Broadleaf plants/sm Insects	0.01	#DIV/0!
Fruits/pods/seeds/lg insects	0.00	#DIV/0!

Mammalian Results

Mammalian Class	Body Weight	Ingestion (Fdry) (g bwt/day)	Ingestion (Fwet) (g/day)	% body wgt consumed	FI (kg-diet/day)
Herbivores/ insectivores	15	3	14	95	1.43E-02
	35	5	23	66	2.31E-02
	1000	31	153	15	1.53E-01
Grainvores	15	3	3	21	3.18E-03
	35	5	5	15	5.13E-03
	1000	31	34	3	3.40E-02

Mammalian Class	Body Weight	Adjusted LD50	Adjusted NOAEL
Herbivores/ insectivores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09
Grainvores	15	29.89	0.26
	35	24.18	0.21
	1000	10.46	0.09

Dose-Based EECs (mg/kg-bw)	Mammalian Classes and Body weight					
	Herbivores/ insectivores (grams)			Granivores (grams)		
	15	35	1000	15	35	1000
Short Grass	8.13	5.62	1.30			
Tall Grass	3.73	2.58	0.60			
Broadleaf plants/sm Insects	4.57	3.16	0.73			
Fruits/pods/seeds/lg insects	0.51	0.35	0.08	0.11	0.08	0.02

Dose-based RQs (Dose-based EEC/LD50 or NOAEL)	Small mammal 15 grams		Medium mammal 35 grams		Large mammal 1000 grams	
	Acute	Chronic	Acute	Chronic	Acute	Chronic
Short Grass	0.27	30.84	0.23	26.34	0.12	14.12
Tall Grass	0.12	14.13	0.11	12.07	0.06	6.47
Broadleaf plants/sm insects	0.15	17.34	0.13	14.82	0.07	7.94
Fruits/pods/lg insects	0.02	1.93	0.01	1.65	0.01	0.88
Seeds (granivore)	0.00	0.43	0.00	0.37	0.00	0.20

Dietary-based RQs (Dietary-based EEC/LC50 or NOAEC)	Mammal RQs	
	Acute	Chronic
Short Grass	#DIV/0!	3.55
Tall Grass	#DIV/0!	1.63
Broadleaf plants/sm insects	#DIV/0!	2.00
Fruits/pods/seeds/lg insects	#DIV/0!	0.22

Appendix D. Summary of Toxicity Data for Abamectin

Toxicity studies of technical grade abamectin with aquatic plants

Organism	% ai	Endpoint (ppb)	Source (Study Classification)
Duckweed (<i>Lemna gibba</i>), freshwater, static	91.4	14-d IC ₅₀ = 3900 (nominal, total form) ^(a) (95% CL 2300-6500) Visual Observed NOAEC = 1,200	00088787 (Supplemental)
Green algae (<i>Selenastrum capricornutum</i>), freshwater, static	91.4	9-d IC ₅₀ >100,000 (nominal, total form) ^(a, b)	00088780 (Supplemental)

^(a) Concentrations tested were above the solubility in water (7.8 ppb in distilled). Acetone was used to increase solubility in water.

^(b) Precipitate was observed at concentrations of 25,000 ppb and above.

Acute toxicity studies of technical grade abamectin with aquatic invertebrates

Organism	% ai	Endpoint (ppb)	Source (Study Classification)
Water flea (<i>Daphnia magna</i>) age <24 hr, static	91.43	48 hr EC ₅₀ = 0.34 (effect measured is immobilization as surrogate for mortality) (95% CL 0.28-0.41) slope = 10.1	00088784 (Acceptable)
Mysid (<i>Americamysis bahia</i>) age N.R., static	91	96 hr LC ₅₀ = 0.21 (95% CL 0.1-0.32)	00150565 (Acceptable)
Eastern oyster (<i>Crassostrea virginica</i>), age embryos, static	90.5	48 hr IC ₅₀ = 430 (nominal, total form) ^(a) (95% CL 280-580)	00159158 (Supplemental)
Mysid (<i>Americamysis bahia</i>) age <24 hr, flow through	Tritium labeled	96 hr LC ₅₀ = 0.020 (measured) (95% CL 0.015-0.027)	40856305 (Acceptable)
Mysid (<i>Americamysis bahia</i>) age 4 days, flow through	Tritium labeled	96 hr LC ₅₀ = 0.024 (measured)	40856305 (Acceptable)
Mysid (<i>Americamysis bahia</i>) age 10 days, flow through	Tritium labeled	96 hr LC ₅₀ = 0.032 (measured)	40856305 (Acceptable)
Mysid (<i>Americamysis bahia</i>) age 21 days, flow through	Tritium labeled	96 hr LC ₅₀ = 0.033 (measured)	40856305 (Acceptable)

^(a) Concentrations tested were above the solubility in water (7.8 ppb in distilled). Acetone was used to increase solubility in water.

Acute toxicity studies of abamectin formulations with aquatic invertebrates

Organism	Formulation % ai	Endpoint	Source (Study Classification)
Water flea (<i>D. magna</i>) age <24 hr, static	Fire Ant Bait 0.022 ^(a)	48 hr EC ₅₀ = 1.68 ppb ai (7600 ppb product) (95% CL 1.3 -2.18 ppb ai) slope = 5.0	00088785 (Supplemental)

^(a) 100 mg abamectin/100 lbs of product * 100 = 0.022% abamectin

Acute toxicity studies of abamectin degradates with aquatic invertebrates

Organism	% Purity	Endpoint	Source (Study Classification)
Water flea (<i>D. magna</i>) age <24 hr, static	Moderately polar photodegrade group 87.7%	48 hr EC ₅₀ = 6.3 (95% CL 2.5-16) slope = 1.3	ACC258746 (Acceptable)

Water flea (<i>D. magna</i>) age <24 hr, static	Polar photodegradate group 94.3%	48 hr EC ₅₀ = 4.2	ACC258746 (Acceptable)
Water flea (<i>D. magna</i>) age <24 hr, static	Non-polar photodegradate group 94.3%	48 hr EC ₅₀ = 25.9	ACC258746 (Acceptable)
Water flea (<i>D. magna</i>) age <24 hr, static	8α – hydroxy abamectin B1 (major soil metabolite)	48 hr EC ₅₀ = 25.54 (95% CL 18-32)	00153540 (Acceptable)

Acute toxicity studies of technical grade abamectin with freshwater and marine/estuarine fish

Organism	% ai	Endpoint (ppb)	Source (Study Classification)
Carp (<i>Cyprinus carpio</i>), freshwater, size 5.34 g, flow through	97	96 hr LC ₅₀ = 42 (nominal, total form) ^a (95% CL =32-56)	00153797 (Supplemental)
Rainbow trout (<i>Oncorhynchus mykiss</i>), freshwater, size 0.31 g, static	91.4	96 hr LC ₅₀ = 3.6 (nominal, total form) ^(b) (95% CL =2.2-6)	00088780 (Supplemental)
Bluegill sunfish (<i>Lepomis macrochirus</i>), freshwater size 0.34 g, static	91	96 hr LC ₅₀ = 9.6 (nominal, total form) ^(b) (95% CL =5.8-16)	00088782 (Supplemental)
Sheepshead minnow (<i>Cyprinodon variegatus</i>), estuarine/marine, size 41 mg, static renewal	91	96 hr LC ₅₀ = 15 (nominal, total form) ^(b) (95% CL =11-20)	00150910 (Supplemental)
Channel catfish (<i>Ictalurus punctatus</i>), freshwater size 0.8 g, static	91	96 hr LC ₅₀ = 24 (nominal, total form) ^(c) (95% CL =18-32)	00153588 (Supplemental)

^(a) Concentrations tested were above the solubility in water (7.8 ppb in distilled, <1 ppb in tap). No solvent was used to increase solubility in water.

^(b) Concentrations tested were above the solubility in water (7.8 ppb in distilled, < 1 ppb in tap). Acetone was used to increase solubility in water.

^(c) Concentrations tested were above the solubility in water (7.8 ppb in distilled, < 1 ppb in tap). DMF was used to increase solubility in water.

Acute toxicity studies of formulations of abamectin with fish

Organism	Formulation, % ai	Endpoint	Source (Study Classification)
Rainbow trout (<i>O. mykiss</i>), freshwater, size 0.14 g, static	Fire Ant Bait 0.022 ^(a)	96 hr LC ₅₀ = 5.06 ppb ai (23,000 ppb product) (95% CL 3.52 -7.04 ppb ai) slope = 3.7	00088781 (Supplemental)
Bluegill sunfish (<i>L. macrochirus</i>), freshwater, size 0.34 g, static	Fire Ant Bait	96 hr LC ₅₀ = 57.2 ppb ai (260,000 ppb product) (95% CL 39.6-85.8 ppb ai) slope = 2.14	00088783 (Supplemental)

^(a) 100 mg abamectin/100 lbs of product * 100 = 0.022% abamectin

Fish early life stage and invertebrate life cycle studies with abamectin

Organism	% ai	Endpoint (ppb)	Source (Study Classification)
Rainbow trout (<i>O. mykiss</i>), freshwater, flow through	Tech	NOAEC=0.52 LOAEC 0.96 Based on wet weight	40069609 (Acceptable)
Water flea (<i>D. magna</i>), freshwater, flow through	91.43 (tritium labeled)	21-d NOAEC = 0.03 LOAEC 0.093	00153570 (Acceptable)
Mysid (<i>A. bahia</i>), estuarine/marine, flow through	>99% (tritium labeled)	28-d NOAEC = 0.0035 LOAEC=0.0093	40856306 (Supplemental)

Acute and sub-acute toxicity studies with abamectin technical grade

Organism	% ai	Endpoint	Source (Study Classification)
Mallard duck (<i>Anas platyrhynchos</i>), age 5 months, oral dosing	91.4	14-d (post-dosing observation) LD ₅₀ = 85 mg/kg-bw (95% CL 67-120) slope = 7.3	ACC246358 (Supplemental)
Bobwhite quail (<i>C. virginianus</i>), age 12 months, oral dosing	91	14 D (post-dosing observation) LD ₅₀ = >2000 mg/kg-bw	ACC250762 (Acceptable)
Mallard duck (<i>Anas platyrhynchos</i>), age 10 days, dietary dosing	91	8-d (3-d post-dosing observation) LC ₅₀ = 383 ppm (95% CL 302-487) slope = 7.25	ACC250761 (Acceptable)
Bobwhite quail (<i>C. virginianus</i>), age 14 days, dietary dosing	91	8 D (3 day post-dosing observation) LC ₅₀ = 3102 ppm (95% CL 2344 - 4415) slope = 4.4	ACC250763 (Acceptable)

Avian reproduction studies with abamectin technical grade

Organism	% ai	Endpoint	Source (Study Classification)
Mallard duck (<i>Anas platyrhynchos</i>), dietary	94.7	NOAEL = 12 ppm LOAEL = 64 ppm (from pilot study)	40318601 (Acceptable)

Terrestrial invertebrate toxicity studies with abamectin

Organism	% ai	Endpoint	Source (Study Classification)
Honey bee (Honey bee), age Worker, contact	Tech	48 hr (3 day post-dosing observation) LD ₅₀ = 0.41 µg ai/bee	00159162 (Acceptable)
Honey bee (Honey bee), age Adult, foliar residues	FORM	8 hr (3 day post-dosing observation) LD ₅₀ = <0.05 lbs ai/A	00159161 (Acceptable)
Earthworm (Earthworm), age Adult, soil exposure	97	28-d LC ₅₀ = 18 ppm ai (95% CL 14 - 32)	40318603 (Supplemental)

Mammalian toxicity profile of abamectin^(a)

Guideline No./ Study Type	Results	MRID #, Study Classification, Dosage
---------------------------	---------	--------------------------------------

Guideline No./ Study Type	Results	MRID #, Study Classification, Dosage
81-1 Acute oral – rat (sesame oil vehicle)	LD ₅₀ = 13.6 mg/kg-bw	006894
81-1 Acute oral – rat (methyl cellulose vehicle)	LD ₅₀ = 214 – 232 mg/kg-bw	45607202
81-2 Acute Dermal – rabbit	LD ₅₀ = 2000 mg/kg-bw	0025978
81-3 Acute Inhalation – rat	LC ₅₀ ≤ 0.21 mg/L (nose only)	45623501
81-4 Primary Eye Irritation	Not an irritant	45063501
81-5 Primary Skin Irritation	Slight irritation	41123904
81-6 Dermal Sensitization	Negative in Buehler	--
81-8 Acute Neurotoxicity	None	None
870.3700a Prenatal developmental in rodents-rats	<u>Maternal NOAEL</u> > 1.6 mg/kg-bw/day <u>Maternal LOAEL</u> = not established <u>Developmental NOAEL</u> > 1.6 mg/kg-bw/day <u>Developmental LOAEL</u> = not established	Accession: 249152 (1982) Acceptable/guideline 0, 0.4, 0.8, 1.6 mg/kg- bw/day
870.3700a Prenatal developmental in rodents-CD-1 mouse	<u>Maternal NOAEL</u> = 1.5 mg/kg-bw/day <u>Maternal LOAEL</u> = 3.0 mg/kg-bw/day based on hind limb splay <u>Developmental NOAEL</u> < 0.75 mg/kg-bw/day <u>Developmental LOAEL</u> = 0.75 mg/kg-bw/day based on cleft palate and hindlimb extension	44179901 (1999) Acceptable/Non-Guideline 0, 0.75, 1.5, 3.0 mg/kg- bw/day
870.3700b Prenatal developmental in nonrodentsrabbits	<u>Maternal NOAEL</u> = 1.0 mg/kg-bw/day <u>Maternal LOAEL</u> = 2.0 mg/kg-bw/day based on decreased body weight, food consumption and water consumption <u>Developmental NOAEL</u> = 1.0 mg/kg-bw/day <u>Developmental LOAEL</u> = 2.0 mg/kg-bw/day based on cleft palate, clubbed foot, delayed ossification of sternebrae, metacarpals, phalanges	Accession: 249152 (1989) Acceptable./Guideline 0, 1.0, 2.0 mg/kg-bw/day
870.3800a 2-Generation Reproduction and fertility effects-rat	<u>Parental/Systemic NOAEL</u> = 0.40 mg/kg/day <u>Parental/systemics LOAEL</u> = not established <u>Reproductive NOAEL</u> = 0.40 mg/kg/day <u>Reproductive LOAEL</u> = not established <u>Offspring NOAEL</u> = 0.12 mg/kg-bw/day <u>Offspring LOAEL</u> = 0.40 mg/kg-bw/day based on increased retinal folds, increased dead pups at birth, decreased viability and lactation indices, decreased pup body weight	00164151 (1984) Acceptable/Guideline 0, 0.05, 0.12, 0.40 mg/kg-bw/day

Guideline No./ Study Type	Results	MRID #, Study Classification, Dosage
870.3800b 1-Generation Reproduction and fertility effects-rat	<u>Parental/Systemic NOAEL</u> = 1.0 mg/kg-bw/day. <u>Parental/Systemic LOAEL</u> =1.5/2.0 mg/kg- bw/day based on whole body tremors, ataxia, ptyalis, ocular/nasal discharges and mortality <u>Reproductive NOAEL</u> = 3.0 mg/kg-bw/day <u>Offspring NOAEL</u> < 0.5 mg/kg/day <u>Offspring LOAEL</u> = 0.5 mg/kg/day based on decreased pup survival and body weight between days 1-21 and delay in opening of eyes	00096450 Unacceptable/Non- Guideline 0, 0.5, 1.0, 1.5/2.0 mg/kg-bw/day
870.3800c 1-Generation Reproduction and fertility effects- rat	<u>Parental/Systemic NOAEL</u> = 0.4 mg/kg-bw/day <u>Parental/Stemic LOAEL</u> = not established <u>Reproductive NOAEL</u> = 0.4 mg/kg-bw/day <u>Offspring NOAEL</u> =0.1 mg/kg-bw/day <u>Offspring LOAEL</u> = 0.2 mg/kg-bw/day based on reduced pup weight, spastic movements, delayed incisor eruption	00096451 Unacceptable/Non- guideline 0, 0.1, 0.2, 0.4 mg/kg-bw/day
870.3800c 1-Generation Reproduction and fertility effects- rat	<u>Parental/Systemic NOAEL</u> = 0.4 mg/kg-bw/day <u>Parental/Systemic LOAEL</u> = not established <u>Reproductive NOAEL</u> = 0.4 mg/kg-bw/day <u>Offspring NOAEL</u> =0.4 mg/kg-bw/day LOAEL = not established	40713404 (1988) Acceptable/Nonguideline 0, 0.1, 0.2, 0.4 mg/kg- bw/day with delta-8,9 isomer 0, 0.06, 0.12, 0.40 mg/kg-bw/day
870.4300a Combined Chronic toxicity/carcinogenicity- rats	NOAEL = 1.5 mg/kg-bw/day LOAEL = 2.0 mg/kg-bw/day based on tremors. No evidence of carcinogenicity	40069601, 40375511, 40517801 (1985) Acceptable/Guideline 0, 0.75, 1.5, 2.0 mg/kg-bw/day
870.3150a Subchronic toxicity dogs	NOAEL = 0.25 mg/kg-bw/day LOAEL = 0.50 mg/kg/day based on body tremors, one death, liver pathology, decreased body weight	00131082 Acceptable/Guideline 0, 0.25, 0.5, 2.0, 8.0 mg/kg/day
870.4100b Chronic toxicity dogs	40375510 (1987) Acceptable/Guideline 0, 0.25, 0.5, 1.0 mg/kg-bw/day	NOAEL = 0.25 mg/kg/day LOAEL = 0.5 mg/kg/day based on mydriasis, death at 1.0 mg/kg/day
870.4300b Combined Chronic toxicity/Carcinogenicity- mice	NOAEL = 4.0 mg/kg-bw/day LOAEL = 8.0 mg/kg-bw/day based on increased mortality in males, tremors, body weight decreases in females, dermatitis in males, extramedullary hematopoiesis in spleen of males No evidence of carcinogenicity	40069602, 40375512, 40517801 (1985) Acceptable/Guideline 0, 2, 4, 8 mg/kg-bw/day

Guideline No./ Study Type	Results	MRID #, Study Classification, Dosage
Gene Mutation 870.5100 Ames/Salmonella E.coli/mammalian gene mutation assay	negative both with and without S-9	Accession: 246894, 265568, 265569 (1986) Acceptable/Guideline Three studies: (1) 0, 3, 10, 30, 100, 1000 ug/plate, (2) 0, 100, 300, 1000, 3000, 10,000 ug/plate both with and without S-9, (3) doses not specified
Gene Mutation 870.5100 Ames/Salmonella E.coli/mammalian gene mutation assay	negative both with and without S-9 up to 3000 ug/plate	40713402 (1988) Acceptable/Guideline doses not specified up to 3000 ug/plate both with and without S-9 using delta-8,9 isomer
Gene Mutation 870.5100 Ames/SalmonellaE.coli/ mammalian gene mutation assay	negative both with and without S-9	40713405 (1988) Acceptable/Guideline doses up to 10,000 ug/plate both with and without S-9 using polar degradates
Gene Mutation 870.5300 CHO/HGPRT Forward Mutation Assay	Negative	265570 (1986) Acceptable/Guideline both with and without S-9
Gene Mutation 870.5300 Mammalian cells in culture in V79 cells	Not mutagenic for V79 cells in absence of S-9, but in the presence of S-9 appeared to have a mutagenic potential, provided the test cells had an appropriate level of sensitivity	MRID Unavailable 1983 Acceptable/Guideline
Cytogenetics 870.5395 in vivo micronucleus assay -male mice	No chromosomal aberrations in male mice; but females not tested.	MRID Unavailable Acceptable/non-Guideline 0, 1.2, 12.0 mg/kg i.p.
Other Effects 870.5550	single strand DNA breaks at 0.3 and 0.6 mM in rat hepatocytes in vitro, but negative when hepatocytes from rat at LD50 dose level was used	MRID Unavailable (1983) 0.3 and 0.6 mM
Metabolism	Avermectin B1a did not bioaccumulate in rat tissues. Half-life slightly longer in females than in males for several tissues.	No MRID (1985) Nonguideline
Metabolism	The metabolism of avermectin B1 in rats results in the formation of 24-OH-Me-B1a and accounts for most of the radiolabeled residues. Avermectin B1a does not bioaccumulate.	No MRID (1985) Nonguideline

Guideline No./ Study Type	Results	MRID #, Study Classification, Dosage
870.7600 Dermal penetration	Dermal penetration is 1%	Accession: 265590 (1986) Acceptable/Nonguideline in Monkeys.

^(a) Source: Rourke *et al.* November 2, 1994 Human Health Risk Assessment for New uses on Plums/Prunes, Leafy Vegetables, Fruiting Vegetables, Herb Subgroup (except chives), Avocado, Mint, and Food Handling Establishments. DB Barcode: D297225

Appendix E. RQ Method and LOCs

Risk Presumption	RQ	LOC
Birds and Wild Mammals		
Acute Risk	Dietary based: EEC ^a (ppm) ^b / LC ₅₀ (ppm) Dose based: EEC (mg/kg-bw/d) / LD ₅₀ (mg/kg-bw/d) ^c	0.5
Acute Restricted Use	Dietary based: EEC (ppm) / LC ₅₀ (ppm) Dose based: EEC (mg/kg-bw/d) / LD ₅₀ (mg/kg-bw/d)	0.2
Acute Listed Species	Dietary based: EEC (ppm) / LC ₅₀ (ppm) Dose based: EEC (mg/kg-bw/d) / LD ₅₀ (mg/kg-bw/d)	0.1
Chronic Risk	Dietary based: EEC (ppm) / NOAEC (ppm) Dose based: EEC (mg/kg-bw/d) / NOAEL (mg/kg-bw/d)	1.0
Aquatic Animals		
Acute Risk	EEC (ppm) / (LC ₅₀ (ppm) or EC ₅₀ (ppm))	0.5
Acute Restricted Use	EEC (ppm) / (LC ₅₀ (ppm) or EC ₅₀ (ppm))	0.1
Acute Listed Species	EEC (ppm) / (LC ₅₀ (ppm) or EC ₅₀ (ppm))	0.05
Chronic Risk	EEC (ppm) / NOAEC (ppm)	1.0
Terrestrial Plants and Plants Inhabiting Semi-Aquatic Areas		
Acute Risk	EEC (lbs ai/A) / EC ₂₅ (lbs ai/A)	1.0
Acute Listed Use	EEC (lbs ai/A) / (EC ₀₅ or NOAEC (lbs ai/A))	1.0
Aquatic Plants		
Risk	EEC (ppm) / EC ₅₀ (ppm)	1.0
Listed Species	EEC (ppm) / (EC ₀₅ or NOAEC (ppm))	1.0

^aEEC = estimated environmental concentration

^bppm = parts per million

^cmg/kg-bw/d = milligrams per kilogram of body weight per day

Appendix F. Locates Output

All Medium Types Reported

Mammal, Marine mml, Bird, Amphibian, Reptile, Crustacean, Bivalve, Gastropod, Arachnid, Insect, Dicot, Monocot, Ferns, Conf/cycds, Coral, Lichen

almonds, walnuts, english, apples, avocados, avocados (PR), citrus fruit, all, cotton, all, cantaloups, cucumbers and pickles, honeydew melons, pumpkins, squash, watermelons, eggplant, peppers, bell, peppers, chile (all peppers - excluding bell), pimientos, tomatoes, grapes, dill for oil, dill for oil (irrigated), herbs and spice plants harvested for sale (PR), herbs, dried, herbs, fresh cut, mustard seed, parsley, amaranth, celery, escarole and endive, lettuce, all, rhubarb, spinach, mint for oil, all (irrigated), mint for oil, peppermint (irrigated), mint for oil, spearmint (irrigated), pears, all, plums and prunes, potatoes

AL, AK, AZ, AR, CA, CO, CT, DE, DC, FL, GA, HI, ID, IL, IN, IA, KS, KY, LA, ME, MD, MA,

MI, MN, MS, MO, MT, NE, NV, NH, NJ, NM, NY, NC, ND, OH, OK, OR, PA, PR, RI, SC, SD, TN, TX, UT, VT, VA, WA, WV, WI, WY

1145 Species Affected:

Inverse Name:	Taxa:	Co. occurrence:
Status:		
Abalone, White	Gastropod	118
Endangered		
Abutilon eremipetalum (ncn)	Dicot	20
Endangered		
Abutilon sandwicense (ncn)	Dicot	17
Endangered		
Achyranthes mutica (ncn)	Dicot	20
Endangered		
Achyranthes splendens var. rotundata (ncn)	Dicot	17
Endangered		
A'e (Zanthoxylum dipetalum var. tomentosum)	Dicot	20
Endangered		
A'e (Zanthoxylum hawaiiense)	Dicot	56
Endangered		
'Aiea (Nothocestrum breviflorum)	Dicot	20
Endangered		
'Aiea (Nothocestrum peltatum)	Dicot	16
Endangered		
'Akepa, Hawaii	Bird	20
Endangered		
'Akepa, Maui	Bird	20
Endangered		
'Akia Loa, Kauai (Hemignathus procerus)	Bird	16
Endangered		
'Akia Pola'au (Hemignathus munroi)	Bird	20
Endangered		
'Akoko (Chamaesyce celastroides var. kaenana)	Dicot	17
Endangered		
'Akoko (Chamaesyce deppeana)	Dicot	17
Endangered		
'Akoko (Chamaesyce herbstii)	Dicot	17
Endangered		
'Akoko (Chamaesyce kuwaleana)	Dicot	17
Endangered		
'Akoko (Chamaesyce rockii)	Dicot	17
Endangered		
'Akoko (Chamaesyce skottsbergii var. skottsbe)	Dicot	37
Endangered		
'Akoko (Euphorbia haelecleana)	Dicot	33

Endangered		
Alani (Melicope adscendens)	Dicot	20
Endangered		
Alani (Melicope balloui)	Dicot	20
Endangered		
Alani (Melicope haupuensis)	Dicot	16
Endangered		
Alani (Melicope knudsenii)	Dicot	36
Endangered		
Alani (Melicope lydgatei)	Dicot	17
Endangered		
Alani (Melicope mucronulata)	Dicot	20
Endangered		
Alani (Melicope munroi)	Dicot	20
Endangered		
Alani (Melicope ovalis)	Dicot	20
Endangered		
Alani (Melicope pallida)	Dicot	16
Endangered		
Alani (Melicope quadrangularis)	Dicot	16
Endangered		
Alani (Melicope reflexa)	Dicot	20
Endangered		
Alani (Melicope saint-johnii)	Dicot	17
Endangered		
Alani (Melicope zahlbruckneri)	Dicot	20
Endangered		
Albatross, Short-tailed	Bird	17
Endangered		
Allocarya, Calistoga	Dicot	21
Endangered		
Alopecurus, Sonoma	Monocot	44
Endangered		
Alsinidendron obovatum (ncn)	Dicot	17
Endangered		
Alsinidendron trinerve (ncn)	Dicot	17
Endangered		
Alsinidendron viscosum (ncn)	Dicot	16
Endangered		
Amaranthus brownii (ncn)	Dicot	17
Endangered		
Ambersnail, Kanab	Gastropod	23
Endangered		
Ambrosia, San Diego	Dicot	54
Endangered		
Ambrosia, South Texas	Dicot	43
Endangered		
Amphipod, Illinois Cave	Crustacean	27
Endangered		
Amphipod, Kauai Cave	Crustacean	16
Endangered		
Amphipod, Noel's	Crustacean	4
Endangered		
Amphipod, Peck's Cave	Crustacean	75
Endangered		
'Anaunau (Lepidium arbuscula)	Dicot	17
Endangered		
'Anunu (Sicyos alba)	Dicot	20
Endangered		
Aristida chaseae (ncn)	Monocot	3
Endangered		
Arrowhead, Bunched	Monocot	41
Endangered		
Asplenium fragile var. insulare (ncn)	Ferns	20
Endangered		
Aster, Florida Golden	Dicot	29
Endangered		

Aster, Ruth's Golden	Dicot	5
Endangered		
Auerodendron pauciflorum (ncn)	Dicot	2
Endangered		
Aupaka (Isodendron hosakae)	Dicot	20
Endangered		
Aupaka (Isodendron laurifolium)	Dicot	33
Endangered		
Avens, Spreading	Dicot	111
Endangered		
'Awikiwiki (Canavalia molokaiensis)	Dicot	20
Endangered		
'Awiwi (Centaurium sebacoides)	Dicot	36
Endangered		
'Awiwi (Hedyotis cookiana)	Dicot	36
Endangered		
Ayenia, Texas	Dicot	41
Endangered		
Barberry, Island	Dicot	25
Endangered		
Barberry, Nevin's	Dicot	53
Endangered		
Bariaco	Dicot	7
Endangered		
Bat, Gray	Mammal	1801
Endangered		
Bat, Hawaiian Hoary	Mammal	73
Endangered		
Bat, Indiana	Mammal	9735
Endangered		
Bat, Lesser (=Sanborn's) Long-nosed	Mammal	154
Endangered		
Bat, Mexican Long-nosed	Mammal	28
Endangered		
Bat, Ozark Big-eared	Mammal	49
Endangered		
Bat, Virginia Big-eared	Mammal	259
Endangered		
Beardtongue, Penland	Dicot	2
Endangered		
Beargrass, Britton's	Monocot	51
Endangered		
Bear-poppy, Dwarf	Dicot	13
Endangered		
Bedstraw, El Dorado	Dicot	20
Endangered		
Bedstraw, Island	Dicot	25
Endangered		
Beetle, American Burying	Insect	450
Endangered		
Beetle, Coffin Cave Mold	Insect	16
Endangered		
Beetle, Comal Springs Dryopid	Insect	75
Endangered		
Beetle, Comal Springs Riffle	Insect	75
Endangered		
Beetle, Helotes Mold	Insect	17
Endangered		
Beetle, Hungerford's Crawling Water	Insect	17
Endangered		
Beetle, Kretschmarr Cave Mold	Insect	13
Endangered		
Beetle, Mount Hermon June	Insect	22
Endangered		
Beetle, Ohlone Tiger	Insect	22
Endangered		
Beetle, Salt Creek Tiger	Insect	35
Endangered		
Beetle, Tooth Cave Ground	Insect	29
Endangered		

Bellflower, Brooksville Endangered	Dicot	7
Bird's-beak, Palmate-bracted Endangered	Dicot	133
Bird's-beak, Pennell's Endangered	Dicot	27
Bird's-beak, salt marsh Endangered	Dicot	142
Bird's-beak, Soft Endangered	Dicot	62
Bittercress, Small-anthered Endangered	Dicot	39
Blackbird, Yellow-shouldered Endangered	Bird	19
Bladderpod, Kodachrome Endangered	Dicot	3
Bladderpod, San Bernardino Mountains Endangered	Dicot	25
Bladderpod, Spring Creek Endangered	Dicot	12
Bladderpod, White Endangered	Dicot	6
Bladderpod, Zapata Endangered	Dicot	12
Blazing Star, Scrub Endangered	Dicot	19
Bluegrass, Hawaiian Endangered	Monocot	16
Bluegrass, Mann's (<i>Poa mannii</i>) Endangered	Monocot	16
Bluegrass, Napa Endangered	Monocot	21
Bluegrass, San Bernardino Endangered	Monocot	51
Blue-star, Kearney's Endangered	Dicot	18
Bluet, Roan Mountain Endangered	Dicot	50
Boa, Puerto Rican Endangered	Reptile	24
Bobwhite, Masked Endangered	Bird	18
Bonamia menziesii (ncn) Endangered	Dicot	73

Boxwood, Vahl's	Dicot	4
Endangered		
Broom, San Clemente Island	Dicot	25
Endangered		
Buckwheat, Cushenbury	Dicot	25
Endangered		
Buckwheat, Ione (incl. Irish Hill)	Dicot	14
Endangered		
Buckwheat, Steamboat	Dicot	8
Endangered		
Bulrush, Northeastern (=Barbed Bristle)	Monocot	268
Endangered		
Bush-mallow, San Clemente Island	Dicot	25
Endangered		
Bush-mallow, Santa Cruz Island	Dicot	25
Endangered		
Buttercup, Autumn	Dicot	6
Endangered		
Butterfly, Behren's Silverspot	Insect	48
Endangered		
Butterfly, Callippe Silverspot	Insect	30
Endangered		
Butterfly, El Segundo Blue	Insect	25
Endangered		
Butterfly, Fender's Blue	Insect	83
Endangered		
Butterfly, Karner Blue	Insect	552
Endangered		
Butterfly, Lange's Metalmark	Insect	18
Endangered		
Butterfly, Lotis Blue	Insect	21
Endangered		
Butterfly, Mission Blue	Insect	36
Endangered		
Butterfly, Mitchell's Satyr	Insect	273
Endangered		
Butterfly, Myrtle's Silverspot	Insect	44
Endangered		
Butterfly, Palos Verdes Blue	Insect	25
Endangered		
Butterfly, Quino Checkerspot	Insect	54
Endangered		
Butterfly, Saint Francis' Satyr	Insect	25
Endangered		
Butterfly, San Bruno Elfin	Insect	19
Endangered		
Butterfly, Schaus Swallowtail	Insect	15
Endangered		
Butterfly, Smith's Blue	Insect	27
Endangered		
Butterfly, Uncompahgre Fritillary	Insect	23
Endangered		
Button-celery, San Diego	Dicot	54
Endangered		
Cactus, Arizona Hedgehog	Dicot	47
Endangered		
Cactus, Bakersfield	Dicot	25
Endangered		
Cactus, Black Lace	Dicot	33
Endangered		
Cactus, Brady Pincushion	Dicot	9
Endangered		
Cactus, Key Tree	Dicot	1
Endangered		
Cactus, Knowlton	Dicot	23
Endangered		
Cactus, Kuenzler Hedgehog	Dicot	29
Endangered		
Cactus, Nellie Cory	Dicot	15
Endangered		

Cactus, Nichol's Turk's Head	Dicot	38
Endangered		
Cactus, Peebles Navajo	Dicot	4
Endangered		
Cactus, Pima Pineapple	Dicot	25
Endangered		
Cactus, San Rafael	Dicot	5
Endangered		
Cactus, Sneed Pincushion	Dicot	43
Endangered		
Cactus, Star	Dicot	40
Endangered		
Cactus, Tobusch Fishhook	Dicot	47
Endangered		
Cactus, Wright Fishhook	Dicot	19
Endangered		
Campeloma, Slender	Gastropod	14
Endangered		
Campion, Fringed	Dicot	45
Endangered		
Capa Rosa	Dicot	4
Endangered		
Caribou, Woodland	Mammal	21
Endangered		
Catesbaea Melanocarpa (ncn)	Dicot	1
Endangered		
Cat's-eye, Terlingua Creek	Dicot	15
Endangered		
Cavesnail, Tumbling Creek	Gastropod	5
Endangered		
Ceanothus, Coyote	Dicot	20
Endangered		
Ceanothus, Pine Hill	Dicot	20
Endangered		
Chaffseed, American	Dicot	243
Endangered		
Chamaecrista glandulosa (ncn)	Dicot	5
Endangered		
Chamaesyce Halemanui (ncn)	Dicot	16
Endangered		
Checker-mallow, Keck's	Dicot	53
Endangered		
Checker-mallow, Kenwood Marsh	Dicot	27
Endangered		

Checker-mallow, Pedate	Dicot	25
Endangered		
Checker-mallow, Wenatchee Mountains	Dicot	10
Endangered		
Chupacallos	Dicot	6
Endangered		
Cladonia, Florida Perforate	Lichen	58
Endangered		
Clarkia, Pismo	Dicot	24
Endangered		
Clarkia, Presidio	Dicot	11
Endangered		
Clarkia, Vine Hill	Dicot	27
Endangered		
Cliffrose, Arizona	Dicot	68
Endangered		
Clover, Leafy Prairie	Dicot	140
Endangered		
Clover, Monterey	Dicot	27
Endangered		
Clover, Running Buffalo	Dicot	519
Endangered		
Clover, Showy Indian	Dicot	44
Endangered		
Combshell, Southern (=Penitent mussel)	Bivalve	30
Endangered		
Combshell, Upland	Bivalve	93
Endangered		
Condor, California	Bird	214
Endangered		
Coneflower, Smooth	Dicot	447
Endangered		
Coneflower, Tennessee Purple	Dicot	33
Endangered		
Coot, Hawaiian (=Alae keo keo)	Bird	73
Endangered		
Cordia bellonis (ncn)	Dicot	2
Endangered		
Coyote-thistle, Loch Lomond	Dicot	18
Endangered		
Crane, Mississippi Sandhill	Bird	13
Endangered		
Crane, Whooping	Bird	2256
Endangered		
Cranichis Ricartii	Monocot	4
Endangered		
Crayfish, Cave (Cambarus aculabrum)	Crustacean	18
Endangered		
Crayfish, Cave (Cambarus zophonastes)	Crustacean	3
Endangered		
Crayfish, Nashville	Crustacean	19
Endangered		
Crayfish, Shasta	Crustacean	22
Endangered		
Creeper, Hawaii	Bird	20
Endangered		
Creeper, Molokai (Kakawahie)	Bird	20
Endangered		
Creeper, Oahu (Alauwahio)	Bird	17
Endangered		
Crow, Hawaiian ('Alala)	Bird	20
Endangered		
Crownscale, San Jacinto Valley	Dicot	28
Endangered		
Curlew, Eskimo	Bird	42
Endangered		
Cyanea undulata (ncn)	Dicot	16
Endangered		
Cypress, Santa Cruz	Conf/cycds	41
Endangered		

Daisy, Willamette	Dicot	113
Endangered		
<i>Daphnopsis hellerana</i> (ncn)	Dicot	2
Endangered		
Dawn-flower, Texas Prairie (=Texas Bitterweed)	Dicot	42
Endangered		
Deer, Columbian White-tailed	Mammal	94
Endangered		
Deer, Key	Mammal	1
Endangered		
<i>Delissea rhytidisperma</i> (ncn)	Dicot	16
Endangered		
<i>Diellia erecta</i> (ncn)	Ferns	57
Endangered		
<i>Diellia falcata</i> (ncn)	Ferns	17
Endangered		
<i>Diellia pallida</i> (ncn)	Ferns	16
Endangered		
<i>Diellia unisora</i> (ncn)	Ferns	37
Endangered		
<i>Diplazium molokaiense</i> (ncn)	Ferns	20
Endangered		
Dogweed, Ashy	Dicot	16
Endangered		
Dragonfly, Hine's Emerald	Insect	163
Endangered		
Dropwort, Canby's	Dicot	272
Endangered		
<i>Dubautia latifolia</i> (ncn)	Dicot	16
Endangered		
<i>Dubautia pauciflora</i> (ncn)	Dicot	16
Endangered		
Duck, Hawaiian (Koloa)	Bird	53
Endangered		
Duck, Laysan	Bird	17
Endangered		
Dudleya, Santa Clara Valley	Dicot	128
Endangered		
Elepaio, Oahu	Bird	17
Endangered		
Elktoe, Appalachian	Bivalve	97
Endangered		
Erubia	Dicot	2
Endangered		

Eugenia Woodburyana	Dicot	5
Endangered		
Evening-primrose, Antioch Dunes	Dicot	41
Endangered		
Evening-primrose, Eureka Valley	Dicot	6
Endangered		
Fairy Shrimp, Conservancy Fairy	Crustacean	133
Endangered		
Fairy Shrimp, Longhorn	Crustacean	105
Endangered		
Fairy Shrimp, Riverside	Crustacean	121
Endangered		
Fairy Shrimp, San Diego	Crustacean	26
Endangered		
Falcon, Northern Aplomado	Bird	305
Endangered		
Fanshell	Bivalve	676
Endangered		
Fern, Adiantum vivesii	Ferns	2
Endangered		
Fern, Aleutian Shield	Ferns	3
Endangered		
Fern, Elaphoglossum serpens	Ferns	2
Endangered		
Fern, Pendant Kihi (Adenophorus periens)	Ferns	56
Endangered		
Fern, Thelypteris inabonensis	Ferns	4
Endangered		
Fern, Thelypteris verecunda	Ferns	5
Endangered		
Fern, Thelypteris yaucoensis	Ferns	4
Endangered		
Ferret, Black-footed	Mammal	708
Endangered		
Fiddleneck, Large-flowered	Dicot	53
Endangered		
Finch, Laysan	Bird	17
Endangered		
Finch, Nihoa	Bird	17
Endangered		
Flannelbush, Mexican	Dicot	26
Endangered		
Flannelbush, Pine Hill	Dicot	20
Endangered		
Fly, Delhi Sands Flower-loving	Insect	76
Endangered		
Flycatcher, Southwestern Willow	Bird	641
Endangered		
Fox, San Joaquin Kit	Mammal	360
Endangered		
Fox, San Miguel Island	Mammal	25
Endangered		
Fox, Santa Catalina Island	Mammal	25
Endangered		
Fox, Santa Cruz Island	Mammal	25
Endangered		
Fox, Santa Rosa Island	Mammal	25
Endangered		
Frankenia, Johnston's	Dicot	16
Endangered		
Fringe Tree, Pygmy	Dicot	45
Endangered		
Fringepod, Santa Cruz Island	Dicot	25
Endangered		
Fritillary, Gentner's	Monocot	39
Endangered		
Frog, Dusky Gopher (Mississippi DPS)	Amphibian	12
Endangered		
Frog, Mountain Yellow-legged	Amphibian	78
Endangered		

Gahnia Lanaiensis (ncn)	Monocot	20
Endangered		
Gecko, Monito	Reptile	2
Endangered		
Geranium, Hawaiian Red-flowered	Dicot	20
Endangered		
Gerardia, Sandplain	Dicot	119
Endangered		
Gilia, Hoffmann's Slender-flowered	Dicot	25
Endangered		
Gilia, Monterey	Dicot	27
Endangered		
Goetzea, Beautiful (Matabuey)	Dicot	4
Endangered		
Golden Sunburst, Hartweg's	Dicot	76
Endangered		
Goldenrod, Short's	Dicot	63
Endangered		
Goldfields, Burke's	Dicot	66
Endangered		
Goldfields, Contra Costa	Dicot	166
Endangered		
Goose, Hawaiian (Nene)	Bird	56
Endangered		
Gouania hillebrandii (ncn)	Dicot	20
Endangered		
Gouania meyenii (ncn)	Dicot	33
Endangered		
Gouania vitifolia (ncn)	Dicot	17
Endangered		
Gourd, Okeechobee	Dicot	22
Endangered		
Grass, California Orcutt	Monocot	79
Endangered		
Grass, Eureka Dune	Monocot	6
Endangered		
Grass, Fosberg's Love	Monocot	17
Endangered		
Grass, Hairy Orcutt	Dicot	200
Endangered		
Grass, Sacramento Orcutt	Dicot	37
Endangered		
Grass, Solano	Monocot	48
Endangered		

Grass, Tennessee Yellow-eyed	Monocot	46
Endangered		
Grasshopper, Zayante Band-winged	Insect	22
Endangered		
Ground-plum, Guthrie's	Dicot	21
Endangered		
Haha (<i>Cyanea acuminata</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea asarifolia</i>)	Dicot	16
Endangered		
Haha (<i>Cyanea copelandii</i> ssp. <i>copelandii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea Crispa</i>) (=Rollandia <i>crispa</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea dunbarii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea glabra</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea grimesiana</i> ssp. <i>grimesiana</i>)	Dicot	37
Endangered		
Haha (<i>Cyanea grimesiana</i> ssp. <i>obatae</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea humboldtiana</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea koolauensis</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea longiflora</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea Macrostegia</i> var. <i>gibsonii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea mannii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea mceldowneyi</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea pinnatifida</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea platyphylla</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea procera</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea remyi</i>)	Dicot	16
Endangered		
Haha (<i>Cyanea shipmanii</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea stictophylla</i>)	Dicot	20
Endangered		
Haha (<i>Cyanea St-Johnii</i>) (=Rollandia <i>St-Johnii</i>)	Dicot	17
Endangered		
Haha (<i>Cyanea superba</i>)	Dicot	17
Endangered		
Ha'Iwale (<i>Cyrtandra crenata</i>)	Dicot	17
Endangered		
Ha'Iwale (<i>Cyrtandra dentata</i>)	Dicot	17
Endangered		
Ha'Iwale (<i>Cyrtandra giffardii</i>)	Dicot	20
Endangered		
Ha'Iwale (<i>Cyrtandra munroi</i>)	Dicot	20
Endangered		
Ha'Iwale (<i>Cyrtandra polyantha</i>)	Dicot	17
Endangered		
Ha'Iwale (<i>Cyrtandra subumbellata</i>)	Dicot	17
Endangered		
Ha'Iwale (<i>Cyrtandra tintinnabula</i>)	Dicot	17
Endangered		

Ha'Iwale (<i>Cyrtandra viridiflora</i>)	Dicot	17
Endangered		
Hala Pepe (<i>Pleomele hawaiiensis</i>)	Monocot	20
Endangered		
Haplostachys <i>Haplostachya</i> (ncn)	Dicot	20
Endangered		
Harebells, Avon Park	Dicot	19
Endangered		
Harperella	Dicot	260
Endangered		
Harvestman, Bee Creek Cave	Arachnid	41
Endangered		
Harvestman, Bone Cave	Arachnid	29
Endangered		
Harvestman, Robber Baron Cave	Arachnid	17
Endangered		
Hau Kauhiwi (<i>Hibiscadelphus woodi</i>)	Dicot	16
Endangered		
Hau Kuahiwi (<i>Hibiscadelphus distans</i>)	Dicot	16
Endangered		
Hawk, Hawaiian (Io)	Bird	20
Endangered		
Hawk, Puerto Rican Broad-winged	Bird	4
Endangered		
Hawk, Puerto Rican Sharp-shinned	Bird	6
Endangered		
Heau (<i>Exocarpos luteolus</i>)	Dicot	16
Endangered		
Hedyotis <i>degeneri</i> (ncn)	Dicot	17
Endangered		
Hedyotis <i>parvula</i> (ncn)	Dicot	17
Endangered		
Hedyotis <i>St.-Johnii</i> (ncn)	Dicot	16
Endangered		
Hesperomannia <i>arborescens</i> (ncn)	Dicot	37
Endangered		
Hesperomannia <i>arbuscula</i> (ncn)	Dicot	37
Endangered		
Hesperomannia <i>lydgatei</i> (ncn)	Dicot	16
Endangered		
Hibiscus, Clay's	Dicot	16
Endangered		
Higuero De Sierra	Dicot	7
Endangered		

1/28/2010 10:50:28 AM Ver. 2.10.4

Page 113 of 128

Hilo Ischaemum (<i>Ischaemum byrone</i>)	Monocot	56
Endangered		
Holei (<i>Ochrosia kilaueaensis</i>)	Dicot	20
Endangered		
Holly, Cook's	Dicot	4
Endangered		
Honeycreeper, Crested ('Akohekohe)	Bird	20
Endangered		
Hypericum, Highlands Scrub	Dicot	19
Endangered		
'Ihi'Ihi (<i>Marsilea villosa</i>)	Ferns	37
Endangered		
Ilex sintenisii (ncn)	Dicot	4
Endangered		
Iliau (<i>Wilkesia hobbii</i>)	Dicot	16
Endangered		
Ipomopsis, Holy Ghost	Dicot	7
Endangered		
Irisette, White	Monocot	51
Endangered		
Isopod, Lee County Cave	Crustacean	9
Endangered		
Isopod, Socorro	Crustacean	11
Endangered		
Jacquemontia, Beach	Dicot	40
Endangered		
Jaguar	Mammal	92
Endangered		
Jaguarundi, Gulf Coast	Mammal	156
Endangered		
Jaguarundi, Sinaloa	Mammal	108
Endangered		
Jewelflower, California	Dicot	146
Endangered		
Jewelflower, Tiburon	Dicot	17
Endangered		
Kamakahala (<i>Labordia cyrtandrae</i>)	Dicot	17
Endangered		
Kamakahala (<i>Labordia lydgatei</i>)	Dicot	16
Endangered		
Kamakahala (<i>Labordia tinifolia</i> var. <i>lanaiensis</i>)	Dicot	20
Endangered		
Kamakahala (<i>Labordia tinifolia</i> var. <i>wahiawaen</i>)	Dicot	16
Endangered		
Kamakahala (<i>Labordia triflora</i>)	Dicot	20
Endangered		
Kamanomano (<i>Cenchrus agrimonoides</i>)	Monocot	37
Endangered		
Kanaloa kahoolawensis (ncn)	Dicot	20
Endangered		
Kangaroo Rat, Fresno	Mammal	93
Endangered		
Kangaroo Rat, Giant	Mammal	219
Endangered		
Kangaroo Rat, Morro Bay	Mammal	24
Endangered		
Kangaroo Rat, San Bernardino Merriam's	Mammal	53
Endangered		
Kangaroo Rat, Stephens'	Mammal	79
Endangered		
Kangaroo Rat, Tipton	Mammal	68
Endangered		
Kauila (<i>Colubrina oppositifolia</i>)	Dicot	20
Endangered		
Kaulu (<i>Pteralyxia kauaiensis</i>)	Dicot	16
Endangered		
Kidneyshell, Triangular	Bivalve	167
Endangered		
Kio'Ele (<i>Hedyotis coriacea</i>)	Dicot	40
Endangered		

Kiponapona (<i>Phyllostegia racemosa</i>)	Dicot	20
Endangered		
Kite, Everglade Snail	Bird	179
Endangered		
Koki'o (<i>Kokia drynarioides</i>)	Dicot	20
Endangered		
Koki'o (<i>Kokia kauaiensis</i>)	Dicot	16
Endangered		
Koki'o Ke'oke'o (<i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i>)	Dicot	20
Endangered		
Koki'o Ke'oke'o (<i>Hibiscus waimeae</i> ssp. <i>hannerae</i>)	Dicot	16
Endangered		
Kolea (<i>Myrsine juddii</i>)	Dicot	17
Endangered		
Ko'oko'olau (<i>Bidens micrantha</i> ssp. <i>kalcalaha</i>)	Dicot	20
Endangered		
Ko'oko'olau (<i>Bidens wiebkei</i>)	Dicot	20
Endangered		
Ko'olua'ula (<i>Abutilon menziesii</i>)	Dicot	40
Endangered		
Kopa (<i>Hedyotis schlechtendahlia</i> var. <i>remyi</i>)	Dicot	20
Endangered		
Kuawawaenuhu (<i>Alsiniidendron lychnoides</i>)	Dicot	16
Endangered		
Kulu'l (<i>Nototrichium humile</i>)	Dicot	37
Endangered		
Ladies'-tresses, Canelo Hills	Monocot	26
Endangered		
Ladies'-tresses, Navasota	Monocot	131
Endangered		
Larkspur, Baker's	Dicot	44
Endangered		
Larkspur, San Clemente Island	Dicot	25
Endangered		
Larkspur, Yellow	Dicot	44
Endangered		
Lau'ehu (<i>Panicum niihauense</i>)	Monocot	16
Endangered		
Laukahi Kuahiwi (<i>Plantago hawaiiensis</i>)	Dicot	20
Endangered		
Laukahi Kuahiwi (<i>Plantago princeps</i>)	Dicot	53
Endangered		
Laulihilihi (<i>Schiedea stellarioides</i>)	Dicot	16
Endangered		

Layia, Beach Endangered	Dicot	89
Lead-plant, Crenulate Endangered	Dicot	14
Leather-flower, Alabama Endangered	Dicot	37
Leather-flower, Morefield's Endangered	Dicot	15
Lepanthes eltoensis (ncn) Endangered	Monocot	4
Lessingia, San Francisco Endangered	Dicot	19
Lichen, Rock Gnome Endangered	Lichen	125
Lily, Minnesota Trout Endangered	Monocot	44
Lily, Pitkin Marsh Endangered	Monocot	27
Lily, Western Endangered	Monocot	29
Limpet, Banbury Springs Endangered	Gastropod	29
Lipochaeta venosa (ncn) Endangered	Dicot	20
Liveforever, Santa Barbara Island Endangered	Dicot	25
Lizard, Blunt-nosed Leopard Endangered	Reptile	264
Lo'ulu (Pritchardia affinis) Endangered	Monocot	20
Lo'ulu (Pritchardia kaalae) Endangered	Monocot	17
Lo'ulu (Pritchardia munroi) Endangered	Monocot	20
Lo'ulu (Pritchardia napaliensis) Endangered	Monocot	16
Lo'ulu (Pritchardia remota) Endangered	Monocot	17
Lo'ulu (Pritchardia schattaueri) Endangered	Monocot	20
Lo'ulu (Pritchardia viscosa) Endangered	Monocot	16
Lobelia monostachya (ncn) Endangered	Dicot	17
Lobelia niihauensis (ncn) Endangered	Dicot	33
Lobelia oahuensis (ncn) Endangered	Dicot	17
Lomatium, Bradshaw's Endangered	Dicot	113
Lomatium, Cook's Endangered	Dicot	39
Loosestrife, Rough-leaved Endangered	Dicot	237
Lousewort, Furbish Endangered	Dicot	20
Lupine, Clover Endangered	Dicot	93
Lupine, Nipomo Mesa Endangered	Dicot	24
Lupine, Scrub Endangered	Dicot	19
Lyonia truncata var. proctorii (ncn) Endangered	Dicot	3
Lysimachia filifolia (ncn) Endangered	Dicot	33
Lysimachia lydgatei (ncn) Endangered	Dicot	37
Lysimachia maxima (ncn) Endangered	Dicot	20

Mahoe (<i>Alectryon macrococcus</i>)	Dicot	53
Endangered		
Malacothrix, Island	Dicot	25
Endangered		
Malacothrix, Santa Cruz Island	Dicot	50
Endangered		
Mallow, Kern	Dicot	25
Endangered		
Mallow, Peter's Mountain	Dicot	5
Endangered		
Manatee, West Indian	Marine mml	606
Endangered		
Manioc, Walker's	Dicot	28
Endangered		
Manzanita, Del Mar	Dicot	43
Endangered		
Manzanita, Santa Rosa Island	Dicot	25
Endangered		
Ma'o Hau Hele (<i>Hibiscus brackenridgei</i>)	Dicot	57
Endangered		
Ma'oli'oli (<i>Schiedea apokremnos</i>)	Dicot	16
Endangered		
Ma'oli'oli (<i>Schiedea kealiae</i>)	Dicot	17
Endangered		
Mapele (<i>Cyrtandra cyaneoides</i>)	Dicot	16
Endangered		
Mariscus <i>fauriei</i> (ncn)	Monocot	40
Endangered		
Mariscus <i>pennatifolius</i> (ncn)	Monocot	73
Endangered		
Marstonia, Royal (=Royal Snail)	Gastropod	5
Endangered		
Meadowfoam, Butte County	Dicot	43
Endangered		
Meadowfoam, Large-flowered Woolly	Dicot	21
Endangered		
Meadowfoam, Sebastopol	Dicot	27
Endangered		
Meadowrue, Cooley's	Dicot	66
Endangered		
Mehamchame (<i>Flueggea neowawraea</i>)	Dicot	73
Endangered		
Meshweaver, Braken Bat Cave	Arachnid	17
Endangered		

Milkpea, Small's	Dicot	14
Endangered		
Milk-vetch, Applegate's	Dicot	14
Endangered		
Milk-vetch, Braunton's	Dicot	67
Endangered		
Milk-vetch, Clara Hunt's	Dicot	48
Endangered		
Milk-vetch, Coachella Valley	Dicot	28
Endangered		
Milk-vetch, Coastal Dunes	Dicot	27
Endangered		
Milk-vetch, Cushenbury	Dicot	25
Endangered		
Milk-vetch, Holmgren	Dicot	27
Endangered		
Milk-vetch, Jesup's	Dicot	29
Endangered		
Milk-vetch, Lane Mountain	Dicot	25
Endangered		
Milk-vetch, Mancos	Dicot	29
Endangered		
Milk-vetch, Osterhout	Dicot	2
Endangered		
Milk-vetch, Sentry	Dicot	9
Endangered		
Milk-vetch, Shivwits	Dicot	13
Endangered		
Milk-vetch, Triple-ribbed	Dicot	53
Endangered		
Milk-vetch, Ventura Marsh	Dicot	50
Endangered		
Millerbird, Nihoa	Bird	17
Endangered		
Mint, Garrett's	Dicot	8
Endangered		
Mint, Lakela's	Dicot	11
Endangered		
Mint, Longspurred	Dicot	12
Endangered		
Mint, Otay Mesa	Dicot	54
Endangered		
Mint, San Diego Mesa	Dicot	26
Endangered		
Mint, Scrub	Dicot	8
Endangered		
Mitracarpus Maxwelliae	Dicot	4
Endangered		
Mitracarpus Polycladus	Dicot	4
Endangered		
Monardella, Willowy	Dicot	26
Endangered		
Monkey-flower, Michigan	Dicot	59
Endangered		
Moorhen, Hawaiian Common	Bird	53
Endangered		
Morning-glory, Stebbins	Dicot	20
Endangered		
Moth, Blackburn's Sphinx	Insect	40
Endangered		
Mountain Beaver, Point Arena	Mammal	21
Endangered		
Mountainbalm, Indian Knob	Dicot	24
Endangered		
Mountain-mahogany, Catalina Island	Dicot	25
Endangered		
Mouse, Alabama Beach	Mammal	18
Endangered		
Mouse, Anastasia Island Beach	Mammal	8
Endangered		

Mouse, Choctawhatchee Beach Endangered	Mammal	18
Mouse, Key Largo Cotton Endangered	Mammal	1
Mouse, Pacific Pocket Endangered	Mammal	68
Mouse, Perdido Key Beach Endangered	Mammal	25
Mouse, Salt Marsh Harvest Endangered	Mammal	156
Mucket, Pink (Pearlymussel) Endangered	Bivalve	1006
Munroidendron racemosum (ncn) Endangered	Dicot	16
Mussel, Acornshell Southern Endangered	Bivalve	48
Mussel, Black (=Curtus' Mussel) Clubshell Endangered	Bivalve	13
Mussel, Clubshell Endangered	Bivalve	732
Mussel, Coosa Moccasinshell Endangered	Bivalve	74
Mussel, Cumberland Combshell Endangered	Bivalve	174
Mussel, Cumberland Elktoe Endangered	Bivalve	66
Mussel, Cumberland Pigtoe Endangered	Bivalve	25
Mussel, Dark Pigtoe Endangered	Bivalve	41
Mussel, Dwarf Wedge Endangered	Bivalve	566
Mussel, Fine-rayed Pigtoe Endangered	Bivalve	239
Mussel, Flat Pigtoe (=Marshall's Mussel) Endangered	Bivalve	10
Mussel, Gulf Moccasinshell Endangered	Bivalve	145
Mussel, Heavy Pigtoe (=Judge Tait's Mussel) Endangered	Bivalve	72
Mussel, Heelsplitter Carolina Endangered	Bivalve	116
Mussel, Ochlockonee Moccasinshell Endangered	Bivalve	21

1/28/2010 10:51:01 AM Ver. 2.10.4

Page 116 of 128

Mussel, Oval Pigtoe	Bivalve	184
Endangered		
Mussel, Ovate Clubshell	Bivalve	190
Endangered		
Mussel, Oyster	Bivalve	207
Endangered		
Mussel, Ring Pink (=Golf Stick Pearly)	Bivalve	416
Endangered		
Mussel, Rough Pigtoe	Bivalve	518
Endangered		
Mussel, Scaleshell	Bivalve	178
Endangered		
Mussel, Shiny Pigtoe	Bivalve	198
Endangered		
Mussel, Shiny-rayed Pocketbook	Bivalve	150
Endangered		
Mussel, Southern Clubshell	Bivalve	190
Endangered		
Mussel, Southern Pigtoe	Bivalve	98
Endangered		
Mussel, Speckled Pocketbook	Bivalve	7
Endangered		
Mussel, Winged Mapleleaf	Bivalve	105
Endangered		
Mustard, Carter's	Dicot	26
Endangered		
Mustard, Slender-petaled	Dicot	25
Endangered		
Myrcia Paganii	Dicot	4
Endangered		
Na'ena'e (Dubautia herbstobatae)	Dicot	17
Endangered		
Na'ena'e (Dubautia plantaginea ssp. humilis)	Dicot	20
Endangered		
Nani Wai'ale'ale (Viola kauensis var. wahiawaensis)	Dicot	16
Endangered		
Nanu (Gardenia mannii)	Dicot	17
Endangered		
Na'u (Gardenia brighamii)	Dicot	37
Endangered		
Naupaka, Dwarf (Scaevola coriacea)	Dicot	20
Endangered		
Navarretia, Few-flowered	Dicot	139
Endangered		
Navarretia, Many-flowered	Dicot	139
Endangered		
Nehe (Lipochaeta fauriei)	Dicot	16
Endangered		
Nehe (Lipochaeta kamolensis)	Dicot	20
Endangered		
Nehe (Lipochaeta lobata var. leptophylla)	Dicot	17
Endangered		
Nehe (Lipochaeta micrantha)	Dicot	16
Endangered		
Nehe (Lipochaeta tenuifolia)	Dicot	17
Endangered		
Nehe (Lipochaeta waimeaensis)	Dicot	16
Endangered		
Neraudia angulata (ncn)	Dicot	17
Endangered		
Neraudia ovata (ncn)	Dicot	20
Endangered		
Neraudia sericea (ncn)	Dicot	40
Endangered		
Nightjar, Puerto Rico	Bird	13
Endangered		
Nioi (Eugenia koolauensis)	Dicot	17
Endangered		
Niterwort, Amargosa	Dicot	20
Endangered		

Nohoanu (<i>Geranium multiflorum</i>)	Dicot	40
Endangered		
Nuku Pu'u	Bird	36
Endangered		
Ooclot	Mammal	214
Endangered		
'Oha (<i>Delissea rivularis</i>)	Dicot	16
Endangered		
'Oha (<i>Delissea subcordata</i>)	Dicot	17
Endangered		
'Oha (<i>Delissea undulata</i>)	Dicot	20
Endangered		
'Oha (<i>Lobelia gaudichaudii koolauensis</i>)	Dicot	17
Endangered		
'Oha Wai (<i>Clermontia drepanomorpha</i>)	Dicot	20
Endangered		
'Oha Wai (<i>Clermontia lindseyana</i>)	Dicot	40
Endangered		
'Oha Wai (<i>Clermontia oblongifolia</i> ssp. <i>brevipes</i>)	Dicot	20
Endangered		
'Oha Wai (<i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i>)	Dicot	20
Endangered		
'Oha Wai (<i>Clermontia peleana</i>)	Dicot	20
Endangered		
'Oha Wai (<i>Clermontia pyralaria</i>)	Dicot	20
Endangered		
'Oha Wai (<i>Clermontia samuelii</i>)	Dicot	20
Endangered		
'Ohai (<i>Sesbania tomentosa</i>)	Dicot	73
Endangered		
'Ohe'ohe (<i>Tetraplasandra gymnocarpa</i>)	Dicot	17
Endangered		
'Olulu (<i>Brighamia insignis</i>)	Dicot	16
Endangered		
Onion, Munz's	Monocot	28
Endangered		
'O'o, Kauai (= 'A'a)	Bird	16
Endangered		
Opuhe (<i>Urera kaalae</i>)	Dicot	17
Endangered		
'O'u (Honeycreeper)	Bird	36
Endangered		
Oxytheca, Cushenbury	Dicot	25
Endangered		

Paintbrush, San Clemente Island Indian	Dicot	25
Endangered		
Paintbrush, Soft-leaved	Dicot	25
Endangered		
Paintbrush, Tiburon	Dicot	58
Endangered		
Palila	Bird	20
Endangered		
Palo Colorado (<i>Ternstroemia luquillensis</i>)	Dicot	2
Endangered		
Palo de Jazmin	Dicot	2
Endangered		
Palo de Nigua	Dicot	11
Endangered		
Palo de Rosa	Dicot	7
Endangered		
Pamakani (<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>)	Dicot	17
Endangered		
Panicgrass, Carter's (<i>Panicum fauriei</i> var. <i>carteri</i>)	Monocot	37
Endangered		
Panther, Florida	Mammal	106
Endangered		
Parrot, Puerto Rican	Bird	2
Endangered		
Parrotbill, Maui	Bird	20
Endangered		
Pauoa (<i>Ctenitis squamigera</i>)	Ferns	37
Endangered		
Pawpaw, Beautiful	Dicot	25
Endangered		
Pawpaw, Four-petal	Dicot	28
Endangered		
Pawpaw, Rugel's	Dicot	11
Endangered		
Pearlymussel, Alabama Lamp	Bivalve	51
Endangered		
Pearlymussel, Appalachian Monkeyface	Bivalve	86
Endangered		
Pearlymussel, Birdwing	Bivalve	185
Endangered		
Pearlymussel, Cracking	Bivalve	220
Endangered		
Pearlymussel, Cumberland Bean	Bivalve	215
Endangered		
Pearlymussel, Cumberland Monkeyface	Bivalve	166
Endangered		
Pearlymussel, Curtis'	Bivalve	18
Endangered		
Pearlymussel, Dromedary	Bivalve	255
Endangered		
Pearlymussel, Fat Pocketbook	Bivalve	379
Endangered		
Pearlymussel, Green-blossom	Bivalve	114
Endangered		
Pearlymussel, Higgins' Eye	Bivalve	514
Endangered		
Pearlymussel, Little-wing	Bivalve	211
Endangered		
Pearlymussel, Orange-footed	Bivalve	440
Endangered		
Pearlymussel, Pale Lilliput	Bivalve	69
Endangered		
Pearlymussel, Purple Cat's Paw	Bivalve	129
Endangered		
Pearlymussel, Tubercled-blossom	Bivalve	347
Endangered		
Pearlymussel, Turgid-blossom	Bivalve	89
Endangered		
Pearlymussel, White Cat's Paw	Bivalve	34
Endangered		

Pearlymussel, White Wartyback
 Endangered
 Pearlymussel, Yellow-blossom
 Endangered
 Pebblesnail, Flat
 Endangered
 Pelos del Diablo
 Endangered
 Penny-cress, Kneeland Prairie
 Endangered
 Pennyroyal, Todsens
 Endangered
 Penstemon, Blowout
 Endangered
 Pentachaeta, Lyon's
 Endangered
 Pentachaeta, White-rayed
 Endangered
 Peperomia, Wheeler's
 Endangered
 Petrel, Hawaiian Dark-rumped
 Endangered
 Phacelia, Clay
 Endangered
 Phacelia, Island
 Endangered
 Phlox, Texas Trailing
 Endangered
 Phlox, Yreka
 Endangered
 Phyllostegia hirsuta (ncn)
 Endangered
 Phyllostegia kaalaensis (ncn)
 Endangered
 Phyllostegia knudsenii (ncn)
 Endangered
 Phyllostegia mannii (ncn)
 Endangered
 Phyllostegia mollis (ncn)
 Endangered
 Phyllostegia parviflora (ncn)
 Endangered
 Phyllostegia velutina (ncn)
 Endangered

Bivalve	246
Bivalve	177
Gastropod	20
Monocot	7
Dicot	20
Dicot	27
Dicot	12
Dicot	50
Dicot	58
Dicot	2
Bird	56
Dicot	16
Dicot	25
Dicot	37
Dicot	20
Dicot	17
Dicot	17
Dicot	16
Dicot	20
Dicot	37
Dicot	17
Dicot	20

Phyllostegia waimeae (ncn)	Dicot	16
Endangered		
Phyllostegia warshaueri (ncn)	Dicot	20
Endangered		
Phyllostegia wawrana (ncn)	Dicot	16
Endangered		
Pigeon, Puerto Rican Plain	Bird	9
Endangered		
Pilo (Hedyotis mannii)	Dicot	20
Endangered		
Pinkroot, Gentian	Dicot	17
Endangered		
Piperia, Yadon's	Monocot	27
Endangered		
Pitaya, Davis' Green	Dicot	15
Endangered		
Pitcher-plant, Alabama Canebrake	Dicot	38
Endangered		
Pitcher-plant, Green	Dicot	106
Endangered		
Pitcher-plant, Mountain Sweet	Dicot	52
Endangered		
Platanthera holochila (ncn)	Monocot	36
Endangered		
Plover, Piping	Bird	2476
Endangered		
Plum, Scrub	Dicot	36
Endangered		
Poa siphonoglossa (ncn)	Monocot	16
Endangered		
Po'e (Portulaca sclerocarpa)	Dicot	40
Endangered		
Polygala, Lewton's	Dicot	48
Endangered		
Polygala, Tiny	Dicot	51
Endangered		
Polygonum, Scott's Valley	Dicot	22
Endangered		
Polystichum calderonense (ncn)	Ferns	2
Endangered		
Pondberry	Dicot	190
Endangered		
Pondweed, Little Aguja Creek	Monocot	2
Endangered		
Po'ouli	Bird	20
Endangered		
Popcornflower, Rough	Dicot	19
Endangered		
Popolo 'Aiakeakua (Solanum sandwicense)	Dicot	33
Endangered		
Popolo Ku Mai (Solanum incompletum)	Dicot	20
Endangered		
Poppy, Sacramento Prickly	Dicot	13
Endangered		
Poppy-mallow, Texas	Dicot	16
Endangered		
Potentilla, Hickman's	Dicot	46
Endangered		
Prairie-chicken, Attwater's Greater	Bird	70
Endangered		
Prickly-apple, Fragrant	Dicot	4
Endangered		
Prickly-ash, St. Thomas	Dicot	4
Endangered		
Pronghorn, Sonoran	Mammal	59
Endangered		
Pseudoscorpion, Tooth Cave	Arachnid	13
Endangered		
Pteris lidgatei (ncn)	Ferns	37
Endangered		

Pua'ala (<i>Brighamia rockii</i>)	Dicot	20
Endangered		
Purple Bean	Bivalve	109
Endangered		
Pu'uka'a (<i>Cyperus trachysanthos</i>)	Monocot	33
Endangered		
Pygmy-owl, Cactus Ferruginous	Bird	130
Endangered		
Quillwort, Black-spored	Ferns	54
Endangered		
Quillwort, Louisiana	Ferns	124
Endangered		
Quillwort, Mat-forming	Ferns	33
Endangered		
Rabbit, Lower Keys Marsh	Mammal	1
Endangered		
Rabbit, Pygmy	Mammal	79
Endangered		
Rabbit, Riparian Brush	Mammal	24
Endangered		
Rabbitsfoot, Rough	Bivalve	64
Endangered		
Rail, California Clapper	Bird	207
Endangered		
Rail, Light-footed Clapper	Bird	118
Endangered		
Rail, Yuma Clapper	Bird	172
Endangered		
Rattleweed, Hairy	Dicot	22
Endangered		
Reed-mustard, Barneby	Dicot	11
Endangered		
Reed-mustard, Shrubby	Dicot	23
Endangered		
<i>Remya kauaiensis</i> (ncn)	Dicot	16
Endangered		
<i>Remya montgomeryi</i> (ncn)	Dicot	16
Endangered		
<i>Remya</i> , Maui	Dicot	20
Endangered		
<i>Rhadine exilis</i> (ncn)	Insect	17
Endangered		
<i>Rhadine infernalis</i> (ncn)	Insect	17
Endangered		

Rhododendron, Chapman	Dicot	20
Endangered		
Rice Rat (=Silver Rice Rat)	Mammal	1
Endangered		
Ridge-cress (=Pepper-cress), Barneby	Dicot	13
Endangered		
Riffleshell, Northern	Bivalve	362
Endangered		
Riffleshell, Tan	Bivalve	254
Endangered		
Riversnail, Anthony's	Gastropod	66
Endangered		
Rock-cress, Hoffmann's	Dicot	25
Endangered		
Rock-cress, Large (=Braun's)	Dicot	67
Endangered		
Rock-cress, McDonald's	Dicot	21
Endangered		
Rock-cress, Santa Cruz Island	Dicot	25
Endangered		
Rock-cress, Shale Barren	Dicot	81
Endangered		
Rock-cress, Small	Dicot	46
Endangered		
Rock-pocketbook, Ouachita (=Wheeler's pm)	Bivalve	38
Endangered		
Rocksail, Plicate	Gastropod	25
Endangered		
Rosemary, Etonia	Dicot	8
Endangered		
Rosemary, Short-leaved	Dicot	19
Endangered		
Rush-pea, Slender	Dicot	6
Endangered		
Salamander, Barton Springs	Amphibian	28
Endangered		
Salamander, California Tiger	Amphibian	480
Endangered		
Salamander, Desert Slender	Amphibian	28
Endangered		
Salamander, Santa Cruz Long-toed	Amphibian	49
Endangered		
Salamander, Shenandoah	Amphibian	39
Endangered		
Salamander, Sonora Tiger	Amphibian	26
Endangered		
Salamander, Texas Blind	Amphibian	75
Endangered		
Sandalwood, Lanai (=Iliahi)	Dicot	20
Endangered		
Sandlace	Dicot	36
Endangered		
Sand-verbena, Large-fruited	Dicot	38
Endangered		
Sandwort, Cumberland	Dicot	36
Endangered		
Sandwort, Marsh	Dicot	24
Endangered		
Sanicula maritima (ncn)	Dicot	17
Endangered		
Sanicula purpurea (ncn)	Dicot	20
Endangered		
Schiedea haleakalensis (ncn)	Dicot	20
Endangered		
Schiedea helleri (ncn)	Dicot	16
Endangered		
Schiedea hookeri (ncn)	Dicot	17
Endangered		
Schiedea kaalae (ncn)	Dicot	17
Endangered		

Schiedea kauaiensis (ncn)	Dicot	16
Endangered		
Schiedea lydgatei (ncn)	Dicot	20
Endangered		
Schiedea membranacea (ncn)	Dicot	16
Endangered		
Schiedea nuttallii (ncn)	Dicot	33
Endangered		
Schiedea sarmentosa (ncn)	Dicot	20
Endangered		
Schiedea spargulina var. leiopoda (ncn)	Dicot	16
Endangered		
Schiedea verticillata (ncn)	Dicot	17
Endangered		
Schiedea, Diamond Head (Schiedea adamantis)	Dicot	17
Endangered		
Sea turtle, green	Reptile	624
Endangered		
Sea turtle, hawksbill	Reptile	364
Endangered		
Sea turtle, Kemp's ridley	Reptile	323
Endangered		
Sea turtle, leatherback	Reptile	566
Endangered		
Sea-blite, California	Dicot	24
Endangered		
Seal, Caribbean Monk	Marine mml	2
Endangered		
Seal, Hawaiian Monk	Marine mml	74
Endangered		
Sedge, Golden	Monocot	20
Endangered		
Sedge, White	Monocot	27
Endangered		
Sheep, Peninsular Bighorn	Mammal	92
Endangered		
Sheep, Sierra Nevada Bighorn	Mammal	43
Endangered		
Shrew, Buena Vista Lake Ornate	Mammal	25
Endangered		
Shrike, San Clemente Loggerhead	Bird	25
Endangered		
Shrimp, Alabama Cave	Crustacean	15
Endangered		

1/28/2010 10:51:46 AM Ver. 2.10.4

Page 120 of 128

Shrimp, California Freshwater	Crustacean	65
Endangered		
Shrimp, Kentucky Cave	Crustacean	31
Endangered		
Silene alexandri (ncn)	Dicot	20
Endangered		
Silene lanceolata (ncn)	Dicot	73
Endangered		
Silene perlmanii (ncn)	Dicot	17
Endangered		
Silversword, Ka'u (Argyroxiphium kauense)	Dicot	20
Endangered		
Silversword, Mauna Kea ('Ahinahina)	Dicot	40
Endangered		
Skipper, Carson Wandering	Insect	19
Endangered		
Skipper, Laguna Mountain	Insect	26
Endangered		
Snail, Armored	Gastropod	14
Endangered		
Snail, Iowa Pleistocene	Gastropod	78
Endangered		
Snail, Lioplax Cylindrical	Gastropod	20
Endangered		
Snail, Morro Shoulderband	Gastropod	24
Endangered		
Snail, O'ahu Tree (Achatinella abbreviata)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella apexfulva)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella bellula)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella buddii)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella bulimoides)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella byronii)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella caesia)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella casta)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella cestus)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella concavospira)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella curta)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella decipiens)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella decora)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella dimorpha)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella elegans)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella fulgens)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella fuscobasis)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella juddii)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella juncea)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella lehuiensis)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella leucorraphe)	Gastropod	17
Endangered		
Snail, O'ahu Tree (Achatinella lila)	Gastropod	17
Endangered		

Snail, O'ahu Tree (<i>Achatinella livida</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella lorata</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella mustelina</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella papyracea</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella phaeozona</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella pulcherrima</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella pupukanioe</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella rosea</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella sowerbyana</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella spaldingi</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella stewartii</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella swiftii</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella taeniolata</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella thaanumi</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella turgida</i>)	Gastropod	17
Endangered		
Snail, O'ahu Tree (<i>Achatinella valida</i>)	Gastropod	17
Endangered		
Snail, Pecos Assiminea	Gastropod	15
Endangered		
Snail, Snake River Physa	Gastropod	52
Endangered		
Snail, Tulotoma	Gastropod	63
Endangered		
Snail, Utah Valvata	Gastropod	35
Endangered		
Snail, Virginia Fringed Mountain	Gastropod	7
Endangered		
Snake, San Francisco Garter	Reptile	41
Endangered		

Snakeroot	Dicot	28
Endangered		
Snowbells, Texas	Dicot	40
Endangered		
Sparrow, Cape Sable Seaside	Bird	27
Endangered		
Sparrow, Florida Grasshopper	Bird	46
Endangered		
Spermolepis hawaiiensis (ncn)	Dicot	73
Endangered		
Spider, Government Canyon Cave	Arachnid	17
Endangered		
Spider, Kauai Cave Wolf	Arachnid	16
Endangered		
Spider, Madla's Cave	Arachnid	17
Endangered		
Spider, Robber Baron Cave	Arachnid	17
Endangered		
Spider, Spruce-fir Moss	Arachnid	67
Endangered		
Spider, Tooth Cave	Arachnid	13
Endangered		
Spider, Vesper Cave	Arachnid	17
Endangered		
Spineflower, Ben Lomond	Dicot	22
Endangered		
Spineflower, Howell's	Dicot	21
Endangered		
Spineflower, Orcutt's	Dicot	43
Endangered		
Spineflower, Robust	Dicot	49
Endangered		
Spineflower, Scotts Valley	Dicot	22
Endangered		
Spineflower, Slender-horned	Dicot	104
Endangered		
Spineflower, Sonoma	Dicot	44
Endangered		
Spiny mussel, James River	Bivalve	176
Endangered		
Spiny mussel, Tar River	Bivalve	90
Endangered		
Springsnail, Alamosa	Gastropod	11
Endangered		
Springsnail, Bruneau Hot	Gastropod	9
Endangered		
Springsnail, Koster's	Gastropod	4
Endangered		
Springsnail, Roswell	Gastropod	4
Endangered		
Springsnail, Socorro	Gastropod	11
Endangered		
Spurge, Deltoid	Dicot	15
Endangered		
Squirrel, Carolina Northern Flying	Mammal	132
Endangered		
Squirrel, Delmarva Peninsula Fox	Mammal	178
Endangered		
Squirrel, Mount Graham Red	Mammal	14
Endangered		
Stenogyne angustifolia (ncn)	Dicot	20
Endangered		
Stenogyne bifida (ncn)	Dicot	20
Endangered		
Stenogyne campanulata (ncn)	Dicot	16
Endangered		
Stenogyne kanehoana (ncn)	Dicot	17
Endangered		
Stickseed, Showy	Dicot	10
Endangered		

Stickyseed, Baker's Endangered	Dicot	27
Stilt, Hawaiian (=Ac'o) Endangered	Bird	73
Stirrupshell Endangered	Bivalve	28
Stonecrop, Lake County Endangered	Dicot	139
Stork, Wood Endangered	Bird	1428
Sumac, Michaux's Endangered	Dicot	292
Sunflower, San Mateo Woolly Endangered	Dicot	19
Sunflower, Schweinitz's Endangered	Dicot	193
Tadpole Shrimp, Vernal Pool Endangered	Crustacean	484
Taraxacum, California Endangered	Dicot	25
Tarplant, Gaviota Endangered	Dicot	25
Tectaria Estremerana Endangered	Ferns	2
Tern, California Least Endangered	Bird	239
Tern, Interior (population) Least Endangered	Bird	1622
Tern, Roseate Endangered	Bird	208
Ternstroemia subsessilis (ncn) Endangered	Dicot	2
Tetramolopium arenarium (ncn) Endangered	Dicot	20
Tetramolopium capillare (ncn) Endangered	Dicot	20
Tetramolopium filiforme (ncn) Endangered	Dicot	17
Tetramolopium lepidotum ssp. lepidotum (ncn) Endangered	Dicot	17
Tetramolopium remyi (ncn) Endangered	Dicot	20
Thistle, Chorro creek Bog Endangered	Dicot	24

Thistle, Fountain Endangered	Dicot	64
Thistle, La Graciosa Endangered	Dicot	49
Thistle, Suisun Endangered	Dicot	23
Thormint, San Mateo Endangered	Dicot	19
Threeidge, Fat (Mussel) Endangered	Bivalve	31
Thrush, Large Kauai Endangered	Bird	16
Thrush, Molokai (Oloma'o) Endangered	Bird	20
Thrush, Small Kauai (Puaiohi) Endangered	Bird	16
Toad, Arroyo Southwestern Endangered	Amphibian	219
Toad, Houston Endangered	Amphibian	157
Torreya, Florida Endangered	Conf/cycds	30
Tree Fern, Elfin Endangered	Ferns	2
Trematolobelia singularis (ncn) Endangered	Dicot	17
Trillium, Persistent Endangered	Monocot	32
Trillium, Relict Endangered	Monocot	89
Tuctoria, Green's Endangered	Dicot	202
Turtle, Alabama Red-bellied Endangered	Reptile	33
Turtle, Plymouth Red-bellied Endangered	Reptile	17
Uhiuhi (Caesalpinia kavaensis) Endangered	Dicot	57
Ulihi (Phyllostegia glabra var. lanaiensis) Endangered	Dicot	20
Umbel, Huachuca Water Endangered	Dicot	44
Uvillo Endangered	Dicot	6
Vernonia Proctorii (ncn) Endangered	Dicot	3
Vetch, Hawaiian (Vicia menziesii) Endangered	Dicot	20
Vigna o-wahuensis (ncn) Endangered	Dicot	56
Viola helenae (ncn) Endangered	Dicot	16
Viola lanaiensis (ncn) Endangered	Dicot	20
Viola oahuensis (ncn) Endangered	Dicot	17
Vireo, Black-capped Endangered	Bird	590
Vireo, Least Bell's Endangered	Bird	253
Vole, Amargosa Endangered	Mammal	31
Vole, Florida Salt Marsh Endangered	Mammal	8
Vole, Hualapai Mexican Endangered	Mammal	23
Wahane (Pritchardia aylmer-robinsonii) Endangered	Monocot	16
Wahine Noho Kula (Isodendron pyrifolium) Endangered	Dicot	20

Wallflower, Ben Lomond Endangered	Dicot	22
Wallflower, Contra Costa Endangered	Dicot	18
Wallflower, Menzie's Endangered	Dicot	76
Walnut, Nogal Endangered	Dicot	2
Warbler (=Wood), Golden-cheeked Endangered	Bird	359
Warbler (=Wood), Kirtland's Endangered	Bird	229
Warbler, Bachman's Endangered	Bird	50
Warea, Wide-leaf Endangered	Dicot	28
Watercress, Gambel's Endangered	Dicot	125
Water-willow, Cooley's Endangered	Dicot	7
Wawac'lole (Phlegmariurus (=Huperzia) mannii) Endangered	Ferns	40
Wawac'lole (Phlegmariurus (=Lycopodium) nutans) Endangered	Ferns	17
Whale, Finback Endangered	Marine mml	87
Whale, Humpback Endangered	Marine mml	90
Whale, northern right Endangered	Marine mml	18
Wild-buckwheat, Clay-loving Endangered	Dicot	29
Wild-rice, Texas Endangered	Monocot	75
Wire-lettuce, Malheur Endangered	Dicot	1
Wireweed Endangered	Dicot	19
Woodland-star, San Clemente Island Endangered	Dicot	25
Woodpecker, Ivory-billed Endangered	Bird	35
Woodpecker, Red-cockaded Endangered	Bird	3401

Woodrat, Key Largo	Mammal	1
Endangered		
Woodrat, Riparian	Mammal	24
Endangered		
Woolly-star, Santa Ana River	Dicot	70
Endangered		
Woolly-threads, San Joaquin	Dicot	169
Endangered		
Xylosma crenatum (ncn)	Dicot	16
Endangered		
Yerba Santa, Lompoc	Dicot	25
Endangered		
Ziziphus, Florida	Dicot	19
Endangered		
Adobe Sunburst, San Joaquin	Dicot	76
Threatened		
Amaranth, Seabeach	Dicot	171
Threatened		
Amole, Cammatita Canyon	Monocot	24
Threatened		
Amole, Purple	Monocot	51
Threatened		
Amphianthus, Little	Dicot	174
Threatened		
Aster, Decurrent False	Dicot	301
Threatened		
Aupaka (Isodendron longifolium)	Dicot	33
Threatened		
Baccharis, Encinitas	Dicot	43
Threatened		
Bankclimber, Purple	Bivalve	126
Threatened		
Barbara Buttons, Mohr's	Dicot	66
Threatened		
Beaked-rush, Knieskern's	Monocot	93
Threatened		
Bear, Grizzly	Mammal	371
Threatened		
Bear, Louisiana Black	Mammal	679
Threatened		
Beetle, Delta Green Ground	Insect	23
Threatened		
Beetle, Northeastern Beach Tiger	Insect	119
Threatened		
Beetle, Puritan Tiger	Insect	73
Threatened		
Beetle, Valley Elderberry Longhorn	Insect	377
Threatened		
Birch, Virginia Round-leaf	Dicot	10
Threatened		
Birds-in-a-nest, White	Dicot	6
Threatened		
Bladderpod, Dudley Bluffs	Dicot	1
Threatened		
Bladderpod, Lyrate	Dicot	31
Threatened		
Bladderpod, Missouri	Dicot	66
Threatened		
Blazing Star, Ash Meadows	Dicot	14
Threatened		
Blazing Star, Heller's	Dicot	51
Threatened		
Bluecurls, Hidden Lake	Dicot	25
Threatened		
Boa, Mona	Reptile	2
Threatened		
Bonamia, Florida	Dicot	82
Threatened		
Brodiaea, Chinese Camp	Monocot	9
Threatened		

Brodiaea, Thread-leaved Threatened	Monocot	121
Buckwheat, Scrub Threatened	Dicot	56
Buckwheat, Southern Mountain Wild Threatened	Dicot	25
Butterfly Plant, Colorado Threatened	Dicot	16
Butterfly, Bay Checkerspot (Wright's euphydryas) Threatened	Insect	68
Butterfly, Oregon Silverspot Threatened	Insect	68
Butterweed, Layne's Threatened	Dicot	29
Butterwort, Godfrey's Threatened	Dicot	6
Cactus, Bunched Cory Threatened	Dicot	15
Cactus, Chisos Mountain Hedgehog Threatened	Dicot	15
Cactus, Cochise Pincushion Threatened	Dicot	19
Cactus, Lee Pincushion Threatened	Dicot	8
Cactus, Lloyd's Mariposa Threatened	Dicot	21
Cactus, Mesa Verde Threatened	Dicot	29
Cactus, Siler Pincushion Threatened	Dicot	39
Cactus, Uinta Basin Hookless Threatened	Dicot	86
Cactus, Winkler Threatened	Dicot	11
Caracara, Audubon's Crested Threatened	Bird	146
Catchfly, Spalding's Threatened	Dicot	119
Ceanothus, Vail Lake Threatened	Dicot	28
Centaury, Spring-loving Threatened	Dicot	20
Checker-mallow, Nelson's Threatened	Dicot	169

Chumbo, Higo	Dicot	2
Threatened		
Clarkia, Springville	Dicot	24
Threatened		
Clover, Fleshy Owl's	Dicot	149
Threatened		
Clover, Prairie Bush	Dicot	1054
Threatened		
Cobana Negra	Dicot	5
Threatened		
Coqui, Golden	Amphibian	2
Threatened		
Crocodile, American	Reptile	46
Threatened		
Crownbeard, Big-leaved	Dicot	43
Threatened		
Cycladenia, Jones	Dicot	38
Threatened		
Cypress, Gowen	Conf/cyds	27
Threatened		
Daisy, Lakeside	Dicot	59
Threatened		
Daisy, Maguire	Dicot	11
Threatened		
Daisy, Parish's	Dicot	53
Threatened		
Dudleya, Conejo	Dicot	25
Threatened		
Dudleya, Marcescent	Dicot	67
Threatened		
Dudleya, Santa Cruz Island	Dicot	25
Threatened		
Dudleya, Santa Monica Mountains	Dicot	67
Threatened		
Dudleya, Verity's	Dicot	25
Threatened		
Dwarf-flax, Marin	Dicot	17
Threatened		
Eagle, Bald	Bird	115
Threatened		
Elimia, Lacy	Gastropod	11
Threatened		
Evening-primrose, San Benito	Dicot	23
Threatened		
Fairy Shrimp, Vernal Pool	Crustacean	615
Threatened		
Fatmucket, Arkansas	Bivalve	33
Threatened		
Fern, Alabama Streak-sorus	Ferns	10
Threatened		
Fern, American hart's-tongue	Ferns	114
Threatened		
Fleabane, Zuni	Dicot	27
Threatened		
Four-o'clock, Macfarlane's	Dicot	22
Threatened		
Frog, California Red-legged	Amphibian	478
Threatened		
Frog, Chiricahua Leopard	Amphibian	174
Threatened		
Fruit, Earth (=geocarpon)	Dicot	115
Threatened		
Gesneria pauciflora (n:n)	Dicot	3
Threatened		
Gnatcatcher, Coastal California	Bird	146
Threatened		
Goldenrod, Blue Ridge	Dicot	35
Threatened		
Goldenrod, Houghton's	Dicot	78
Threatened		

Goldenrod, White-haired Threatened	Dicot	20
Gooseberry, Miccosukee Threatened	Dicot	29
Grass, Colusa Threatened	Monocot	139
Grass, San Joaquin Valley Orcutt Threatened	Monocot	185
Grass, Slender Orcutt Threatened	Dicot	156
Groundsel, San Francisco Peaks Threatened	Dicot	9
Guajon Threatened	Amphibian	4
Gumplant, Ash Meadows Threatened	Dicot	20
Haha (Cyanea recta) Threatened	Dicot	16
Ha'Iwale (Cyrtandra limahuliensis) Threatened	Dicot	16
Heartleaf, Dwarf-flowered Threatened	Dicot	119
Heather, Mountain Golden Threatened	Dicot	13
Howellia, Water Threatened	Dicot	108
Iguana, Mona Ground Threatened	Reptile	2
Iris, Dwarf Lake Threatened	Monocot	113
Isopod, Madison Cave Threatened	Crustacean	34
Ivesia, Ash Meadows Threatened	Dicot	20
Joint-vetch, Sensitive Threatened	Dicot	273
Kolea (Myrsine linearifolia) Threatened	Dicot	16
Ladies'-tresses, Ute Threatened	Monocot	142
Liveforever, Laguna Beach Threatened	Dicot	17
Lizard, Coachella Valley Fringe-toed Threatened	Reptile	28

Lizard, Island Night Threatened	Reptile	75
Locoweed, Fassett's Threatened	Dicot	31
Lupine, Kincaid's Threatened	Dicot	139
Lynx, Canada Threatened	Mammal	286
Makou (Peucedanum sandwicense) Threatened	Dicot	53
Manaca, palma de Threatened	Monocot	11
Manzanita, Ione Threatened	Dicot	31
Manzanita, Morro Threatened	Dicot	24
Manzanita, Pallid Threatened	Dicot	29
Milk-vetch, Ash Meadows Threatened	Dicot	14
Milk-vetch, Deseret Threatened	Dicot	16
Milk-vetch, Fish Slough Threatened	Dicot	17
Milk-vetch, Heliotrope Threatened	Dicot	16
Milk-vetch, Picrson's Threatened	Dicot	21
Milkweed, Mead's Threatened	Dicot	315
Milkweed, Welsh's Threatened	Dicot	12
Monkshood, Northern Wild Threatened	Dicot	217
Moth, Kern Primrose Sphinx Threatened	Insect	25
Mouse, Preble's Meadow Jumping Threatened	Mammal	92
Mouse, Southeastern Beach Threatened	Mammal	18
Mucket, Orangenacre Threatened	Bivalve	107
Murrelet, Marbled Threatened	Bird	643
Mussel, Alabama Moccasinshell Threatened	Bivalve	142
Mussel, Fine-lined Pocketbook Threatened	Bivalve	269
Mussel, Heelsplitter Inflated Threatened	Bivalve	131
Naucorid, Ash Meadows Threatened	Insect	14
Navarretia, Spreading Threatened	Dicot	79
Oak, Hinckley Threatened	Dicot	21
Orchid, Eastern Prairie Fringed Threatened	Monocot	822
Orchid, Western Prairie Fringed Threatened	Monocot	1161
Otter, Northern Sea Threatened	Marine mml	3
Otter, Southern Sea Threatened	Marine mml	73
Owl, Mexican Spotted Threatened	Bird	591
Owl, Northern Spotted Threatened	Bird	893
Paintbrush, Ash-grey Indian Threatened	Dicot	25

Paintbrush, Golden Threatened	Dicot	47
Pearlshell, Louisiana Threatened	Bivalve	18
Pink, Swamp Threatened	Monocot	454
Plover, Western Snowy Threatened	Bird	445
Pogonia, Small Whorled Threatened	Monocot	994
Potato-bean, Price's Threatened	Dicot	194
Prairie Dog, Utah Threatened	Mammal	46
Primrose, Maguire Threatened	Dicot	9
Pussypaws, Mariposa Threatened	Dicot	68
Rattlesnake, New Mexican Ridge-nosed Threatened	Reptile	26
Reed-mustard, Clay Threatened	Dicot	10
Rocksnail, Painted Threatened	Gastropod	50
Rocksnail, Round Threatened	Gastropod	20
Rosemary, Cumberland Threatened	Dicot	61
Roseroot, Leedy's Threatened	Dicot	63
Rush-rose, Island Threatened	Dicot	25
Salamander, Cheat Mountain Threatened	Amphibian	32
Salamander, Flatwoods Threatened	Amphibian	236
Salamander, Red Hills Threatened	Amphibian	38
Salamander, San Marcos Threatened	Amphibian	75
Sandwort, Bear Valley Threatened	Dicot	25
Schiedea spergulina var. spergulina (ncn) Threatened	Dicot	16

1/28/2010 10:52:53 AM Ver. 2.10.4

Page 126 of 128

Schoepfia arenaria (ncn)	Dicot	4
Threatened		
Scrub-Jay, Florida	Bird	310
Threatened		
Sea turtle, loggerhead	Reptile	816
Threatened		
Sea turtle, olive ridley	Reptile	96
Threatened		
Seagrass, Johnson's	Monocot	51
Threatened		
Seal, Guadalupe Fur	Marine mml	27
Threatened		
Sea-lion, Steller (eastern)	Marine mml	5
Threatened		
Sedge, Navajo	Monocot	28
Threatened		
Shagreen, Magazine Mountain	Gastropod	13
Threatened		
Shearwater, Newell's Townsend's	Bird	53
Threatened		
Shrimp, Squirrel Chimney Cave	Crustacean	19
Threatened		
Silene hawaiiensis (ncn)	Dicot	20
Threatened		
Silversword, Haleakala ('Ahinahina)	Dicot	40
Threatened		
Skink, Blue-tailed Mole	Reptile	25
Threatened		
Skink, Sand	Reptile	56
Threatened		
Skipper, Pawnee Montane	Insect	22
Threatened		
Skullcap, Large-flowered	Dicot	58
Threatened		
Slabshell, Chipola	Bivalve	17
Threatened		
Snail, Bliss Rapids	Gastropod	43
Threatened		
Snail, Chittenango Ovate Amber	Gastropod	17
Threatened		
Snail, Flat-spined Three-toothed	Gastropod	22
Threatened		
Snail, Newcomb's	Gastropod	16
Threatened		
Snail, Noonday	Gastropod	4
Threatened		
Snail, Painted Snake Coiled Forest	Gastropod	3
Threatened		
Snail, Stock Island Tree	Gastropod	1
Threatened		
Snake, Atlantic Salt Marsh	Reptile	25
Threatened		
Snake, Concho Water	Reptile	78
Threatened		
Snake, Eastern Indigo	Reptile	1251
Threatened		
Snake, Giant Garter	Reptile	208
Threatened		
Snake, Lake Erie Water	Reptile	27
Threatened		
Snake, Northern Copperbelly Water	Reptile	136
Threatened		
Sneezeweed, Virginia	Dicot	59
Threatened		
Sparrow, San Clemente Sage	Bird	25
Threatened		
Spineflower, Monterey	Dicot	49
Threatened		
Spiraea, Virginia	Dicot	372
Threatened		

Spurge, Garber's Threatened	Dicot	15
Spurge, Hoover's Threatened	Dicot	165
Spurge, Telephus Threatened	Dicot	6
Squirrel, Northern Idaho Ground Threatened	Mammal	11
Staghorn coral Threatened	Coral	1
Sunflower, Pecos Threatened	Dicot	37
Sunray, Ash Meadows Threatened	Dicot	14
Tarplant, Otay Threatened	Dicot	26
Tarplant, Santa Cruz Threatened	Dicot	67
Tetramolopium rockii (ncn) Threatened	Dicot	20
Thelypody, Howell's Spectacular Threatened	Dicot	24
Thistle, Pitcher's Threatened	Dicot	355
Thistle, Sacramento Mountains Threatened	Dicot	13
Thornmint, San Diego Threatened	Dicot	26
Toad, Puerto Rican Crested Threatened	Amphibian	8
Tortoise, Desert Threatened	Reptile	205
Tortoise, Gopher Threatened	Reptile	179
Towhee, Inyo Brown Threatened	Bird	6
Townsendia, Last Chance Threatened	Dicot	19
Turtle, Bog (Northern population) Threatened	Reptile	631
Turtle, Flattened Musk Threatened	Reptile	97
Turtle, Ringed Sawback Threatened	Reptile	104

Turtle, Yellow-blotched Map Threatened	Reptile	68
Twinpod, Dudley Bluffs Threatened	Dicot	1
Vervain, California Threatened	Dicot	9
Water-plantain, Kral's Threatened	Monocot	34
Whipsnake (=Striped Racer), Alameda Threatened	Reptile	29
Whitlow-wort, Papery Threatened	Dicot	44
Wild-buckwheat, Gypsum Threatened	Dicot	11
Wings, Pigeon Threatened	Dicot	30
Yellowhead, Desert Threatened	Dicot	6

No species were selected for exclusion.

Dispersed species included in report. 1/28/2010 10:53:15 AM Ver. 2.10.4 Page 128 of 128

Species in Counties by State and Taxa

No species were excluded

Minimum of 1 Acre

All Medium Types Reported

Amphibian, Reptile, Crustacean, Bivalve, Gastropod, Arachnid, Insect, Dicot, Monocot, Ferns

root celery (PR)

AL, AK, AZ, AR, CA, CO, CT, DE, DC, FL, GA, HI, ID, IL, IN, IA, KS, KY, LA, ME, MD,
MA,
MI, MN, MS, MO, MT, NE, NV, NH, NJ, NM, NY, NC, ND, OH, OK, OR, PA, PR, RI, SC, SD,
TN, TX, UT, VT, VA, WA, WV, WI, WY

28 Species Affected:

Inverse Name:	Taxa:	Co. occurrence:
Status:		
Bariaco	Dicot	2
Endangered		
Boa, Puerto Rican	Reptile	4
Endangered		
Capa Rosa	Dicot	1
Endangered		
Chupacallos	Dicot	1
Endangered		
Erubia	Dicot	1
Endangered		
Fern, Elaphoglossum serpens	Ferns	1
Endangered		
Fern, Thelypteris inabonensis	Ferns	2
Endangered		

Fern, Thelypteris yaucoensis	Ferns	2
Endangered		
Higuero De Sierra	Dicot	1
Endangered		
Holly, Cook's	Dicot	2
Endangered		
Ilex sintenisii (ncn)	Dicot	1
Endangered		
Lepanthes eltoensis (ncn)	Monocot	1
Endangered		
Palo Colorado (Ternstroemia luquillensis)	Dicot	1
Endangered		
Palo de Jazmin	Dicot	1
Endangered		
Palo de Nigua	Dicot	4
Endangered		
Palo de Rosa	Dicot	1
Endangered		
Prickly-ash, St. Thomas	Dicot	1
Endangered		
Sea turtle, green	Reptile	2
Endangered		
Sea turtle, hawksbill	Reptile	2
Endangered		
Sea turtle, leatherback	Reptile	3
Endangered		
Tree Fern, Elfin	Ferns	1
Endangered		
Uvillo	Dicot	2
Endangered		
Walnut, Nogal	Dicot	1
Endangered		
Cobana Negra	Dicot	1
Threatened		
Coqui, Golden	Amphibian	1
Threatened		
Guajon	Amphibian	1
Threatened		
Manaca, palma de	Monocot	1
Threatened		
Toad, Puerto Rican Crested	Amphibian	1
Threatened		

No species were selected for exclusion.

Dispersed species included in report.

Species in Counties by State and Taxa

No species were excluded

Minimum of 1 Acre

Freshwater

Fish

apples, citrus fruit, all, cotton, all, grapes, potatoes, cantaloups, cucumbers and pickles, honeydew melons, pumpkins, squash, watermelons, eggplant, peppers, bell, peppers, chile

(all peppers - excluding bell), pimientos, tomatoes, amaranth, celery, lettuce, all, escarole

and endive, lettuce, head, lettuce, leaf, lettuce, romaine, parsley, rhubarb, spinach, root celery (PR)

AL, AK, AZ, AR, CA, CO, CT, DE, DC, FL, GA, HI, ID, IL, IN, IA, KS, KY, LA, ME, MD, MA,

MI, MN, MS, MO, MT, NE, NV, NH, NJ, NM, NY, NC, ND, OH, OK, OR, PA, PR, RI, SC, SD, TN, TX, UT, VT, VA, WA, WV, WI, WY

140 Species Affected:

Inverse Name:	Taxa:	Co. occurrence:
Status:		
Cavefish, Alabama	Fish	11
Endangered		
Chub, Bonytail	Fish	148
Endangered		
Chub, Gila	Fish	105
Endangered		
Chub, Humpback	Fish	78
Endangered		
Chub, Mohave Tui	Fish	88
Endangered		
Chub, Oregon	Fish	104
Endangered		
Chub, Owens Tui	Fish	13
Endangered		
Chub, Pahrnagat Roundtail	Fish	1
Endangered		
Chub, Virgin River	Fish	30
Endangered		
Chub, Yaqui	Fish	16
Endangered		
Cui-ui	Fish	7
Endangered		
Dace, Ash Meadows Speckled	Fish	16
Endangered		
Dace, Clover Valley Speckled	Fish	1
Endangered		
Dace, Independence Valley Speckled	Fish	1
Endangered		
Dace, Kendall Warm Springs	Fish	1
Endangered		
Dace, Moapa	Fish	10
Endangered		
Darter, Amber	Fish	47
Endangered		
Darter, Bluemask (=jewel)	Fish	23
Endangered		
Darter, Boulder	Fish	31
Endangered		
Darter, Duskytail	Fish	29

Endangered		
Darter, Etowah	Fish	28
Endangered		
Darter, Fountain	Fish	71
Endangered		
Darter, Maryland	Fish	18
Endangered		
Darter, Okaloosa	Fish	23
Endangered		
Darter, Relict	Fish	10
Endangered		
Darter, Vermilion	Fish	8
Endangered		
Darter, Watercress	Fish	8
Endangered		
Gambusia, Big Bend	Fish	12
Endangered		
Gambusia, Clear Creek	Fish	2
Endangered		
Gambusia, Pecos	Fish	25
Endangered		
Gambusia, San Marcos	Fish	62
Endangered		
Goby, Tidewater	Fish	307
Endangered		
Logperch, Conasauga	Fish	24
Endangered		
Logperch, Roanoke	Fish	125
Endangered		
Madtom, Pygmy	Fish	16
Endangered		
Madtom, Scioto	Fish	45
Endangered		
Madtom, Smoky	Fish	15
Endangered		
Minnow, Rio Grande Silvery	Fish	108
Endangered		
Poolfish, Pahrump (= Pahrump Killifish)	Fish	25
Endangered		
Pupfish, Ash Meadows Amargosa	Fish	11
Endangered		
Pupfish, Comanche Springs	Fish	19
Endangered		
Pupfish, Desert	Fish	198
Endangered		
Pupfish, Devils Hole	Fish	21
Endangered		
Pupfish, Leon Springs	Fish	7
Endangered		
Pupfish, Owens	Fish	13
Endangered		
Pupfish, Warm Springs	Fish	11
Endangered		
Salmon, Atlantic	Fish	73
Endangered		
Salmon, Chinook (Sacramento River Winter Run)	Fish	249
Endangered		
Salmon, Chinook (Upper Columbia River Spring)	Fish	209
Endangered		
Salmon, Coho (Central California Coast population)	Fish	109
Endangered		
Salmon, Sockeye (Snake River population)	Fish	187
Endangered		
Sawfish, Smalltooth	Fish	50
Endangered		
Shiner, Cahaba	Fish	49
Endangered		

Shiner, Cape Fear	Fish	65
Endangered		
Shiner, Palezone	Fish	42
Endangered		
Shiner, Topeka	Fish	294
Endangered		
Spinedace, White River	Fish	15
Endangered		
Springfish, Hiko White River	Fish	3
Endangered		
Springfish, White River	Fish	1
Endangered		
Squawfish, Colorado	Fish	185
Endangered		
Steelhead, (Southern California population)	Fish	126
Endangered		
Stickleback, Unarmored Threespine	Fish	88
Endangered		
Sturgeon, Alabama	Fish	33
Endangered		
Sturgeon, Pallid	Fish	915
Endangered		
Sturgeon, Shortnose	Fish	1090
Endangered		
Sturgeon, White	Fish	8
Endangered		
Sucker, June	Fish	13
Endangered		
Sucker, Lost River	Fish	25
Endangered		
Sucker, Modoc	Fish	9
Endangered		
Sucker, Razorback	Fish	282
Endangered		
Sucker, Shortnose	Fish	13
Endangered		
Topminnow, Gila (Yaqui)	Fish	124
Endangered		
Trout, Gila	Fish	49
Endangered		
Woundfin	Fish	30
Endangered		
Catfish, Yaqui	Fish	16
Threatened		
Cavefish, Ozark	Fish	87
Threatened		
Chub, Chihuahua	Fish	12
Threatened		
Chub, Hutton Tui	Fish	8
Threatened		
Chub, Slender	Fish	76
Threatened		
Chub, Sonora	Fish	7
Threatened		
Chub, Spotfin	Fish	200
Threatened		
Dace, Blackside	Fish	81
Threatened		
Dace, Desert	Fish	6
Threatened		
Dace, Foskett Speckled	Fish	1
Threatened		
Darter, Bayou	Fish	18
Threatened		
Darter, Cherokee	Fish	28
Threatened		
Darter, Goldline	Fish	32
Threatened		
Darter, Leopard	Fish	30
Threatened		

Darter, Niangua	Fish	103
Threatened		
Darter, Slackwater	Fish	59
Threatened		
Darter, Snail	Fish	187
Threatened		
Madtom, Neosho	Fish	48
Threatened		
Madtom, Yellowfin	Fish	93
Threatened		
Minnow, Devils River	Fish	3
Threatened		
Minnow, Loach	Fish	123
Threatened		
Salmon, Chinook (California Coastal Run)	Fish	60
Threatened		
Salmon, Chinook (Central Valley Fall Run)	Fish	40
Threatened		
Salmon, Chinook (Central Valley Spring Run)	Fish	319
Threatened		
Salmon, Chinook (Lower Columbia River)	Fish	119
Threatened		
Salmon, Chinook (Puget Sound)	Fish	171
Threatened		
Salmon, Chinook (Snake River Fall Run)	Fish	190
Threatened		
Salmon, Chinook (Snake River spring/summer)	Fish	206
Threatened		
Salmon, Chinook (Upper Willamette River)	Fish	212
Threatened		
Salmon, Chum (Columbia River population)	Fish	90
Threatened		
Salmon, Chum (Hood Canal Summer population)	Fish	52
Threatened		
Salmon, Coho (Southern OR/Northern CA Coast)	Fish	164
Threatened		
Salmon, Sockeye (Ozette Lake population)	Fish	7
Threatened		
Sculpin, Pygmy	Fish	10
Threatened		
Shiner, Arkansas River	Fish	260
Threatened		
Shiner, Beautiful	Fish	42
Threatened		

Shiner, Blue Threatened	Fish	73
Shiner, Pecos Bluntnose Threatened	Fish	18
Silverside, Waccamaw Threatened	Fish	12
Smelt, Delta Threatened	Fish	82
Spikedace Threatened	Fish	123
Spinedace, Big Spring Threatened	Fish	1
Spinedace, Little Colorado Threatened	Fish	21
Springfish, Railroad Valley Threatened	Fish	13
Steelhead, (California Central Valley population) Threatened	Fish	394
Steelhead, (Central California Coast population) Threatened	Fish	151
Steelhead, (Lower Columbia River population) Threatened	Fish	136
Steelhead, (Middle Columbia River population) Threatened	Fish	195
Steelhead, (Northern California population) Threatened	Fish	99
Steelhead, (Snake River Basin population) Threatened	Fish	219
Steelhead, (South-Central California population) Threatened	Fish	102
Steelhead, (Upper Columbia River population) Threatened	Fish	210
Steelhead, (Upper Willamette River population) Threatened	Fish	197
Steelhead, Puget Sound Threatened	Fish	197
Sturgeon, green Threatened	Fish	75
Sturgeon, Gulf Threatened	Fish	587
Sucker, Santa Ana Threatened	Fish	85
Sucker, Warner Threatened	Fish	8
Trout, Apache Threatened	Fish	43
Trout, Bull Threatened	Fish	712
Trout, Bull (Columbia River population) Threatened	Fish	508
Trout, Bull (Klamath River population) Threatened	Fish	505
Trout, Greenback Cutthroat Threatened	Fish	53
Trout, Lahontan Cutthroat Threatened	Fish	129
Trout, Little Kern Golden Threatened	Fish	44
Trout, Paiute Cutthroat Threatened	Fish	50

No species were selected for exclusion.

Species in Counties by State and Taxa

No species were excluded

Minimum of 1 Acre

All Medium Types Reported

Mammal, Marine mml, Bird, Amphibian, Reptile, Crustacean, Bivalve, Gastropod, Arachnid, Insect, Dicot,
Monocot, Ferns, Conf/cycds, Coral, Lichen
hops, sugarbeets for sugar (irrigated)

40 Species Affected:

Inverte Name:	Taxa:	Co. occurrence:
Status:		
Bat, Indiana	Mammal	11
Endangered		
Butterfly, Fender's Blue	Insect	1
Endangered		
Butterfly, Karner Blue	Insect	2
Endangered		
Butterfly, Mitchell's Satyr	Insect	1
Endangered		
Cactus, Wright Fishhook	Dicot	1
Endangered		
Caribou, Woodland	Mammal	1
Endangered		
Crane, Whooping	Bird	12
Endangered		
Daisy, Willamette	Dicot	2
Endangered		
Ferret, Black-footed	Mammal	16
Endangered		
Limpet, Banbury Springs	Gastropod	2
Endangered		
Lomatium, Bradshaw's	Dicot	2
Endangered		
Penstemon, Blowout	Dicot	4
Endangered		
Plover, Piping	Bird	10
Endangered		
Rabbit, Pygmy	Mammal	2
Endangered		
Riffleshell, Northern	Bivalve	1
Endangered		
Snail, Snake River Physa	Gastropod	4
Endangered		
Snail, Utah Valvata	Gastropod	3
Endangered		
Springsnail, Bruneau Hot	Gastropod	1
Endangered		
Tern, Interior (population) Least	Bird	7
Endangered		
Bear, Grizzly	Mammal	7
Threatened		
Butterfly Plant, Colorado	Dicot	3
Threatened		
Checker-mallow, Nelson's	Dicot	3
Threatened		
Clover, Prairie Bush	Dicot	1
Threatened		
Daisy, Lakeside	Dicot	1
Threatened		
Ladies'-tresses, Ute	Monocot	3

Threatened		
Lupine, Kincaid's	Dicot	1
Threatened		
Milk-vetch, Heliotrope	Dicot	1
Threatened		
Mouse, Preble's Meadow Jumping	Mammal	5
Threatened		
Murrelet, Marbled	Bird	1
Threatened		
Orchid, Eastern Prairie Fringed	Monocot	8
Threatened		
Orchid, Western Prairie Fringed	Monocot	3
Threatened		
Owl, Mexican Spotted	Bird	1
Threatened		
Owl, Northern Spotted	Bird	4
Threatened		
Prairie Dog, Utah	Mammal	1
Threatened		
Snail, Bliss Rapids	Gastropod	3
Threatened		
Snake, Lake Erie Water	Reptile	1
Threatened		
Thelypody, Howell's Spectacular	Dicot	1
Threatened		
Thistle, Pitcher's	Dicot	1
Threatened		
Townsendia, Last Chance	Dicot	1
Threatened		
Yellowhead, Desert	Dicot	1
Threatened		

No species were selected for exclusion.

Dispersed species included in report. 1/28/2010 11:13:28 AM Ver. 2.10.4 Page 4 of 5 1/28/2010 11:13:33 AM Ver. 2.10.4 Page 5 of 5