

014302
014307

DATA EVALUATION RECORD

FENPYROXIMATE
(5% SC FORMULATION)

Study Type: §81-5, Primary Dermal Irritation

Work Assignment No. 2-01-56M (MRID 44781009)

Prepared for

Health Effects Division
Office of Pesticide Programs
U.S. Environmental Protection Agency
1921 Jefferson Davis Highway
Crystal Mall II
Arlington, VA 22202

Prepared by

Pesticide Health Effects Group
Sciences Division
Dynamac Corporation
2275 Research Boulevard
Rockville, MD 20850-3268

Primary Reviewer:
Christie E. Padova, B.S.

Signature: Christie E. Padova
Date: 2-11-00

Project Manager:
Mary L. Menettez, Ph.D.

Signature: Mary L. Menettez *CDP*
Date: 2-11-00

Disclaimer

This Data Evaluation Record may have been altered by the Health Effects Division subsequent to signing by Dynamac Corporation personnel.

1

014302

FENPYROXIMATE (5% SC FORMULATION)

Primary Dermal Irritation Study (§81-5)

EPA Reviewer: Marion Copley
Registration Action Branch 1/HED (7509C)

M. Copley 5/23/00

EPA Work Assignment Manager: Marion Copley
Registration Action Branch 1/HED (7509C)

M. Copley 5/23/00

DATA EVALUATION RECORD

STUDY TYPE: Primary Dermal Irritation - Rabbit
OPPTS Number: 870.2500

OPP Guideline Number: §81-5

DP BARCODE: D259079
P.C. CODE: 129131

SUBMISSION CODE: S564863
TOX. CHEM. NO.: None

TEST MATERIAL (PURITY): NNI-850 Flowable-R (5% fenpyroximate)

SYNONYMS: Fenpyroximate 5% SC Formulation

CITATION: Haynes, G. (1990) Fenpyroximate 5% SC: primary dermal irritation in rabbits. Toxicol Laboratories, Ltd., Herefordshire, England. Laboratory Report No. A/E 24652. May 1990. MRID 44781009. Unpublished.

SPONSOR: Nihon Nohyaku Company, Ltd., 2-5 Nihonbashi, 1-Chome, Chuo-ku, Tokyo 103, Japan.

EXECUTIVE SUMMARY: In a primary dermal irritation study (MRID 44781009), six young adult New Zealand White female rabbits were dermally exposed to 0.5 mL of NNI-850 Flowable-R (5% fenpyroximate) for 4 hours. The test material was applied as received to a single intact 6.25-cm² site/animal. Animals were observed for dermal irritation for up to 7 days following patch removal, and irritation was scored by the Draize scale.

Very slight erythema was observed at 1/6 test sites 1 hour following patch removal. No other dermal irritation was observed during the 7-day study. In this study, NNI-850 Flowable-R (5% fenpyroximate) is not a notable dermal irritant, and is classified as TOXICITY CATEGORY IV for primary dermal irritation.

This study is classified acceptable (§81-5) and satisfies the guideline requirement for a primary dermal irritation study in the rabbit.

COMPLIANCE: Signed and dated GLP, Quality Assurance, and Data Confidentiality statements were provided.

I. MATERIALS AND METHODS**A. MATERIALS:**

1. **Test Material:** NNI-850 Flowable-R
Description: White liquid
Lot/Batch #: 900216
Purity: 5% Fenpyroximate
CAS #: Not provided
2. **Vehicle and/or positive control:** None employed
3. **Test animals:** Species: Albino rabbit
Strain: New Zealand White
Age: Approximately 10-12 weeks
Weight: Approximately 2 kg (not further specified)
Source: Ranch Rabbits, Crawley Down, Sussex, England
Acclimation period: ≥5 Days
Diet: SQC Standard Rabbit Pellets (Special Diets Services, Witham, Essex, England), ad libitum
Water: Tap water, ad libitum
Housing: Individually in grid-bottomed metal cages
Environmental conditions:
Temperature: 17-21 °C
Humidity: 45-65%
Air changes: Not specified
Photoperiod: 12-hour light/dark cycle

B. STUDY DESIGN and METHODS:

1. **In-life dates:** April 19-26, 1990
2. **Animal assignment and treatment:** Fur from the dorsal area of six young adult New Zealand White female rabbits was clipped 1 day prior to dermal administration with 0.5 mL of NNI-850 Flowable-R (5% fenpyroximate). The test material was applied as received to a single intact site per animal using a 6.25-cm² piece of surgical lint. The patches were held in place by wrapping the torso of each animal with Elastoplast elastic adhesive bandage. The coverings were removed 4 hours following application and residual test material was removed using warm water. Dermal irritation was observed at 1, 24, 48, and 72 hours and 7 days following patch removal. Erythema and edema were scored separately using the Draize scale.

II. RESULTS AND DISCUSSION:

- A. Clinical observations: Very slight erythema (score of 1) was observed at 1/6 test sites 1 hour following patch removal. No other dermal irritation was observed during the 7-day study. In this study, NNI-850 Flowable-R (5% fenpyroximate) is not a notable dermal irritant.
- B. Deficiencies: There were no deficiencies that affected the validity of the study results.