

5% CHLORDANE

5% CHLORDANE

The 1 pound package may be used as a duster for small garden or home use. Shake can to fluff powder. Open one or more holes and press sides of container with fingers, using rapid motion. Cover all parts of the plant with a thin uniform film of dust, top and undersides of leaves.

DIRECTIONS

Apply thoroughly when damage or insects are noticed and repeat treatments within 1 year should not be necessary.

TURF AND LAWN INSECTS — To control white grub: Apply to the soil's surface at the rate of 1 lb. to 436 sq. ft.; 2½ to 5 lbs. per 1,000 sq. ft. 20' x 50'; 100 to 200 pounds per acre. Wash the insecticide into the soil to a depth of 4 to 6 inches with ample quantities of water. Apply in early spring or late fall.

TO CONTROL JAPANESE BEETLE LARVAE — Use same as for white grubs, apply in early spring or late fall.

TO CONTROL EARWIGS — Apply to lawn or areas infested at the rate of ¼ pound per 1,000 sq. ft.; 30 pounds per acre. Make application in the evening during fair weather when rain is not expected. Avoid wetting down the treated area for at least two nights.

TO CONTROL CHINCH BUGS — Apply to lawn or turf at the rate of 1 lb. per 1,000 sq. ft.; 4 lbs. to 4,000 sq. ft.; 40 lbs. per acre.

TO CONTROL SOD WEBWORMS — Apply to lawn or turf as soon as worms appear at the rate of 1 lb. to 400 sq. ft.; 2½ lbs. per 1,000 sq. ft.; 40 to 100 lbs. per acre. Water the lawn thoroughly before treatment. Do not water again for several days.

TO CONTROL MOLE CRICKETS — Apply to lawn or turf at the rate of 1 lb. per 1,000 sq. ft.; 4 lbs. per 4,000 sq. ft.; 40 lbs. per acre. Apply generous quantities of water prior to the application, then follow with additional water to carry the insecticide into the soil.

TO CONTROL CHIGGERS — On lawn, ground, litter and vegetation at the rate of 1 lb. per 1,000 sq. ft.; 4 pounds per 4,000 sq. ft.; 40 lbs. per acre.

TO CONTROL WIREWORMS — Apply to lawn or turf at the rate of 1 lb. per 400 sq. ft.; 2½ lbs. per 1,000 sq. ft.; 80 to 100 lbs. per acre. Water the same as for white grubs.

TO CONTROL ANTS — Apply on lawn or turf at the rate of 30 lbs. per acre (¾ lb. per 1,000 sq. ft.). For spot control apply liberally to ant hills and wash the insecticide into the soil.

GARDEN INSECTS

May be used on potatoes, tomatoes and cabbage to control flea beetles, cabbage worms, tomato hornworms, Colorado potato beetles and potato psyllids. Do not apply when plants are in bloom as Chlordane is toxic to beneficial insects as well as pests. Apply uniform thin film of dust as soon as insects appear and repeat treatment as necessary.

Potatoes: Do not exceed 1 pound to 1,500 sq. ft., 4 pounds to 6,000 sq. ft. or 30 pounds per acre.

Tomatoes: Do not exceed 1 pound to 436 sq. ft., 4 pounds to 1,722 sq. ft. or 100 pounds per acre. Do not apply after fruit begins to form.

Cabbage: Do not exceed 1 pound to 1,100 sq. ft.; 4 pounds to 4,350 sq. ft. or 40 lbs. per acre. Do not apply after edible parts start to form.

CAUTION: Avoid inhalation of dust. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Avoid contamination of foodstuff and use on leafy vegetables, such as lettuce. Do not apply to food crops after edible parts begin to form. Remove residue at harvest by washing, wiping or other means. Avoid use on gourds or other varieties of cucurbits, which may be injured by Chlordane.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Buyer's guarantee limited to label claims.

BONIDE CHEMICAL COMPANY, INC., UTICA, NEW YORK 13502