


B O N I D E

CHLORDANE 10% DUST



DIRECTIONS

Apply thoroughly when damage or insects are noticed and make repeat treatments as often as necessary.

TURF AND LAWN INSECTS — To control white grubs: Apply to the soil's surface at the rate of 50 to 100 pounds per acre (1½ to 2½ lbs. per 1,000 sq. ft. 20' x 50'). Wash the insecticide into the soil to a depth of 4 to 6 inches with ample quantities of water. Apply in early spring or late fall. (Keep children and pets off treated areas until thoroughly dried.)

TO CONTROL JAPANESE BEETLE LARVAE — Use same as for white grubs, apply in early spring or late fall.

TO CONTROL EARWIGS — Apply to lawn or areas infested at the rate of 15 pounds per acre (½ pound per 1,000 sq. ft.). Make application in the evening during fair weather when rain is not expected. Avoid wetting down the treated area for at least two nights.

TO CONTROL CHINCH BUGS — Apply to lawn or turf at the rate of 20 lbs. per acre (½ lb. per 1,000 sq. ft.).

TO CONTROL SOD WEBWORMS — Apply to lawn or turf as soon as worms appear at the rate of 20 to 50 lbs. per acre (1¼ lbs. per 1,000 sq. ft.). Water the lawn thoroughly before treatment. Do not water again for several days.

TO CONTROL MOLE CRICKETS — Apply to lawn or turf at the rate of 20 lbs. per acre (½ lb. per 1,000 sq. ft.). Apply generous quantities of water prior to the application, then follow with additional water to carry the insecticide into the soil.

TO CONTROL CHIGGERS — Dust on lawn, ground, litter and vegetation at the rate of 20 lbs. per acre (½ lb. per 1,000 sq. ft.).

TO CONTROL WIREWORMS — Apply to lawn or turf at the rate of 40 to 50 lbs. per acre (1¼ lbs. per 1,000 sq. ft.). Water the same as for white grubs.

TO CONTROL ANTS — Dust on lawn or turf at the rate of 15 lbs. per acre (¾ lb. per 1,000 sq. ft.). For spot control apply liberally to ant hills and wash the insecticide into the soil.

TO CONTROL SURFACE CUTWORMS — Apply insecticide to soil surface around base of plants at time of planting or transplanting. Cover the soil with a visible layer of dust, to protect seedlings and small transplants such as tomato, pepper, cabbage. Make application late in the day.

GARDEN INSECTS

This dust may be used on potatoes to control Colorado Potato Beetle Larvae, repel Potato Flea Beetle at the rate of 15 lbs. per acre. Tomatoes — to repel Potato Flea Beetle and control Colorado Potato Beetle Larvae, use at the rate of 50 lbs. per acre. (Do not apply after fruit begins to form). Cabbage, Broccoli and Brussels Sprouts to control Imported Cabbage Worms, Diamond Back Moths use at the rate of 20 lbs. per acre. (Do not apply after edible parts start to form). Do not apply when plants are in bloom, since Chlordane is toxic to beneficial insects as well as pests. Apply dust as soon as insects appear and repeat treatment as necessary.


CAUTION: Avoid inhalation of dust. Avoid contamination of feed and foodstuff. Avoid use on Lettuce and Endive. Do not apply to food crops after edible parts begin to form. Remove residue at harvest by washing, wiping or other means. Avoid use on gourds or other varieties of cucurbits, which may be injured by chlordane. Do not contaminate any body of water nor apply to any area not specified on the label.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed.

USDA Reg. No. 4-67

Mfg. by Bonide Chemical Co., Inc., Utica, N.Y.

Buyer's Guarantee Limited to Label Claims



4 67