

Reg # 4-81
PM 17
P/11

BONIDE®

ACCEPTED

AUG 18 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under 4-81 EPA Reg. No.

BLUE DEATH .75% ROTENONE GARDEN DUST

CONTROLS
MEXICAN BEAN BEETLES, IMPORTED CABBAGE WORMS,
DIAMOND BACK MOTH, AND CERTAIN OTHER INSECTS
ON VEGETABLES, FLOWERS AND BUSH FRUITS.

ACTIVE INGREDIENTS:
Rotenone .75%
Other cube salts 1.50%
INERT INGREDIENTS 97.75%
EPA Est. 4-NV-1 EPA Reg. No. 4-81

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions)

1 LB. NET 4 LBS. NET

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION
May be harmful if swallowed. Avoid breathing dust or spray. Avoid contact with eyes, skin, clothing. Wash thoroughly after use. In case of contact rinse with plenty of water.
ENVIRONMENTAL HAZARDS
This product is toxic to aquatic organisms. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply while bees are actively visiting the area. Additional protective information may be obtained from your Cooperative Extension Service.

DIRECTIONS FOR USE
It is violation of Federal law to use this product in a manner inconsistent with its labeling.
Storage: Store in a cool dry place which is inaccessible to children and pets.
Container Disposal: Do not reuse empty container. Wrap container and put in trash collection.

Blue Death .75 Rotenone Dust is a high quality insecticide containing special micronized cube powder and fibrous talc, attritionized for better dispersion of toxicant and super adhesive qualities.
Recommended for the control of certain insects on beans, such as Mexican bean beetles, Cautiflower: flea beetles, cabbage worms. Cucumbers, Melons: flea beetles, striped cucumber beetles. Tomatoes, Eggplant's: flea beetles, horn worm, stink bugs. Roses, Asters, Dahlias, Gladiolus: flea beetles, leafroller, Japanese beetle, leaf chafer. Bush fruits - currants, gooseberries: flea beetles, imported currant worms, climbing cutworms. Do not apply to food crops within one day of harvest.

DIRECTIONS
FOR DUSTING:
Use a good duster and apply a uniform film of dust to all parts of the plant, upper and under surfaces of the leaves. Begin when insects first appear and repeat treatment weekly or as often as necessary.
FOR SPRAYING:
Mix at the rate of 1 pound to 5 gallons. Apply so as to thoroughly wet the plant, both upper and under surfaces of the leaves. Begin application when insects first appear and repeat treatment weekly or as often as necessary.

USE BONIDE VEGETABLE-FLORAL DUST OR SPRAY INSECTICIDE-FUNGICIDE
FOR CERTAIN INSECTS AND DISEASES

Buyer's guarantee limited to label claims.