


4-107

01/30/2004

1/4

MASTER LABEL



GARDEN DUST
for
Vegetables – Flowers

[Additional Brand Names: Fruit Tree Spray E.F.; Rose & Flower Dust E.F.]

Optional Marketing Statements:

CONTROLS Aphids, Certain Beetles, Spider Mites, Roller Worms – Anthracnose, Leaf Spot, Powdery Mildew, Early and Late Blights.

Use as a Dust or Spray

ACTIVE INGREDIENTS:

Pyrethrins	0.03%
Rotenone	0.50%
Other Cube Resins	0.75%
Sulfur (4-5 microns).....	25.00%
Copper (basic copper sulfate)*	5.00%
OTHER INGREDIENTS:	68.72%
TOTAL:	100.00%

*Metallic copper equivalent 2.7%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back/side/other panel(s) for first aid and additional precautionary statements.

NET WEIGHT _____ OZ. (_____ g)

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.	
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If swallowed	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin, inhaled or swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust (vapor or spray mist). Remove contaminated clothing and wash before reuse. Do not reenter treated areas until sprays have dried or dusts have settled.

Environmental Hazards

This pesticide is extremely toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Storage: Store and transport in an upright position. Store in a cool, dry area inaccessible to children or pets.

Disposal: *If empty* -- Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* -- Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GARDEN DUST is a specially formulated insecticide and fungicide designed for convenient use as a dust or spray, colored green to blend with foliage, and packaged in a squeeze duster can.

DIRECTIONS FOR HOME GARDEN USE:

AS A DUST: Shake can to fluff powder. Press sides of can with fingers rapidly for dust effect. Cover both tops and undersides of leaves with thin uniform film of dust or spray.

AS A SPRAY: Mix 2 Tbs. per gallon of water. Make a thin paste, strain, add rest of water, and agitate to prevent nozzle plugging.

Repeat treatment as required, usually every 7 to 10 days. **NOTE** -- Stop application to food crops one day before harvest. Start applying when plants leaf out in the spring before insects and diseases appear, as it is easier to prevent than cure certain diseases. Try to hit as many insects as possible. Best results are obtained in early morning or evening when wind is at ebb. Avoid use during high temperatures on plants susceptible to sulfur injury, such as melons, squash, cantaloupes and related plants. Depending on foliage density, one pound should cover approximately 1,000 sq. ft.

FRUITS -- Peach¹, Sour Cherry², Grape³, Apple⁴, Pear⁵, Nectarine⁶.

Insects -- Aphids, Leafhoppers, Exposed Thrips, Cherry Fruit Flies, Red-necked Borers, Japanese Beetles, Mites and Rose Chafers.

Diseases (*superscript following disease corresponds to above crop listing*) -- Scab⁽⁴⁻⁵⁾, Black Rot⁽³⁻⁴⁾, Powdery Mildew⁽³⁻⁴⁾, Fire Blight⁽⁴⁻⁵⁾, Anthracnose⁽³⁻⁵⁾, Brown Rot Blossom Blight^(1-2,6), Sooty Blotch⁽⁴⁾, Leaf Spot^(2,4), Frog Eye⁽⁴⁾, Stem Blight^(1-2,6), Shothole^(1-2,6), Peach Canker^(1,6), Apple Cedar Rust⁽⁴⁾, Peach Leaf Curl^(1,6).

Comments: Certain Vinifera and French Hybro varieties of grapes may be slightly sensitive to copper sprays resulting in marginal leaf burn; test sprays are recommended. Apply to cherries only at popcorn and petal fall stages. Apply to peaches only at bud swell and pink bud stages.

VEGETABLES -- Bean¹, Beet², Cucumber³, Squash⁴, Muskmelon⁵, Tomato⁶, Celery⁷, Pepper⁸, Onion⁹, Carrot¹⁰, Potato¹¹, Sweet Peas¹²

Insects -- Aphids, Leafhoppers, Exposed Thrips, Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Strick Bugs, Celery Leaf-tiers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites, Colorado Potato Beetle, Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers

Diseases (*superscript following disease corresponds to above crop listing*) –
 Bacterial Blights^(1-2,7), Early Blights^(1-2,6-8,10-11), Powdery Mildew^(1,3-5,12), Downy
 Mildew^(1-5,8-10,12), Alternaria Blight^(3-5,6,8,10), Angular Leaf Spot^(1-5,7,9,12), Bacterial
 Spot^(6,8), Septoria Leaf Spot^(2-7,9,12), Cercospora Leaf Spot^(2-3,7-9,12).

Comments: For European Corn Borers, make 3-4 applications or more as eggs start to hatch and repeat at 5-7 day intervals.

ORNAMENTALS – Pine¹ (Austrian, Ponderosa), Rose².

Insects – Aphids, Leafhoppers, Exposed Thrips, Oblique-banded Leaf Rollers, Sawfly Larvae, Boxelder Bugs, Japanese Beetles, Elm Leaf Beetle Larvae, Mites, Lygus Bugs, Lace Bugs, Greenhouse Whiteflies, Stink Bugs, Garden Fleahoppers, Greenhouse Leafhoppers, Cucumber Beetle, Rose Chafers, Ants, Aster Beetles, Flea Beetles.

Diseases (*superscript following disease corresponds to above crop listing*) –
 Pine Needle Blight¹, Black Spot², Powdery Mildew², Rust².

Comments: For disease control, apply to point of runoff when new growth is emerging.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Bonide Products, Inc.
 6301 Sutliff Road
 Oriskany, NY 13424