

BONIDE

[4-110] (of 2)

25% MALATHION

WETTABLE SPRAY POWDER

ACCEPTED
JUN 13 1975

ACTIVE INGREDIENT:

Malathion

INERT INGREDIENTS:

CAUTION: Keep out of reach of children.

Harmful if swallowed or inhaled. Avoid breathing spray mist or contact with skin, eyes or clothing. May be absorbed through skin. Do not use on household pets or human. Wash thoroughly with soap and water before reuse.

This product is toxic to fish, wildlife, and other aquatic organisms. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where run off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

foodstuff.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

BONIDE 25% MALATHION WETTABLE SPRAY POWDER is a registered pesticide of the United States Department of Agriculture. It is a restricted use pesticide and its use is limited to those persons who are registered with the United States Department of Agriculture. For more information, contact your local Agricultural Extension Service.

NET WEIGHT LBS.

FIRST PROOF

Page (2 of 2)

WATER-SOLUBLE POWDER

INSTRUCTIONS

Mix the proper amount with water in a clean container. The mixture should be applied to the spray tank 1/4 full of water and add mixture to tank. The mixture should be applied to the plants in the morning or evening. Do not apply if rain is expected for the next 24 hours.

Four (4) applications are recommended. Applications call for use of 1/2 lb. of powder per acre. For example, 1/2 lb. of powder per acre requires 4 lbs. of powder per acre.

CROP	INSECT	WHEN TO APPLY	AMOUNT
CUCUMBER, SQUASH	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	4 lbs. gal. water, or 6 lbs. 100 gals. water per acre + 5 lbs. gal. water, or 7 lbs. 100 gals. water per acre
Broccoli	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	4 lbs. gal. water, or 2 lbs. 100 gals. water per acre
Kale	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	2 1/2 - 3 lbs. gal. water, or 4-5 lbs. 100 gals. water per acre
Mustard Green	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 lbs. gal. water, or 5 lbs. 100 gals. water per acre
Turnips	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 lbs. gal. water, or 5 lbs. 100 gals. water per acre
ENDIVE	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	1 1/2 lbs. gal. water, or 2 lbs. 100 gals. water per acre
BEETS	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 1/2 lbs. gal. water, or 5 lbs. 100 gals. water per acre 5 lbs. gal. water, or 8 lbs. 100 gals. water per acre
CARROTS	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	1 1/2 lbs. gal. water, or 2 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
BEAN	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 1/2 lbs. gal. water, or 5 lbs. 100 gals. water per acre
TOMATOES	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	2 1/2 - 3 lbs. gal. water, or 4-5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
EGGPLANT	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	1 1/2 lbs. gal. water, or 2 lbs. 100 gals. water per acre 2 1/2 lbs. gal. water, or 4 lbs. 100 gals. water per acre
ONIONS	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	4 lbs. gal. water, or 6 lbs. 100 gals. water per acre 5 lbs. gal. water, or 8 lbs. 100 gals. water per acre
PEPPERS	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	4 lbs. gal. water, or 6 lbs. 100 gals. water per acre 5 lbs. gal. water, or 8 lbs. 100 gals. water per acre
LETTUCES, WEEB, CHLITONS	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 lbs. gal. water, or 5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
ARTICHOKES	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	3 lbs. gal. water, or 5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	2 1/2 - 3 lbs. gal. water, or 4-5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
BLUEBERRIES	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	2 1/2 - 3 lbs. gal. water, or 4-5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre
STRAWBERRIES	Colorado potato beetle	Apply 1st application at first sign of infestation. Repeat every 7-10 days.	2 1/2 - 3 lbs. gal. water, or 4-5 lbs. 100 gals. water per acre 4 lbs. gal. water, or 6 lbs. 100 gals. water per acre

