

BONIDE

FLOWER-VEGETABLE 4-IN-1 SPRAY

APHIDICIDE - MITICIDE
INSECTICIDE - FUNGICIDE

EPA REG. NO. 4127

ACTIVE INGREDIENTS

Malathion: 0,0-diethyl phosphorodithioate of diethyl mercaptosuccinate... 9.4%

Methoxychlor: 1,1,1-trichloro-2, 2-bis(p-methoxyphenyl) ethane..... 9.5%
(equivalent to 8.36% 2,2-bis(p-methoxyphenyl) 1,1,1-trichloroethane and 1.14% related compounds)

2,4-Dinitro 6-octyl*phenyl crotonate.... 1.4%

2,6 Dinitro-4-octyl*phenyl crotonate

Nitrooctyl*phenols (principally o-nitro) .1%

(*A mixture of 1-methylheptyl, 1-ethylhexyl and 1-propylpentyl isomers)

~~Aromatic Petroleum Solvents 13.5%~~

INERT INGREDIENTS 66.1% 79.6%

Contains Aromatic Petroleum Solvents


ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, recommended for the pesticide registered under EPA Reg. No. 4-127

KEEP OUT OF REACH OF CHILDREN
CAUTION

See label for use directions and cautions

NET CONTENTS 1 Quart



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION

Harmful if swallowed or inhaled. Avoid contact with skin and eyes. May be absorbed through skin. In case of contact wash immediately with soap and water. Avoid breathing spray mist. May be skin irritant and/or sensitizer. Wash hands and contaminated clothing after use. Do not take or administer internally. Avoid contamination of foodstuffs, feed, feeding troughs, watering receptacles, and water supplies. Wear rubber gloves when using indoors. When using outdoors wear long pants, long sleeve shirt, and rubber gloves. Outdoors, spray with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Use of this product may be hazardous to your health. This product has been determined to cause birth defects in laboratory animals.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL & CHEMICAL HAZARDS

DANGER- EXTREMELY FLAMMABLE. KEEP AWAY FROM FIRE, SPARKS AND HEATED SURFACES. ALWAYS KEEP CONTAINER TIGHTLY CLOSED (PREVENTS EVAPORATION).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner consistent with its labeling.

10 Point Type

STORAGE AND DISPOSAL

Storage: Keep container tightly closed to prevent evaporation. Do not store near or contaminate feed or foodstuffs. Do not use, pour, spill or store near heat or open flame. Store in a cool, dry area inaccessible to children and pets.
Disposal: Do not reuse empty container. Wrap container and put in trash collection.

BONIDE FLOWER - VEGETABLE 4-IN-1 SPRAY, EPA Reg. No. 4-126

FOR HOME GARDEN USE.

FLOWER-VEGETABLE 4-in-1 SPRAY is a concentrated combination of aphicide, miticide, insecticide and fungicide. It easily and instantly mixes with water for spraying without laborious pre-mixing or straining. Use a good hand or pressure sprayer and apply thoroughly. After you have applied spray you will notice there is no unsightly residue and the leaves actually look clean and shiny.

FLOWER-VEGETABLE 4-in-1 SPRAY is designed for use on flowers, ornamentals (shrubs), vegetables which are listed below.

FLOWERS, ORNAMENTALS: Aster, Chrysanthemum, Dahlia, Delphinium, Evonymus, Geranium, Gladiolus, Hydrangea, Nasturtium and Rose.

To Control: Aphids (black, pink, white, green), Spider Mites (red, clover, two spotted), Four Lined Lace Bug, Lace Bug, Leaf Hoppers, Mealy Bug, Japanese Beetle (adult), Scale and Powdery Mildew.

VEGETABLES: Cucumber, Cantaloupe, Squash and Melon.

To Control: Cucumber Beetle, Squash Vine Borer, Melon Worm, Aphid, Spider Mite, Leaf Hopper and Powdery Mildew.

DIRECTIONS FOR APPLICATION

Use at the rate of 10 teaspoons or 3 1/3 tablespoons or 1 2/3 oz. per gallon for insects and disease. Frequent and thorough application, particularly undersides of leaves and stems is the key to successful mildew and insect control. Start applications at first sign of insect or mildew activity and repeat applications every 7-10 days. Do not apply during high temperatures (90° F.)

IMPORTANT: Do not apply to cantaloupes, cucumbers, melons or squash within 7 days of harvest. Vegetables should be washed, blanched, wiped or otherwise cleaned to remove residues prior to cooking and eating.

Buyer's guarantee limited to label claims

~~© 1973~~ Mfg. by BONIDE CHEMICAL CO., INC., Yorkville, N.Y. 13495

