

ACCEPTED

APR 4, 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 4-133
SUBJECT
TO ATTACHED COMMENT

U S D E P

SEVIN[®] 50W

Insecticide

A WETTABLE POWDER FOR CONTROL OF INSECT PESTS

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate)

INERT INGREDIENTS:
Union Carbide Corp. Reg. T.M.

EPA Reg. No. 4-133
50% by WT.
50% by WT.

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions) **8 OZ. NET**

NEW INSECTICIDE "Sevin" 50W

Controls the major pests of vegetables. Can be used close to harvest to prevent damage by late broods of insects. Also effective in controlling Japanese Beetle and certain other insects on roses, flowers, shrubs, ornamentals.

DIRECTIONS FOR VEGETABLE INSECT CONTROL

SMALL SCALE OR HOME GARDEN USE: Use 2 level tablespoonfuls per gallon of water. Apply with good hand or pressure sprayer using sufficient spray to obtain full coverage. Keep spray mixture well agitated. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary.

CROP	INSECTS CONTROLLED	Time Between Last Application & Harvest
BEANS	Mexican Bean Beetle, Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles, Japanese Beetles.	Day of Harvest Tolerance: 10 PPM
CUCUMBERS SUMMER SQUASH	Pickle and Melonworms, Cucumber Beetles, Squash Bugs, Flea Beetles.	Day of Harvest Tolerance: 10 PPM
SWEET CORN	Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetles. For larvae in whorl (Budworm and Foliage Feeders) apply to entire plant. Repeat as necessary.	Day of Harvest
Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. For insects attacking silks and ears, apply at 2 to 3 day intervals, as infestation warrants, starting when first silks appear, continuing till silks begin to dry. 3 or more applications may be necessary. Timing and good coverage are absolutely essential for effective control.		
Apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge.		
TOMATOES EGGPLANT PEPPERS	Tomato Fruitworm, Fall Armyworm, Flea Beetles, Leaf Hoppers, Tomato Hornworms, European Corn Borer, Stink Bugs	Day of Harvest Tolerance: 10 PPM
CABBAGE BROCCOLI CAULIFLOWER	Flea Beetles, Harlequin Bug, Armyworm, Cabbage Caterpillars, Corn Earworm. (For small Loopers after heads form use SEVIN every 5 to 7 days.)	3 Days Before Harvest Tolerance: 10 PPM
ROSES, FLOWERS, SHRUBS, ORNAMENTALS: Japanese Beetles, Bagworm, Birch Leaf Miners, Elm Leaf Beetles. Apply spray as insects appear and repeat treatments as needed. (Do not apply on Boston Ivy.) Use 2 level tablespoonfuls per gal. water.		
TURF, LAWN	Chinch Bugs, Fall Armyworm, Leafhoppers, Sod Webworms or Lawn Moths, Ants, Earwigs, Fleas, Millipedes, Mosquitoes. Use 1 lb. in 75 gals. water per 5000 sq. ft. of lawn (50' x 100'). 2 tablespoonfuls to 3 gals. water per 100 sq. ft. (10' x 10').	No Time Limit
Before application, mow the lawn close and remove clippings. Water lawn and then apply the product at the recommended dosage.		

CAUTION: Keep out of reach of children. Harmful if inhaled or swallowed. Do not breathe dust. Avoid breathing of spray. Do not take internally. Avoid unnecessary skin contact. Wash thoroughly after using.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Avoid storage near feed and food products. Avoid contamination of food.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Buyer's guarantee limited to label claims

Manufactured by
BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502