



ACCEPTED
 MAR 6 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 4134 SUBJECT
 TO ATTACHED COMMENTS.

BONIDE

* **SEVIN**

2.50%

Insect Dust

FOR

**VEGETABLES • FLOWERS
 ORNAMENTALS • LAWN INSECTS**

ACTIVE INGREDIENT:
 Carbaryl-1-Naphthyl, N-Methylcarbamate . . . 2.50%
INERT INGREDIENTS 97.50%
 100.0%

Made by

BONIDE CHEMICAL CO., INC.
 UTICA, NEW YORK

Reg. T.M. U.

CAUTION: "KEEP OUT OF REACH OF CHILDREN"
(Note Side or Back Panels, Other Cautions)

NEW INSECTICIDE "Sevin"

Controls the major pests of vegetables with a minimum of applications and reduces the hazards to persons applying it and to field workers and pickers. Can be used close to harvest to prevent damage by late broods of insects. — — — — — Also effective in controlling Japanese Beetle and certain other insects on roses, flowers, shrubs, ornamentals. — — — — — Chinch Bugs, Sod Webworms in lawns.

DIRECTIONS FOR USE

SMALL SCALE OR HOME GARDEN USE Apply with a good hand or power duster evenly on all parts of the plant with a uniform film of dust, especially the underside of leaves. Start applications as soon as insects appear and repeat at 7 to 10 day intervals. For large scale use follow chart below.

CROP	INSECTS CONTROLLED	Dust	RECOMMENDED Dose Per Acre	TIME BETWEEN LAST APPLICATION & HARVEST
① BEANS	MEXICAN BEAN BEETLE	2.5%	20 lbs. 1 lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORM, FLEA BEETLES, AND JAPANESE BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
	CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR	2.5%	40 to 60 lbs. 1 to 1.5 lbs. active SEVIN	
	LIMA BEAN POD BORER, LYGUS AND STINK BUGS	2.5%	80 lbs. 2 lbs. active SEVIN	
② CUCUMBERS SUMMER SQUASH	PICKLE AND MELONWORMS	2.5%	20 lbs. 1 lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	CUCUMBER BEETLES, SQUASH BUGS AND FLEA BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
③ SWEET CORN	CORN EARWORM, EUROPEAN CORN BORER, FALL ARMYWORM, SAP BEETLES, AND JAPANESE BEETLES	2.5%	40 to 80 lbs. 1 to 2 lbs. active SEVIN	Day of harvest for fresh market ears 7 days before harvest or forage and husks fed to live stock
④ TOMATOES EGGPLANT PEPPERS	TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES AND LEAFHOPPERS	2.5%	40 lbs. 1 lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	TOMATO HORNWORMS, EUROPEAN CORN BORER AND STINK BUGS	2.5%	40 to 80 lbs. 1 to 2 lbs. active SEVIN	
⑤ POTATOES	COLORADO POTATO BEETLE	2.5%	20 lbs. 1 lb. active SEVIN	Day of harvest
	LEAFHOPPERS AND FLEA BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
	LYGUS BUGS	2.5%	80 lbs. 2 lbs. active SEVIN	

Apply when insects or their damage appear. Repeat at 7 to 10 day intervals as necessary. Do not feed treated bean vines to dairy or meat animals.

For larvae or whole bodyworm damage and foliage feeding apply to entire plant. Repeat as necessary. For insects attacking stalks and ears apply at 2 to 4 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation.

ROSES, FLOWERS, SHRUBS, ORNAMENTALS: Japanese Beetles and certain other insects — Apply a uniform film of dust to all parts of the plant as soon as insects appear and repeat treatments as needed.

LAWN AND TURF (established): Chinch Bugs, Fall Armyworm, Leafhoppers, Sod Webworms — Apply at the rate of 1 to 2 lbs. per acre (2 lbs. per 500 sq. ft. of lawn 20 x 25 sq. ft.) Apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

CAUTION! HARMFUL IF SWALLOWED OR INHALED.

Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed or foodstuffs. Store out of reach of children.

Guaranteed as label implied.

4 lbs. Net

Form 561

U.S.D.A. REG. NO. 4-134

ACCEPTED

NEW INSECTICIDE "Sevin"

Controls the major pests of vegetables, with a minimum of applications and reduces the hazards to persons applying it and to field workers and pickers. Can be used close to harvest to prevent damage by late broods of insects. — — — — — Also effective in controlling Japanese Beetle and certain other insects on roses, flowers, shrubs, ornamentals . . . Chinch Bugs, Sod Webworms in lawns.

DIRECTIONS FOR USE

SMALL SCALE OR HOME GARDEN USE: Apply with a good hand or power duster, cover all parts of the plant with a uniform film of dust, especially the underside of leaves. Start applications as soon as insects appear and repeat at 7 to 10-day intervals. For large scale use follow chart below:

CROP	INSECTS CONTROLLED	Dust	RECOMMENDED Dose Per Acre	TIME BETWEEN LAST APPLICATION & HARVEST
① BEANS	MEXICAN BEAN BEETLE	2.5%	20 lbs. ½ lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	LEAFHOPPERS, BEAN LEAF BEETLE, CUCUMBER BEETLES, WESTERN BEAN CUTWORM, FLEA BEETLES AND JAPANESE BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
	CORN EARWORM, ARMYWORMS, VELVET BEAN CATERPILLAR	2.5%	40 to 60 lbs. 1 to 1½ lbs. active SEVIN	
	LIMA BEAN POD BORER, LYGUS AND STINK BUGS	2.5%	80 lbs. 2 lbs. active SEVIN	
② CUCUMBERS SUMMER SQUASH	PICKLE AND MELONWORMS	2.5%	20 lbs. ½ lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	CUCUMBER BEETLES, SQUASH BUGS AND FLEA BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
③ SWEET CORN	CORN EARWORM, EUROPEAN CORN BORER, FALL ARMYWORM, SAP BEETLES, AND JAPANESE BEETLES	2.5%	40 to 80 lbs. 1 to 2 lbs. active SEVIN	Day of harvest for fresh market ears, 7 days before harvest on forage and husks fed to livestock
④ TOMATOES EGGPLANT PEPPERS	TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES AND LEAFHOPPERS	2.5%	40 lbs. 1 lb. active SEVIN	Day of harvest Tolerance: 10 PPM
	TOMATO HORNWORMS, EUROPEAN CORN BORER AND STINK BUGS	2.5%	40 to 80 lbs. 1 to 2 lbs. active SEVIN	
⑤ POTATOES	COLORADO POTATO BEETLE	2.5%	20 lbs. ½ lb. active SEVIN	Day of harvest
	LEAFHOPPERS AND FLEA BEETLES	2.5%	40 lbs. 1 lb. active SEVIN	
	LYGUS BUGS	2.5%	80 lbs. 2 lbs. active SEVIN	

① ② ③ ④ Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary. (1) Do not feed treated bean vines to dairy or meat animals.

⑤ For larvae in whorl (Budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation.

ROSES, FLOWERS, SHRUBS, ORNAMENTALS: Japanese Beetles and certain other insects — Apply a uniform film of dust to all parts of the plant as soon as insects appear and repeat treatments as needed.

LAWN AND TURF (established): Chinch Bugs, Fall Armyworm, Leafhoppers, Sod Webworms — Apply at the rate of 17½ lbs. per acre (2 lbs. per 500 sq. ft. of lawn 20 x 25 sq. ft.) Apply after rain or watering and do not water for at least 2 days after application. Repeat 2 to 3 weeks later if necessary.

CAUTION! HARMFUL IF SWALLOWED OR INHALED.

Do not take internally. Avoid breathing dust. Avoid prolonged or repeated