

B O N I D E

P A R I S G R E E N P E L L E T E
M O S Q U I T O L A R V A C I D E

FOR GOVERNMENTAL APPLICATION BY PUBLIC HEALTH DEPTS.

ACTIVE INGREDIENTS:

Copper Acetoarsenite* (from Paris Green.....	8.2%
Petroleum Oil.....	36.0%
INERT INGREDIENTS (Carrier #3 Vermiculite).....	55.8%
	<u>100.0%</u>

U.S.D.A. Reg. No. 4-137

*Total Arsenic expressed as elemental.....	3.64%
Water Soluable Arsenic expressed as elemental.....	.246%
Copper expressed as elemental.....	2.09%

DIRECTIONS:

Apply when larvae are in 2nd, 3rd or 4th instar stages. Application in 1st stage not effective with no toxic effect on pupae. Recommended rate is 0.75 to 1.50 lbs. active ingredient per acre (9 to 18 lbs. pellets). May be applied by duster, rotary seeder, or hand broadcast. Application by bellows dusters or sling seeders is recommended, sling seeders are preferred.

Hand Broadcast Method will average 9-12 lbs. of mix per acre but applicators can check overdosage by test applying 2 quarts per 5,000 sq. ft. (100' x 50' area), to gain accurate rate indication before general application.

Aircraft Application: Recommended altitude 75' using 150 H.P. Super Cub, 115 H.P. Cub or Public Health Service AT-6. Swath width will be 25 feet and application rate should not exceed 10 lbs. per acre or per 1742 ft. long swath (1/3 mile).

WARNING: Keep away from children and domestic animals. May be fatal if swallowed. Do not get in eyes, on skin or on clothing; wash thoroughly after using. Avoid breathing of dust. As some applicators may be allergic to Paris Green compounds use of rubber gloves while hand broadcasting is recommended. Avoid contamination of feed and foodstuffs. Dispose of empty containers and spillage by burying.

BONIDE CHEMICAL CO., INC.

UTICA, NEW YORK

