

to control knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.


For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.

For control of knotweed, chickweed, red sorrel, clover, mouse-ear, spotted spurge, burdock, thistles, bur clover, violets and yarrow, apply one quart of Bonide to 20,000 sq. ft. of area.



# BONIDE

## CHICKWEED & CLOVER KILLER W/BANVEL "D"

Kills knotweed, common chickweed, red sorrel, clover, mouse-ear chickweed, curly dock, dog fennel, stitchwort, henbit, English daisy, spurge, purslane, hawkweed, lawn burweed, carpet weed, pepperweed, chicory, spurry, creeping charly, knawel, black medic, spotted spurge, burdock, thistles, bur clover, violets and yarrow.

Manufactured by **BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502**

NET WEIGHT: 1.05 LBS. (475 GMS.)  
EPA REG. NO. 100-100000-01  
EPA EST. NO. 100-100000-01  
EPA REG. NO. 100-100000-01  
EPA EST. NO. 100-100000-01  
EPA REG. NO. 100-100000-01  
EPA EST. NO. 100-100000-01

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
(See side panel for other cautions)

One Quart Net Treats up to 20,000 Sq. Ft.

CHICKWEED & CLOVER KILLER is especially effective against the majority of weed species found in turf. It is absorbed by the leaves and roots and transmitted readily to all plant parts. It kills even the deepest roots. Containing Banvel "D" it is effective in wet as well as warm weather. As a result it is effective for control of hard to kill turf annuals.

### IMPORTANT

Avoid application on or near desirable trees or other plants, or on areas where their roots may extend in locations where the weeds may be washed or moved into contact with their roots. Do not use in equipment on or near desirable trees or other plants. Do not drift of product to desirable plants. Use a coarse spray. Thoroughly clean application equipment after use. Do not re-use equipment. Destroy by burning or a safe place. Do not burn.

**CAUTION:** Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water for eyes get medical attention if necessary.

Avoid contamination of domestic or irrigation water.

### DIRECTIONS

For control of knotweed, chickweed, red sorrel, clover, mouse-ear and perennial or biennial weeds, apply one quart of Bonide to 20,000 sq. ft. of area. For control of knotweed, chickweed, red sorrel, clover, mouse-ear and perennial or biennial weeds, apply one quart of Bonide to 20,000 sq. ft. of area.

**BIRGRASS GREEN:** Apply one to two quarts per acre (1/2 to 1/3 pound dicamba per acre).

Apply application to root zone areas of trees, shrubs or ornamental plants. (See import at bottom)

For establishment of perennial grasses (including bluegrass, lawn type fescues and other special turf) and pasture grasses grow for seed, and established perennial grasses, grown for seed. Use one to two quarts per acre. To control sheep sorrel and sorrel in established grasses, make application between November 15 and April 1st or prior to boot stage. For new seeding make application in late spring after seed crop has 3 to 5 leaves. Do not use seed from treated grasses for feed or food purposes. Do not graze with animals in treated fields within 30 days before slaughter.

To control nightflowering catchfly and white cockle in established grasses make application to foliage in the spring after seed crop has 3 to 5 leaves. For new seeding make application to foliage in spring after seed crop has 3 to 5 leaves.

To control chickweeds (common, mouse-ear, perennial or stitchwort), clover curlydock, cowcockle, dog fennel, English daisy, spurge, purslane, hawkweed, lawn burweed, carpet weed, pepperweed, chicory, spurry, creeping charity, knotweed, black medic, spotted spurge, burdock, thistles, bur dock, violets and yellow.

For Merion bluegrass (other bluegrasses), lawn type fescues and other turf and pasture grasses. To control downy mildew and rust on turf and pasture grasses. Use 2.4 gallons per acre applied in fall after harvest and in spring before seed crop has more than two leaves. Do not graze meat animals in treated fields.


For fencerows, roadways and similar non-cropland. To control field bindweed, Russian knapweed, Canada thistle, leafy spurge. Use 4 to 8 gallons per acre applied to foliage and/or soil when weed is in actively growing stage. For spot treatment mix 3 to 6 ounces with one gallon of water and treat one square rod (272 sq. ft.).

To control top growth of field bindweed, Russian knapweed, Canada thistle and leafy spurge, chickweeds, curly dock, dog fennel, knotweed, clover and sheep sorrel. Use 1/2 to one gallon per acre applied as a foliar spray in spring or early summer to actively growing weeds.

For wasteland, fencerows and similar non-cropland. To control eastern persimmon. Use 4 to 8 gallons in 100 gallons water, 1/4 to 1/2 pint of spray solution per inch diameter of plant applied to ground under trees as basal treatment.

Tree Kill — To control red alder, white ash, aspen, basswood, beech, yellow paper birch, dogwood, sweet black gum, hickory, huckleberry, red sugar maple, blackack and post oak, persimmon and white pine. Apply undiluted to overlapped axe cuts anytime during year. Cut must penetrate bark and cambium (sapwood). Symptoms of injury will be noted within a few weeks but kill may take several months.

Manufactured by  
BONIDE CHEMICAL CO., INC., UTICA, N. Y. 13502



4-172

BONIDE

**CHICKWEED & CLOVER  
KILLER**  
W/BANVEL "D"

Kills knotweed, common chickweed, red sorrel, clover, mouse-ear chickweed, curly dock, dog fennel, stitchwort, henbit, English daisy, spurge, purslane, hawkweed, lawn burweed, carpet weed, pepperweed, chicory, spurry, creeping charity, knotweed, black medic, spotted spurge, burdock, thistles, bur dock, violets and yellow.

ACTIVE INGREDIENTS:

Dimethopate salt of dicamba (3,6-dichloro-o-anisic acid)	EPA Reg. No. 4.172
NEET (N-ethylmaleimide)	13.0%
NEET (N-ethylmaleimide)	2.1%
918 (N-ethylmaleimide)	84.9%

55

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
(See side panel other cautions)  
One Quart Net Treats up to 20,000 Sq. Ft.

BONIDE CHICKWEED & CLOVER KILLER is especially effective against the majority of weed species found in turf. It is absorbed through leaves and roots and transmitted rapidly to all plant parts. It kills even the deepest roots. Containing Banvel® D, it is effective in cold as well as warm weather. As a result, it is effective for control of hard to kill winter annuals.

**IMPORTANT**

Avoid application on or near desirable trees or other plants, or on areas where their roots may extend or in locations where the weed killer may be washed or moved into contact with their roots. Do not clean equipment on or near desirable trees or other plants. Prevent drift of product to desirable plants, use a coarse spray. Thoroughly clean application equipment after use. Do not reuse container. Destroy by burying in a safe place. Do not burn.

**CAUTION:** Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Store out of reach of children.

Avoid contamination of domestic or irrigation waters.

**DIRECTIONS**

**GOLF COURSES:** For use on golf fairways, tees and rough. To control chickweed (common, mouse-ear and perennial or stitchwort) clover, curly dock, dog fennel, knotweed, sheep sorrel (red sorrel) henbit, English daisy, spurge, purslane, hawkweed, lawn burweed, carpet weed, pepperweed, chicory and spurry, apply one half to one gallon per acre, 1 1/2 oz. (7 teaspoons) to 3 1/4 oz. (6 1/2 tablespoons) in 8 or 1/2 pt. of water per acre to 5,000 sq. ft. Apply as a foliar spray to actively growing weeds in sufficient water for coverage to 1/2 inch. Use 50-100 gallons per acre large areas.

**BENTGRASS GREENS:** Apply one to two quarts per acre (1/4 to 1/2 pound dicamba per acre).

Avoid application to root zone areas of trees, shrubs or ornamental plants (see important ph.)

For establishment of perennial grasses (including bluegrass, lawn type fescues and other special turf and pasture grasses grown for seed) and established perennial grasses grown for seed. Use one to two quarts per acre. To control sheep sorrel (red sorrel) in established grasses, make application between November 15 and April 1st or prior to boot stage. For new seeding make application to foliage in spring after seed crop has 3 to 5 leaves. Do not use seed from treated grasses for feed or food purposes. Do not graze meat animals in treated fields within 30 days before slaughter.

55

37