

BONIDE

Lawn Insect Control Granules

Savin 20%

**STOPS LAWN AND GRASS DAMAGE FROM SOD WEBWORMS (LAWN MOTHS),
FALL ARMYWORMS, CUTWORMS, LEAFHOPPERS, EARWIGS, ANTS, FLEAS,
MOSQUITOES, MILLEPEDS AND ADULT BLUEGRASS BILLBUG (N.Y. STATE)**

EPA Est. 4-NY-1

EPA Reg. No. 4-177

ACTIVE INGREDIENT: Carbaryl (1-naphthyl methylcarbamate) . . . 20.0%
INERT INGREDIENTS 80.0%

CAUTION: Keep Out of Reach of Children.

Harmful if inhaled or swallowed. Avoid breathing of dust or spray. Do not take internally. Avoid contact with skin. In case of contact, and before eating or smoking, thoroughly wash hands and face.

Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Avoid contamination of food, feed, water supplies, streams and ponds. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke and fumes.

DIRECTIONS

Fall Armyworms, Cutworms, Leafhoppers, Earwigs, Ants, Fleas, Mosquitoes, Millepedes — Apply at the rate of 1 lb. per 1,000 sq. ft. (5 lbs./5,000). 50 lbs. will treat 50,000 sq. ft. Water thoroughly after application to wash ingredients into soil.

Sod Webworms (Lawn Moths) — Apply at the rate of 1 lb. per 1,000 sq. ft. (5 lbs./5,000). 50 lbs. will treat 50,000 sq. ft. Soak or irrigate area to be treated to force bugs to surface before application. Do not water for 2-3 days after application. Repeat application if needed.

Adult Bluegrass Billbug (N.Y. State) — Apply at the rate of 1¼ lbs. per 1,000 sq. ft. (6.25 lbs./5,000). Water thoroughly after application to wash ingredients into soil.

50 LBS. NET WEIGHT

Buyer's guarantee limited to label claims

