

## CAUTION

Keep out of reach of children. Avoid breathing spray and contact with eyes or skin. May be absorbed through the skin. In case of contact and after use, 'vash with soap and water. Avoid contamination of feed and foodstuffs

Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 F may cause bursting.

Cover fish bowls or aquariums near use areas before use. Do not spray Maidenhair Fern, Lantana, Cycla-

Cover fish bowls or aquariums near use areas before use. Do not spray Maidenhair Fern, Lantana, Cyclamen or apply to household pets or humans.

FLORA BOMB IS A MULTI PURPOSE SPRAY for house plants, small gardens, and may be used in large gardens for spot treatment to prevent spreading of localized infestations to larger areas. The formula is designed for use on African Violets, Asters, Azaleas, Begonias, Camellias, Carnations, Chrysanthe mums, Dahlias, Delphiniums, Dogwood, English Tyy, Euonymus, Fushia, Geraniums, Crassula, Grape vine, Kentia palm, Laurel, Marigolds, Rhododendrons, Roses, Rubber plants, Snapdragons, Stocks, Wan dering Dew and Zimnias. Insecticide for aphids, spider and clover mites, exposed thrips, lace bugs, flea beetle, bag worms, tent caterpillars, leaf hoppers, rose chafer, Japanese beetles, pavement ants, and army worms. Fungicide for black spot and powdery mildew on roses, alternaria leaf spot and rust on carnations, and powdery mildew and septoria leaf spot on Chrysanthemums.

## Sevon is registered trademant Phattan is a registered tra 2 55-770, 2,55-771

P OUT OF REACH OF

WEIGHT

TOTENIS

Any (C. nauthy) N. methylcarbamate;

et [N. ttrichloromethylthior phthylmide;

tthoor (O. Dirmethyl dithrophosphate of 'afry, myccaprisucc


sis of ediorophemyli 2,2,2 trichloroethar

DIENIS

The is decided trademath of Union Carbane corp

is registered trademath of Union Carbane corp

READY TO USE INDOORS — OUTDOORS ,
JSE PLANTS (AFRICAN VIOLET)
WERS (ROSES) ORNAMENTAL
NSECTICIDE AND FUNGICIDE



## **DIRECTIONS**

Shake can before use, remove cap, hold in upright position with spray opening (at end of arrow on but ton) pointed toward object to be sprayed. Keep spray opening and can 14 (fourteen) inches away from toliage. Propellant will freeze foliage if held closer. Press button downward to release mist. Apply mist to all parts of plant, top and undersides of foliage. Stop application before droplets form on leaf surfaces or liquid starts to drip off foliage. Apply as preventative before diseases (black spot, powdery middew, alternaria leaf spot, rust, septoria leaf spot) occur and every 5-7 days during warm, moist periods when diseases are prevalent. Properly treated foliage will be protected for about one week.

Buyer's guarantee limited to label claims

c 1977 & Mfg. by

BONIDE CHEMICAL CO., INC., YORKVILLE, N.Y. 13495

