

~~ACCEPTED~~

AGRICULTURAL MITICIDE
FUNGICIDE AND INSECTICIDE ACT
AS A FOLIAR AND POISON REGISTERED
ED UNDER SOC.

BONIDE

KELTHANE EC

AGRICULTURAL MITICIDE
EMULSIFIABLE CONCENTRATE

CAUTION: FIRE AND EXPLOSION HAZARD

AMERICAN

For maximum effectiveness, Kelthane EC should be applied at the rate of 1/2 to 1 pint per acre. This will give the best control of weeds and the best economic return. Kelthane EC is a non-selective herbicide and will kill all plants. It is important to apply the product before the weeds have had time to grow and spread. The best time to apply is when the weeds are small and the soil is dry. Kelthane EC should not be applied after the first frost or during the winter months.

SUMMARY OF KELTHANE EC DOSAGE RATES AND LIMITATIONS BETWEEN LAST APPLICATION AND HARVEST

CROP	Dosage - Qts. Per Acre Min	Dosage - Qts. Per Acre Max	Rate Qts. Per 100 Gals. Dilute Sprays*	Cutoff Days Before Harvest
Alfalfa	1/2	1	1/2	45
Barley	1/2	1	1/2	45
Beets	1/2	1	1/2	45
Buckwheat	1/2	1	1/2	45
Corn	1/2	1	1/2	45
Cotton	1/2	1	1/2	45
Grazing Land	1/2	1	1/2	45
Mint	1/2	1	1/2	45
Peas	1/2	1	1/2	45
Rhubarb	1/2	1	1/2	45
Raspberries	1/2	1	1/2	45
Strawberries	1/2	1	1/2	45
Turnips	1/2	1	1/2	45
Wheat	1/2	1	1/2	45

*Dilute sprays should be applied at the rate of 1/2 to 1 pint per acre. This will give the best control of weeds and the best economic return.

For maximum effectiveness, Kelthane EC should be applied at the rate of 1/2 to 1 pint per acre. This will give the best control of weeds and the best economic return.

For maximum effectiveness, Kelthane EC should be applied at the rate of 1/2 to 1 pint per acre. This will give the best control of weeds and the best economic return.

Packed by

KELTHERM COMPANY, INC., 1000 N. W. 21st Street, Oklahoma City, Okla.

NON-SELECTIVE HERBICIDE FOR ECONOMIC POSITION
ED UNDER NO.

ACCEPTED

CA

K

RECEIVED
FEBRUARY 11
1968

CHILDREN

三

A black and white photograph showing a dark, rectangular object at the top and a curved, dark shape below it.

DIRECTIONS FOR USE