


BONIDE



INSECT SPRAY

CONTAINS MALATHION, METHOXYCHLOR, KELTHANE

EPA Reg. No. 4-203

ACTIVE INGREDIENTS:

- *Methoxychlor, Technical.....24.0%
- **Malathion.....13.0%
- ***1,1-bis(p-chlorophenyl)-2,2,2-trichloroethanol.. 3.0%
- Aromatic petroleum derivative solvents.....20.0%

INERT INGREDIENTS:.....40.0%

- *equivalent to 21.1% 2,2-bis (p-methoxyphenyl) 1,1,1-trichloroethane and 2.9% other isomers and related products
- **0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate. Cythion reg. T.M. American Cyanamid
- ***Kelthane reg. T.M. Rohm and Haas Co.

5 GALS. NET
(0.1, 30, 55, 75)

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel other cautions)

DANGER: EXTREMELY FLAMMABLE. KEEP AWAY FROM FIRE, SPARKS AND HEATED SURFACES.

DIRECTIONS

Apply with garden hose applicator, tank or hand pressure sprayer. Choose cool, calm period (early morning or evening) when plants will not receive hot direct rays of sun. Don't spray if rain is expected before leaf surfaces dry. Apply when insects appear and repeat treatment as necessary.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac, arbor vitae, juniper, hydrangea, azalea, and elm, maple, oak, dogwood, birch, boxelder and pine trees. Mix two tablespoons per gallon water, or at the rate of 3 quarts per 100 gallons water, to control birchleaf and boxwood miner, oak kermes, bagworm, spider mites, mealy bugs, four-lined leaf bugs, tarnished plant bugs, European pine shoot moth, and the following scale: oyster shell*, Euonymus, scurfy, azalea, pine leaf, magnolia, fletcher*, Florida red, juniper, monterey pine, black and soft. (*Apply when scale crawlers have settled on foliage.)

For Jap beetles, aphids, lace bug, thrips, flea beetle, plant bugs, white fly, red spider - use two teaspoons per gallon water, or at the rate of 1 quart per 100 gallons water.

For tent caterpillars, rose chafers, grasshoppers - use 4 teaspoons per gallon water, or at the rate of 2 quarts per 100 gallons water.

To control flies, mosquitoes, ants - mix 3 tablespoons per gallon, or at the rate of 5 quarts per 100 gallons water, and spray thoroughly tall grass, shrubbery, around buildings, ant hills or nests and other such places where these pests congregate or harbor, but not near food crops. (1 gallon mixed spray treats 425 sq. ft.)

Do not use on certain ferns including Boston, Maiden Hair, Pteris and Crassula.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of spray mist. Avoid contact with skin, eyes. In case of contact, wash immediately with soap and water. Any liquid that has entered through skin. Wash thoroughly after using. Do not use on painted surfaces. Keep children and pets off treated areas until washed and the glass is dry.

This product is poisonous to fish. Keep away from body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is highly toxic to bees, especially if direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

Buyer's guarantee limited to label claims.

Manufactured by
Bonide Chemical Co., Inc., Utica, N. Y.