


BONIDE

TREATS 2,500 SQUARE FEET
A 50 FOOT BY 50 FOOT LAWN AREA
NET WEIGHT 10 LBS.

Just pour in spreader and walk.
Clean, easy to use, dust free, dry granules.

DIAZINON® 2 $\frac{1}{2}$ G



Kills chinch bugs, sod webworms (lawn moths), ants, fleas, mosquitoes, bermuda and clover mites, earwigs, millipeds, fall army worms, cutworms, leafhoppers, brown dog ticks, crickets, chiggers, sowbugs, collembola (spring tails), spiders and scorpions.

EPA Reg. No. 4-209

ACTIVE INGREDIENT:

*0, 0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 2.50%

INERT INGREDIENTS: 97.50%

*Diazinon®, a registered trademark of Ciba-Geigy Corp.

CAUTION: KEEP OUT OF REACH OF CHILDREN
(See back panel cautions)



DIAZINON[®] 2 $\frac{1}{2}$ G

BONIDE DIAZINON[®] 2 $\frac{1}{2}$ G is a new and effective insecticide in dry granular form. Prevents surface feeding insect damage to lawns, and may also be used on all turf around ornamental plants, picnic areas, patios. Nothing to add! No water to mix! No spray to drift! Easily applied to all lawn areas with any spreader.

DIRECTIONS

Begin protection before lawn shows signs of surface insect injury. Serious damage can be prevented if the first application is made during late spring or early summer. DIAZINON[®] 2 $\frac{1}{2}$ G is easily applied after mowing, when grass is dry. If the infestation is very heavy, a second application will be necessary. Repeat full treatment in 7 to 10 days.

Apply at the rate of 10 lbs. per 2,500 sq. ft., or 20 lbs. per 5,000 sq. ft. Remove leaves, dead or tall grass and other debris before application. Measure lawn. Apply to measured area correct amount of DIAZINON[®] 2 $\frac{1}{2}$ G at recommended rate with drop-type spreader. Do not guess. Spread uniformly for best results. Criss-cross application is recommended; $\frac{1}{2}$ material applied in one direction, the remainder at right angles. Avoid over or under application, skips or misses.

Thoroughly wet down grass a few hours before applying and use at the recommended rate for each square foot.

The following suggested spreader settings are approximate. Adjustments must be made for condition of the spreader, rate of walk and humidity/moisture conditions. Since most machine manufacturers use only whole numbers, fractions will have to be estimated between whole numbers. Increasing numbers usually equals faster rate of application. Close shut off on turns, push shut off open when starting. Keep machine handle low or container of machine vertical to ground level. Adjustments will seldom exceed $\frac{1}{2}$ number.

SPREADER	MODEL	SETTING NO.	$\frac{1}{2}$ RATE
Patco	P220 thru P524	7 $\frac{1}{4}$	6
Lawn Beauty	18 and 24	7 $\frac{1}{2}$	6 $\frac{1}{4}$
Scotts	35 - 8	7 $\frac{3}{4}$	6 $\frac{1}{2}$
Jackson	20 and 26	7 $\frac{1}{4}$	6
Modern	360G - 1	7 $\frac{3}{4}$	6 $\frac{1}{2}$
*Patco-Rorocast	"C"	5	4 $\frac{1}{4}$
*Cyclone-Broadcast	"B"	5	4 $\frac{1}{2}$
*Brisk walking			

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. May be absorbed through the skin. In case of contact, wash thoroughly. Do not contaminate feed, foodstuffs or drinking dishes of pets. Wash after handling.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on the label. Dispose of empty container in safe place. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

BUYER'S GUARANTEE LIMITED TO LABEL CLAIMS
Manufactured by
BONIDE CHEMICAL CO., INC., UTICA, NEW YORK 13502