
USE RECOMMENDATIONS	
---------------------	--

PLANTS	PESTS CONTROLLED	AMOUNT TO MIX WITH WATER
Ornamentals - roses, camellias, honey- suckle, azaleas, other flowering plants, vines, privet, lilac and other herbace- ous shrubs, shade trees including elm,	Loopers, tent caterpillar, European pine shoot moth, birch leaf miners, thrips, red-pine sawfly, lygus and plant bugs, leafhoppers.	4 teaspoons per gal. or 2 qt. per 100 gals.
maple, ash, locust, evergreen — such as pine, spruce, juniper, arborvitae, and various nursery plants.	Aphids, soft scale, scale crawlers, mealy bugs, white fly and mites.	6 teaspoons per gal. or 1½ qt. per 50 gals.
Turf: bluegrass, Bermudagrass, dichon- dra, and other ground cover.	Sod webworms (lawn moth), cutworms, armyworms, millipeds, sowbugs (pill- bugs), snails and slugs.	6 teaspoons per gal. or 1½ qt. per 50 gals.
An outstanding insecticide for the contrange of pests, including snails and slugs on trees, flowers and shrubs. ACTIVE INGREDIENTS Mexacarbate 4: (Dimethylamino) C J xvlyi methylcarbamate Aromatic petroleum derivative solvent INERT INGREDIENTS "Zectran in Trademark of The Dow Chemical Company. U.S. Patent No. 3,084.098 WARNING: KEEP OUT OF REACH OF (Read complete warnings on side panel) IND NOT USE, POUR, SPILL OR STORE NEAR HEAT (ONE QUART NET	Interview Interview	



DIRECTIONS FOR USE

Apply ZECTRAN 1E in water sprays to ornamental plants and turf. Applications may be made with hand sprayers, hose attached sprayers or power spraying equipment. Use enough spray to wet the plants and turf areas thoroughly.

KILLS: Armyworms, cutworms, sod webworms, loopers, birch leaf miners. European pine shoot moth, tent caterpillars, red-pine sawfly, thrips, leafhoppers, white fly, mealy bugs, lygus and other plant bugs, aphids.

It scales and scale crawlers, mites, millipedes, sowbugs, snails and slugs.

FOLIAGE-FEEDING INSECTS AND MITES: Spray infested plants thoroughly at amounts indicated in the dosage table. Re-treat as necessary.

SNAILS, SLUGS AND TURF PESTS: Water turf thoroughly before treating. Spray infested area as indicated in the dosage table. Apply 1 gallon of spray per 130 to 180 sq. ft. of area. Soak the debris and soil under infested plants. Re-treat as necessary.

Note: Feeding stops soon after treatment, but complete kill may be delayed for two or three days.

WARNING: May be fatal if swallowed. Avoid eye and skin contact. Avoid breathing spray mists. Wash well after handling. If swallowed, call a physician immediately. Note to Physician: ZECTRAN 1E INSECTICIDE has chol-inergic effects — treat symptomatically. Atropine is an antidote.

Avoid contamination of feed, foodstuffs, water supplies and fish ponds. Do not use on maidenhair fern, cape chestnut, or Geraldton wax flower. Excessive dosage may cause slight leaf injury to new growth on roses, phlox and petunias.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment. or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.