



5% SEVIN

GENERAL INFORMATION

SEVIN is an outstanding insecticide that controls a wide variety of insect pests. MILLER 5% SEVIN INSECTICIDE DUST is compatible with commonly used miticides, fungicides, and insecticides. Do NOT use with lime or other alkaline material. SEVIN is not effective against mites. Where mites are a problem, use a suitable miticide.

FOR SMALL MEASUREMENTS

57 standard level tablespoons = 1 lb.
1 1/2 standard level tablespoons = 1 oz.
4 1/2 ounces = 1 cupful

DIRECTIONS

Apply MILLER 5% SEVIN INSECTICIDE DUST when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results, apply thoroughly to upper and lower leaf surfaces. If necessary repeat applications every 7 to 10 days. Apply with conventional dusting equipment. Do NOT dust too heavily or crop injury may result.

BEANS: For Control of Mexican Bean Beetle: Apply 1/4 lb. per 1,000 sq. ft. For Control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetles: Apply 1/2 lb. per 1,000 sq. ft. For Control of Corn Earworm, Armyworms and Velvet Bean Caterpillar: Apply 1/2 to 3/4 lb. per 1,000 sq. ft. For Control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 1 lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN: For Control of Corn Earworm, European Corn Borer, Fall Armyworm, Sep Beetle and Japanese Beetle: Apply 1/2 to 1 lb. per 1,000 sq. ft. For Larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to harvest on Sweet Corn. Do NOT apply within 7 days of harvest on corn used for fodder or forage.

CUCUMBERS AND SUMMER SQUASH: For Control of Pickleworm and Melonworm: Apply 1/4 to 1/2 lb. per 1,000 sq. ft. For Control of Cucumber Beetles, Squash Bugs and Flea Beetles: Apply 1/2 lb. per 1,000 sq. ft.

LETUCE: For Control of Armyworm, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus and Stink Bugs: Apply 1/2 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 3 to 7 day intervals. Do NOT apply within 3 days of harvest.

OKRA: For Control of Corn Earworm and Stink Bugs: Apply 1/4 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 5 to 7 day intervals or as necessary.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS: For Control of Colorado Potato Beetle, Flea Beetle and Leafhoppers: Apply 1/2 to 3/4 lb. per 1,000 sq. ft. For Control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs and Stink Bugs: Apply 1 lb. per 1,000 sq. ft.

BLUEBERRIES: For Control of Cherry and Cranberry Fruitworm: Apply 1/4 lb. per 1,000 sq. ft.

CRANBERRIES: For Control of Fruitworms, Fireworms, Leafhoppers, and Cutworms: Apply 1 to 1 1/4 lbs per 1,000 sq. ft. Do not apply within 1 day of harvest.

GRAPES: For Control of Grape Leaf Folder, Grape Leafhopper and Grape Berry Moth: Apply 1/2 lb. per 1,000 sq. ft. Make application when insects are first noted and repeat as necessary.

STRAWBERRIES: For Control of Meadow Spittlebug and Leaf Roller: Apply 1/4 to 1 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.

ORNAMENTALS: Arbor Vitae, Azaleas, Boxwood, Dogwood, Rose, Spirea, Yew: For Control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhoppers, Tent Caterpillar and exposed Thrip: Apply 1 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do NOT use on Boston ivy.


NOTE: SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide. Toxic to honeybees. Avoid use during flowering and pollination period. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do Not Breathe Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

Avoid Contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles. Do not re-use empty container. Destroy by burying with waste in a sanitary land fill, away from water supplies or by burning. Stay away from smoke and fumes.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use, when this product is not used in accordance with the directions as stated on this label.



Manufactured By
MILLER CHEMICAL & FERTILIZER CORPORATION

Hanover, Pennsylvania 17331/Ancor Standard Company

NET WEIGHT 3



5% SEVIN

GENERAL INFORMATION

SEVIN is an outstanding insecticide that controls a wide variety of insect pests. MILLER 5% SEVIN INSECTICIDE DUST is compatible with commonly used miticides, fungicides, and insecticides. Do NOT use with lime or other alkaline material. SEVIN is not effective against mites. Where mites are a problem, use a suitable miticide.

FOR SMALL MEASUREMENTS

1/2 standard level tablespoonfuls	= 1 qt.
1/2 standard level tablespoonfuls	= 1 oz.
4 1/2 ounces	= 1 cup.

DIRECTIONS

Apply MILLER 5% SEVIN INSECTICIDE DUST when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results, apply thoroughly to upper and lower leaf surfaces if necessary repeat applications every 7 to 10 days. Apply with conventional dusting equipment. Do NOT dust too heavily or crop injury may result.

BEANS: For Control of Mexican Bean Beetle. Apply 1/4 lb. per 1,000 sq. ft. For Control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetles. Apply 1/2 lb. per 1,000 sq. ft. For Control of Corn Earworm, Armyworms and Velvet Bean Caterpillar. Apply 1/2 to 1/4 lb. per 1,000 sq. ft. For Control of Lima Bean Pod Borer, Lygus and Stink Bugs. Apply 1 lb. per 1,000 sq. ft. Do not feed treated bean vines to dairy or meat animals.

CORN: For Control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle and Japanese Beetle. Apply 1/2 to 1 lb. per 1,000 sq. ft. For Larvae in whorl (budworm damage) and foliage feeders apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to harvest on Sweet Corn. Do NOT apply within 7 days of harvest on corn used for fodder or forage.

CUCUMBERS AND SUMMER SQUASH: For Control of Pickleworm and Melonworm. Apply 1/4 to 1/2 lb. per 1,000 sq. ft. For Control of Cucumber Beetles, Squash Bugs and Flea Beetles. Apply 1/2 lb. per 1,000 sq. ft.

LETTUCE: For Control of Armyworm, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus and Stink Bugs. Apply 1/4 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals. Do NOT apply within 3 days of harvest.

OKRA: For Control of Corn Earworm and Stink Bugs. Apply 1/4 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 5 to 7 day intervals or as necessary.

POTATOES, EGGPLANTS AND PEPPERS: For Control of Colorado Potato Beetle, Flea Beetle and Leafhoppers. Apply 1/2 to 1 1/2 lb. per 1,000 sq. ft. For Control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs and Stink Bugs. Apply 1 lb. per 1,000 sq. ft.

BLUEBERRIES: For Control of Cherry and Cranberry Fruitworm. Apply 1/4 lb. per 1,000 sq. ft.

CRANBERRIES: For Control of Fruitworms, Fleaworms, Leafhoppers, and Cutworms. Apply 1 to 1 1/2 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.

GRAPES: For Control of Grape Leaf Folder, Grape Leafhopper and Grape Berry Moth. Apply 1/2 lb. per 1,000 sq. ft. Make application when insects are first noted and repeat as necessary.

STRAWBERRIES: For Control of Meadow Spittlebug and Leaf Roller. Apply 1/4 to 1 lb. per 1,000 sq. ft. Do not apply within 1 day of harvest.


ORNAMENTALS: Arbor Vitae, Azalea, Boxwood, Dogwood, Rose, Spires, Yew. For Control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhoppers, Tent Caterpillar and exposed Thrip. Apply 1 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals as necessary. Do NOT use on Boston Ivy.

NOTE: SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide. Toxic to honeybees. Avoid use during flowering and pollination period. Injury to tender foliage may occur if applications are made when plants are wet or in the presence of high humidity of several days duration following treatment.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do Not Breathe Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact With Hands Before Eating. Take Shower & Bath After Work. Wear Regular Long Sleeve Work Clothing. Change to Clean Clothing Daily. Note to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Avoid Contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering facilities. Do not re-use empty containers. Destroy by burying with waste in a sanitary land fill, away from water supplies or by burning. Stay away from smoke and fumes.

The use of this material being beyond our control and involving elements of risk to human beings, animals and vegetation, we do not make any warranty, express or implied, as to the effects of such use. When this product is to be used in accordance with the directions as stated on this label.



Manufactured By
MILLER CHEMICAL & FERTILIZER CORPORATION
Hanover, Pennsylvania 17331 An Alco Standard Company