

BUGONeX

DIRECTIONS: Thoroughly spray both upper and lower leaf surfaces and other injected plant parts. Apply with pressure sprayers, hose-end-type sprayer, sprinkling can, Repeat as necessary. Do not apply to food crops within 3 days of harvest or as otherwise specified.

ORNAMENTALS, ROSES, FLOWERS, EVERGREENS, SHRUBS AND TREES: Aerial, 1 1/2 to 2 tablespoons per gallon of water. **Aphids:** 1 1/2 to 2 tablespoons per gallon of water. **Aphids, Mites, Scale, Spider Mites, Japanese Beetle, Adults, Mealy Bug, Whitefly, Citrus Red Scale, Pear Thrips, Fruit Tree Scale, Bagworm, Tent Caterpillar, Elm Bark Beetle, Red-banded Sawfly, Dogwood Borer, Emerald Ash Borer, Elm Leaf Miner, Yellow-bellied Sapsucker, Oyster Shell Scale, European Scale, Starling Scale, etc.**

VEGETABLES, TOMATOES, PEPPERS, CUCUMBERS, BEANS, CABBAGE AND CAULIFLOWER: Flea Beetles, Aphids, Mexican Bean Beetle, Cucumber Beetles, Leafhoppers, Leaf Miners, etc. **NOTE:** Do not use within 14 days of harvest on cabbage and cauliflower.

OUTDOOR PESTS: For **FLIES** and **MOSQUITOES:** Use 2 tablespoons per gallon of water. Spray patios and foliage and grass around patios and porches at the rate of one gallon per 1000 square feet.

For CLOVER APHIDS, Mites, Spiders, HIGGERS, GARDEN SLUGS, GRASSHOPPERS, AND CERTAIN LAWN INSECTS: Use 8 tablespoons per gallon of water. Spray lawns between window door sills, lawns, etc. at the rate of one gallon of diluted spray per 100 sq. ft.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin, wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep out of reach of children and domestic animals. **NOTE TO PHYSICIAN:** Repeated prolonged use of Malathion may cause **CHOLINESTERASE INHIBITION, ATROPINE IS ANTIDOTAL.** Do not use or store near heat or open flame. Keep container closed when not in use. Do not use on Boston, Maidenhair and Pteris ferns, certain species of Junipers, and some species of Crassula. Destroy container when empty.

NOTE: 3 tablespoons 1 1/2 to 2 gallons of water. 1 ounce 1/2 to 1 gallon of water. 1/2 ounce 1/4 to 1/2 gallon of water. 1/4 ounce 1/8 to 1/4 gallon of water.

GUARANTEE: Effective for purpose claimed.

ACTIVE INGREDIENTS: Malathion 50%
Methylated Naphthalenes 47%

INERT INGREDIENTS: 3%

***G, O-dimethyl dithiophosphate & diethyl mercaptosuccinate**
**Not considered active ingredients in Canada
CANADIAN Reg. No. 7944 P.C.P. A. U.S.D.A. Reg. No. 88 12

NOTICE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Write for free catalog and prices on other OneX products. If dealer cannot supply your needs, order direct from manufacturer.

50% MALATHION BUG KILLER LIQUID CONCENTRATE

SIMPLY ADD TO WATER AND SPRAY PLANTS
CAUTION: Keep out of the reach of children
See warnings on back panel.

CONTENTS:
7 OUNCES
MAKES UP TO
8 GALLONS
OF MIXED
SPRAY IF
USED AS
DIRECTED



79¢
KILLS
CHEWING
SUCKING
CRAWLING
& FLYING
BUGS

CONTROLS MANY GARDEN PESTS
FOR BETTER PLANT PROTECTION

[88-12]

ACCEPTED
MAR 6 1970

UNDER THE FEDERAL INSECTICIDE ACT
AND THE FEDERAL FERTILIZER ACT
AS ENFORCED BY THE ENVIRONMENTAL
PROTECTION AGENCY
FOR THE FOLLOWING COMMENTS.

[88-12]

NET CONT. 2 FL OZS. MFG. IN U.S.A. FOR HYDROPONIC CHEM. CO. INC. (COPY 21 OHIO) REG. U.S. PAT. OFF.

DIRECTIONS: Thoroughly spray both upper and lower leaf surfaces and other infected plant parts. Apply with pressure sprayers, hose-end-type sprayer, sprinkling can. Repeat as necessary. Do not apply to food crops within 3 days of harvest or as otherwise specified.

ORNAMENTALS: ROSES, FLOWERS, EVERGREENS, SHRUBS AND TREES. Aphids, 1½ to 2 teaspoonfuls per gallon of water. Mites, Red Spider Mites, Spruce Mites, Japanese Beetle Adults, Mealy Bugs, Leafhoppers, Flea Beetles, White Fly, Lace Bug, Bagworm, Tortoise, Plant Bug, Birch, Boxwood and Elm Leaf Miners, Tent Caterpillars: 1 tablespoonful per gallon of water. Soft Brown Scale, Green Scale Crawlers, Oyster Shell Scale, Euonymus Scale, Azalea Scale: 1½ to 2 tablespoonfuls per gallon of water.

VEGETABLES: TOMATOES, PEPPERS, CUCUMBERS, BEANS, CABBAGE AND CAULIFLOWER. Flea Beetles, Aphids, Mexican Bean Beetles, Cucumber Beetles, Leafhoppers, Red Spider Mites, Thrips, Cabbage Worms: 1 tablespoonful per gallon of water. Do not use within 7 days of harvest on cabbage and cauliflower.

OUTDOOR PESTS: For **FLIES** and **MOSQUITOES:** Use 5 tablespoonfuls per gallon of water. Spray patios and foliage and grass around patios and porches at the rate of one gallon per 1000 square feet. For **CLOVER MITES, ANTS, SPIDERS, CHIGGERS, CRICKETS, EARWIGS** and certain **LAWN INSECTS:** Use 8 tablespoonfuls per gallon of water. Spray foundations, window-door sills, lawns, picnic area at the rate of one gallon of diluted spray per 100 sq. ft.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep out of reach of children and domestic animals. **NOTE TO PHYSICIAN:** Repeated, prolonged careless use of Malathion may cause **CHOLINESTERASE INHIBITION. ATROPINE IS ANTIDOTAL.** Do not use or store near heat or open flame. Keep container closed when not in use. Do not use on Boston, Maidenhair and Pteris ferns, certain species of Junipers, and some species of Crassula. Destroy container when empty.

NOTE: 3 teaspoonfuls, 1 tablespoonful, 2 tablespoonfuls: 1 ounce. Canadian Users: 1 U. S. Gallon: 5/6 Imperial Gallon.

GUARANTEE: Effective for purpose claimed.

ACTIVE INGREDIENTS: *Malathion 50%
**Methylated Naphthalenes 42%

INERT INGREDIENTS: 8%
*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate
**Not considered active ingredients in Canada.

CANADIAN Reg. No. 7944 P.C.P. Act. U.S.D.A. Reg. No. 88-12.
NOTICE: Sellers guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Write for free catalog and prices of other "ONeX" products. If dealer cannot supply your needs, order direct from manufacturer.

M.A. N.E. CHEM. CO. INC. (COPY 21 OHIO) REG. U.S. PAT. OFF. PRINTED IN U.S.A.

BUGONeX

**50% MALATHION
BUG KILLER
LIQUID CONCENTRATE**

CAUTION: Keep out of the reach of children. See warnings on back.

CONTENTS
2 OUNCES
MAKES UP TO
8 GALLONS
OF MIXED
SPRAY IF
USED AS
DIRECTED



HYDROPONIC
ONeX
CHEMICAL COMPANY

**CONTROLS MANY GARDEN PESTS
FOR BETTER PLANT PROTECTION**

79¢

KILLS
CHEWING,
SUCKING,
CRAWLING
& FLYING
BUGS

CARMEL ETHICAL CHEMICALS

Formula MU-17
(USDA Registration No 5011-71)

(A highly concentrated insecticide to be diluted with fuel oil or special
(fogging oil, for use in Dyna-Fog insecticide applicators, or other thermal
(aerosol fogging machines.

(ACTIVE INGREDIENTS

100%

(Petroleum distillate

80%

(Naled--1,2 Dibromo-2,2-dichloroethyl dimethyl
(phosphate

20%

(SKULL & BONES

DANGER - POISON

SKULL & BONES

(WARNING

(Keep out of the reach of children. See right side panel for additional warnings,
(and antidote statements.

(DIRECTIONS

This concentrate should be diluted at a rate of 1 gallon of concentrate in
19 gallons of No. 2 Fuel Oil, or special fogging oil for use in Dyna-Fog
insecticide applicators, or other thermal aerosol fogging machines. Fog should
be applied by pest control operators or trained personnel.

For houseflies, gnats, certain other nuisance insects such as flying moths,
Craneflies, midges, black flies in residential areas, municipalities, tidal
marshes, swamps, woodlands, livestock pastures, feed lots and pastures
including Dairy cattle. ^{pastures} It is not necessary to avoid farm buildings, feed
or forage areas.

ACCEPTED