

DIRECTIONS

FOR DUSTING: Apply with suitable duster as it comes from the package to both surfaces of the leaves as uniformly as possible.

FOR SNAP AND BUTTER BEANS: To control Mexican Dean Beetle, Bean Leaf Beetle and Cucumber Beetle.

FOR CABBAGE, COLLARDS AND KALE: To control Cabbage Worm, Cabbage Looper and Flea Beatle

FOR CUCUMBERS AND MELONS: To control Cucumber Beetle and to reduce damage by Pickle Worm.

FOR POTATOES AND TOMATOES: To control Colorado Potato Beetle, Aphids and Fiea Beetle.

FOR PUMPKINS AND SQUASH: To control Cucumber Beetle, Squash Vine Borer and to reduce damage by Pickle Worm.

FOR OKRA: To control Cucumber Beetle and Flea Beetle.

FOR PEAS: To control Pea Aphid.

FOR PEPPERS: To control Aphids, Flea Beetle and Pepper Maggot.

CAUTION

Do not apply within one day of harvest.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product, other than indicated on the label. Euger assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instruction.



NET WEIGHT

50 POUNDS

1% ROTENONE DUST

ACTIVE INGREDIENTS:

,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	-
Rotenone	1.00%
	2.00%
INERT INGREDIENTS	97.00%
	100.00%

CAUTION Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 148-1011



ACCETTED