

ACCEPTED

MAY 28 1970

UNDE... ICIDE
FUNGIC... IDE ACT
FOR ECON... IN REG. STER-
ED UNDER NO... SUBJECT
TO ATTACHED COMMENTS.

143-1070

DIRECTIONS FOR USE

CABBAGE, BROCCOLI, AND BRUSSELS SPROUTS: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface-Feeding Cutworms and Aphids—Apply at the rate of 10 lbs. per acre. Do not apply to cabbage after the heads begin to form. Do not apply to Broccoli and Brussels Sprouts after the edible parts begin to form.

COLLARDS AND KALE: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface-Feeding Cutworms and Aphids—Apply at the rate of 10 to 20 lbs. per acre. Do not apply more than 4 times nor more than 3 times after thinning. Do not apply within 14 days of harvest when 10 lbs. per acre is used, nor within 21 days of harvest when 20 lbs. per acre is used.

TOMATOES: To control Tomato Hornworm, Tomato Fruit Worm, Sespentine Leaf Miner, Flea Beetle, Cabbage Looper, Blister Beetle and Aphids—Apply at the rate of 20 lbs. per acre. Do not apply within 10 days of harvest.

PLANTERS

**Toxaphene - Parathion
Dust**

ACTIVE INGREDIENT:

Toxaphene-Technical Chlorinated Camphene 20.00%
(Chlorine Contents 67-69%)

Parathion 2.00%

INERT INGREDIENTS 78.00%

100.00%

O,O-Diethyl O-p-nitrophenyl thiophosphate

DANGER KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-96



WARNING

May be fatal if swallowed, inhaled, or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep away from children and domestic animals. Do not use on cucurbits. Do not use on feed or forage.

This product will kill fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Keep out of lakes, streams, or ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Apply this product only as specified on this label.

Do not re-use empty container. Destroy it by burying with waste in a safe place.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.