

MAY 13 1968

120878

[148-1021]

### GENERAL DIRECTIONS

Use this product only for recommended purposes and at recommended dosages.

**CABBAGE AND BROCCOLI** — For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhopper, Salt-Marsh Caterpillar, Mites, Cutworm (climbing), Dipterous Leafminer (adult), and Lygus Bugs, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 1 day of harvest.

**KALE, COLLARDS, TURNIP TOPS, MUSTARD GREENS, AND BRUSSELS SPROUTS** — For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Salt-Marsh Caterpillar, Mites, Cutworm (climbing), Dipterous Leafminer (adult), False Chinch Bugs, and Lygus Bugs, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 3 days of harvest.

**LETTUCE** — For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Corn Earworm, Cutworm (climbing), Dipterous Leafminer (adult), Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Mites, Lygus Bugs, Salt-Marsh Caterpillar, and False Chinch Bug, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 2 days of harvest.

**SPINACH** — For Aphids, apply at the rate of 8 to 12 lbs. per acre. For Cabbage Looper, Imported Cabbage Worm, Leafhoppers, Mites, Dipterous Leafminer (adult), Cutworm (climbing), Salt-Marsh Caterpillar, and False Chinch Bug, apply at the rate of 15 to 25 lbs. per acre. Repeat applications when necessary. Do not apply within 4 days of harvest.

# 2% PHOSDRIN\* DUST

ACTIVE INGREDIENTS	BY WEIGHT
Alpha Isomer of 2-Carbomethoxy-1-methylvinyl	1.1%**
Related Compounds	0.8%**
INERT INGREDIENTS	98.0%
TOTAL	100.0%

\*\*Equivalent to 2.0%w Phosdrin Insecticide

**DANGER KEEP OUT OF REACH OF CHILDREN.**

USDA REG. No. 1868-98

P-C

**PLANTERS CHEMICAL COMPANY**

Division of Thompson-Hayward Chemical Company

**NORFOLK, VA. FAYETTEVILLE, N. C. ALBANY, GA.**

### PRECAUTIONS IN USING

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected humans and animals should be kept out of treated fields on the day of treatment. Keep out of the reach of children and domestic animals. Do not contaminate food or feed products.

This product will kill fish and wildlife. Do not apply when weather conditions favor drift from areas treated. Do not contaminate streams, lakes or ponds with this material.

### WARNING

**MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN.** Do not swallow or get in eyes or on skin. Wear clean natural rubber gloves and clean protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Phosdrin insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes, immediately flush with water for at least ten minutes and get medical attention. In case of spillage on person or clothing, immediately remove all contaminated clothing and wash skin with soap and water; wash clothing before reuse.

### NON-WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Trademark of Shell Chemical Co.