

**ANTIDOTE
AND FIRST AID TREATMENT**

ATROPINE IS THE EMERGENCY ANTIDOTE FOR METHYL PARATHION POISONING. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use.

CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from vicinity of the Methyl Parathion, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into eyes, wash immediately with running water for at least ten minutes. IF WARNING SYMPTOMS APPEAR: Administer two 1/100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR (See WARNING SYMPTOMS below). Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport the patient immediately to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals, in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until the time for cholinesterase regeneration has been allowed as determined by blood test.

**PLANTERS
METHYL PARATHION
EMULSIBLE CONCENTRATE**

CONTAINS 2.0 LBS. METHYL PARATHION PER GAL.

ACTIVE INGREDIENTS:	BY WEIGHT
O,O-Dimethyl O-p-nitrophenylthiophosphate	24.9%
Petroleum Hydrocarbons	68.9%
INERT INGREDIENTS:	6.2%
TOTAL	100.0%

DANGER KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-109

See side panels for Antidote and Warning statements.

GENERAL DIRECTIONS

Planters Methyl Parathion Emulsible Concentrate can be used in conventional hydraulic sprayers, low-volume ground applicators or airplane sprayers.

FOLIAGE TREATMENT: Mix the required amount of Methyl Parathion Emulsible Concentrate in sufficient water for uniform coverage and time application with state and federal recommendations.

USE AND DIRECTIONS FOR APPLICATION

COTTON: Aphids, bollweevils, leafworm, red spider mite, fleahoppers and thrips. Use one to two pints of Planters Methyl Parathion Emulsible Concentrate per acre. Application should not be made within 15 days of hand picking.

MIXING DIRECTIONS

Mix with water by agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

PLANTERS CHEMICAL CORPORATION

Norfolk, Va.

Fayetteville, N. C.

Albany, Ga.

ACCEPTED
Aug 9, 1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. 148-1028
TO ATTACHED COMMENTS

148-1028

Do not apply to special animals and volatile fields with protective material. Do not feed food or feed empty containers thoroughly.

POISONOUS RAPIDLY CONTACTS CEPTIBILI Do not swallow vapors. Wear goggles and goggles or respirator Methyl P handling: flush with accidental and water large amount a concentrate

Seller with reasonable with direct any other express instruction foreseeal

PLANTERS METHYL PARATHION EMULSIBLE CONCENTRATE

CONTAINS 2.0 LBS. METHYL PARATHION PER GAL.

ACTIVE INGREDIENTS:	BY WEIGHT
O,O-Dimethyl O-p-nitrophenylthiophosphate	24.9%
Petroleum Hydrocarbons	68.9%
INERT INGREDIENTS:	6.2%
TOTAL	100.0%

DANGER KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-109

See side panels for Antidote and Warning statements.

GENERAL DIRECTIONS

Planters Methyl Parathion Emulsible Concentrate can be used in conventional hydraulic sprayers, low-volume ground applicators or airplane sprayers.

FOLIAGE TREATMENT: Mix the required amount of Methyl Parathion Emulsible Concentrate in sufficient water for uniform coverage and time application with state and federal recommendations.

USE AND DIRECTIONS FOR APPLICATION

COTTON: Aphids, bollweevils, leafworm, red spider mite, fleahoppers and thrips. Use one to two pints of Planters Methyl Parathion Emulsible Concentrate per acre. Application should not be made within 15 days of hand picking.

MIXING DIRECTIONS

Mix with water by agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

PLANTERS CHEMICAL CORPORATION

Norfolk, Va.

Fayetteville, N. C.

Albany, Ga.

PRECAUTIONS IN USING

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. When necessary to enter treated Cotton fields within 24 hours after application, protected clothing should be worn. To protect fish and wildlife, do not contaminate streams, lakes, or ponds, with this material. Keep out of reach of children and domestic animals. Do not contaminate food or feed products. Keep away from heat or open flame. To prevent re-use, empty containers to be decontaminated and destroyed. Decontaminate by washing thoroughly with a strong solution of alkali and detergent and rinse with water.

DANGER!

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY, WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING.

Do not swallow or get in eyes, on skin, or on clothing. Do not breathe dust or vapors. Wear clean natural rubber gloves; clean, waterproof protective clothing, and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Methyl Parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If material gets into eyes, immediately flush with water for at least ten minutes and get medical attention. In case of accidental contact, remove all contaminated clothing and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

NOTICE OF WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.