



DIRECTIONS FOR USE

EDB 40 is a preparation made especially for the control of ROOTKNOT NEMATODES, WIREWORMS and SEED-CORN MAGGOTS. For TOBACCO and SWEET CORN, a waiting period of 2 weeks should be allowed before planting.

Before treatment the soil should be prepared as for regular planting. It should be well pulverized, smooth and moist. Old roots and stumps should be removed. The moisture of the soil should be about as needed for good planting conditions.

Soil should be treated with this product 2 weeks in advance of planting or transplanting. For broadcast treatment, EDB 40 should be injected into the soil at least 8 inches beneath the surface with the injectors spaced 10 to 12 inches apart. Apply 15 gallons per acre. The furrows behind the injectors should be closed and packed immediately to retain the fumigant in the soil.

For row or furrow treatment, apply 7½ gallons of EDB 40 per acre in the row or furrow, (example, if a 24 inch strip were treated and the rows were 48 inches apart, then the dosage per acre for row treatment would be ½ the dosage per acre for overall treatment). The injectors should be made at least 8 inches beneath the surface.


EDB 40 should not be applied to soil within 1 foot of growing plants.

WARNING — FLAMMABLE

Keep away from fire, heat and open flame lights. Keep packages closed. Leaking packages should be removed to a safe place.

STORE CONTAINERS IN THE SHADE.

E. D. B. 40



FOR THE CONTROL OF NEMATODES, WIREWORMS AND SEED-CORN MAGGOTS

ACTIVE INGREDIENTS:

Ethylene Dibromide	40.00%
INERT INGREDIENTS:	60.00%
	<hr/> 100.00%

1 gallon of EDB 40 contains 3.67 pounds of Ethylene Dibromide.

DANGER KEEP OUT OF REACH OF CHILDREN.

SEE OTHER WARNINGS ON BACK PANEL

SEE SIDE PANELS FOR WARNINGS & CAUTION STATEMENTS

EPA REG. No. 148-1041 IP 8-70 EPA Est. No. 148-GA-1

WARNING

Vapor Harmful — Absorbed Through Skin.

Avoid Breathing Vapor. Avoid Contact with Eyes, Skin or Clothing. Do Not Take Internally.

In case of contact, immediately remove all contaminated clothing, including shoes and wash skin with soap and plenty of water; flush eyes with water for at least 15 minutes and get medical attention. Wash clothing and air shoes thoroughly before re-use.

If illness results from inhalation, remove to fresh air and give medical attention.

If swallowed, call a physician. Induce vomiting by giving an emetic such as 2 table-spoonsful of salt in a glass of water.

Decontaminate containers by burying unused product at least 18 inches deep and away from water supplies. Allow inside of thoroughly drained container to air dry away from children. Destroy container to prevent re-use.

Not for use or storage in or around the home.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.