

DIRECTIONS

This material is intended for use as a dust on peanuts for the control of Cercospora leaf-spot. Applications should be made at the rate of 20 to 25 pounds per acre with a dusting machine capable of depositing a thin uniform coating of dust on all above ground plant surfaces. Begin applications when disease first appears in the area and repeat at 14 day intervals for a total of three to four applications.

CAUTION

Avoid breathing dust. Avoid contamination of feed and foodstuffs. Harmful if swallowed.

Applications of dusting or spraying materials containing sulphur should not be made just prior to or during periods of excessively high temperatures.

COPPER - SULFUR PEANUT DUST

ACTIVE INGREDIENTS

Sulphur 80.00%

Copper (in copper oxide form)
expressed as metallic 3.40%

INERT INGREDIENTS: 16.60%

TOTAL 100.00%

CAUTION KEEP OUT OF REACH OF CHILDREN.

USDA REG. No. 1868-155

USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES, OR OTHER UNDESIRABLE RESULTS.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.