

user before applying sulphur.  
During periods of high temperature sulphur may burn foliage and fruit. Do not make sulphur applications in excessively high temperatures. Do not use FLOWABLE SULPHUR with water within 4 weeks of an oil application.

### DIRECTIONS FOR MIXING BEFORE USING, STIR UNTIL SMOOTH

Put recommended amount into partially filled spray tank. Keep agitation running during filling and spraying operations. DO NOT ALLOW MIXTURE TO STAND. Failure to maintain agitation will cause FLOWABLE SULPHUR to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of FLOWABLE SULPHUR act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulphur in any form is corrosive material. To reduce the effect, equipment should be flushed daily.

### CAUTION

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing.

Do not contaminate feed or foodstuffs. Do not apply to any area not specified on label.


Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.

Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### NOTICE

**Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.**

# FLOWABLE SULPHUR



ACTIVE INGREDIENT:	
Sulphur as S	52.0%
INERT INGREDIENTS:	48.0%
Total	100.0%

(Contains 6.0 pounds Sulphur per gallon)

## CAUTION: KEEP OUT OF REACH OF CHILDREN

EPA Reg. No. 148-1160

CG-4-75

EPA Est. No. 148-



THOMPSON-HAYWARD  
CHEMICAL COMPANY

P.O. BOX 2383 KANSAS CITY KANSAS 66110

Use of this material in a manner other than in accordance with this label may cause serious plant, animal or personal injury, excessive residues, unsatisfactory control or other unintended consequences. Unless otherwise specified for specific tree fruits, dosage rates are given as pints of FLOWABLE SULPHUR per 100 gallons water for use in a thorough coverage spray.

### APPLES:

Scab Pre-bloom through Calyx sprays	2 quarts
Cover sprays	2 quarts
Powdery Mildew	
Pre-bloom through Calyx sprays	2 quarts
Cover sprays	2 quarts

### PEACH:

Powdery Mildew, Brown Rot, Scab:	
Pink and bloom sprays	2 quarts
Petal fall, shuck and cover sprays	2 quarts

### CITRUS: (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines):

Rust Mite	1 to 4 pints
-----------	--------------

Do not apply within 4 weeks of an oil application.

### PEANUTS:

Rust: Apply 2 to 4 pints per acre in sufficient water for thorough coverage. Apply when disease is just observed on plants. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

**CAUTION:** Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality.