# TO USE AS DUSTER:

Punch out lid liner. Hold container in horizontal position and squeeze container. Changing the position of the container varies the amount discharged.

# FOR BEST RESULTS:

- 1. Apply when you see the first pest appear on a calm day, preferably in the morning on lawns.
- 2. Read caution label.
- 3. Measure erea to be treated for vegetables. Apply dust evenly work into top 4-6" of soil.
- 4. Wash exposed skin surfaces with soap and water after using.

# **CAUTION:**

Avoid inhalation of dust. Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy container when empty.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label. ACCEPTED

ACCEPTED

AAC 2 1/57/

CHLORIDA PRODUCT

ON ECONOMIO POISON REGISTERS

EQ UNDER NO.

EQ ATTENDADO NAMED POISON

SUBJECT

FOR ATTENDADO NAMED POISON

BY ATTENDADO NAMED POISO

READY TO USE • APPLY AS A DUST

# **CAUTION:**

KEEP OUT OF REACH OF CHILDREN SEE CAUTION ON LEFT PANEL

ACTIVE INGREDIENTS: Technical Chlordane\* 10%
INERT INGREDIENTS 90%

\*Equivalent to 6% octachloro-4,7-methanotetrahyroindane and 4% related compounds.

U.S.D.A. Reg. No. 168-319

NET CONTENTS 1 LB.

WASATCH CHEMICAL COMPANY

SALT LARE LITE OF SOC. 4.

168-319

# LAWNS

# INSECTS CONTROLLED

Grasshoppers, Ants, Armyworms, Fall Armyworms, Pillbugs, Sowbugs, Cicada-Killer Wasps, Wild Bees, Chinch Bugs, False Chinch Bugs, Cutworms, European Chafer Grubs, Masked, Chafers, Grubs, Japanese Beetle Grubs, Mole Crickets, Sod Webworms, White Grubs, Wireworms.

## APPLICATION

Apply 4 ounces of MORGRO CHLORDANE 10% DUST per 100 sq. ft. of lawn area (10' x 10'). Apply when grub damage begins to appear, or when you see the moths flying in the lawn. After applying MORGRO CHLORDANE 10 DUST to your lawn, you should water normally. Let the grass dry before children and pets play on the treated area.

# VEGETABLES

Eggplant, Cucumbers, Collard, Cauliflower, Cantaloupe, Cabbage, Broccoli, Table Beets, Beans, Kale, Onion, Peppers, Potatoes, Sweet Corn, Sweet Potato, Watermelon.

#### INSECTS CONTROLLED

Wireworms, Mole Crickets, Root Maggots, White Fringed Beetles, Grubs, Pepper Maggots, White Grubs, Root Worms.

### **APPLICATION**

Apply as a preplanting broadcast soil treatment or at the time of planting, 3 oz. of MORGRO CHLORDANE 10% DUST per 100 sq. ft. of area to treat. Broadcast on soil and thoroughly work into the upper 4 tc 6 inches of soil.

READ CAUTION LABEL.

# OUTSIDE USE

#### INSECTS CONTROLLED

Ants, Box Elder Bugs, Earwigs, Cockroaches.

#### **APPLICATION**

Cust along runways, in cracks in cement walks and patios: around garbage containers; under porches; around outside house foundations; around the base of ornamental plants. Repeat as needed when pests reappear.

02

۵

• •