

CENTER PANEL

Net
Contents
1 lb.

MORGRO

ROSE DUST

INSECTICIDE - MITICIDE - FUNGICIDE

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN
SEE CAUTION ON LEFT PANEL

*revised
change to 1.5%
in letter 1-17-68*

ACTIVE INGREDIENTS:

*CARBARYL (1-naphthyl N-methylcarbamate).....	5.0%
ROTENONE.....	.1.5%
OTHER CUBE RESINS.....	7.0%
**ZINEB.....	5.0%
INERT INGREDIENTS.....	87.0%

1.5

TOTAL..... 100.0%

103.5

*"Sevin" is a trademark of Union Carbide Corp.
1-naphthyl N-methylcarbamate, U. S. Pat. No. 2,903,478
**Zinc Ethylene Bisdithiocarbamate - Rohm & Haas Co.

U.S.D.A. Reg. No. 168-X

Manufactured by:
WASATCH CHEMICAL COMPANY
SALT LAKE CITY, UTAH

ACCEPTED

JUL 6 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 168-389 SUBJECT
TO ATTACHED COMMENTS

LEFT PANEL

TO USE AS DUSTER:

Push out lid liner. Hold container in horizontal position and squeeze container. Changing the position of the container varies the amount discharged.

FOR BEST RESULTS:

1. Apply the dust to the underside, as well as top side of treated foliage when pests first appear. Repeat if needed per directions and cautions at 7 - 10 day intervals.
2. Apply the dust until you see a light film appear on all surfaces of the treated plants.
3. Read the caution on usage of this product.
4. After application, wash all exposed skin surfaces with soap and water.
5. Take cap off and pour into duster when using the refill container.

CAUTION:

Harmful if swallowed or inhaled.
Do not take internally.
Avoid breathing dust.
Avoid prolonged and repeated contact with skin.
Wash thoroughly after handling.
Change contaminated clothing daily.
Avoid contamination of food and foodstuffs.
Keep out of reach of children.

NON-WARRANTY: Manufacturer or seller is not liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically recommended on this label.

RIGHT PANEL

PESTS & DISEASES CONTROLLED ON ROSES, ANNUAL FLOWERS & ORNAMENTAL SHRUBS:

Rose Aphids, Ants, Beetles, such as Japanese, June, Flea, Leaf and Rose chafer Beetles; Boxelder Bugs, Leaf Hoppers, Scales, such as Oyster, Lecanium and Forbes; Spider mites, Pillbugs (Sowbugs), Leaf Rollers, Mealy Bugs, Cicada, Plant Bugs, Psyllids, Tent Caterpillars, Exposed Thrips, Grasshoppers, Downey Mildew, Clover Mites, Fleas, Spittlebugs, White Flies, Slugs, Rose Curculio, Lace Bugs, Black Spots.

FOR BEST RESULTS ON MILDEN: Start treatment before the disease appears. To reduce foliage disease surface water, do not sprinkle foliage.

Do not use on Boston Ivy.