

FOR PERENNIAL WEEDS* IN PASTURE, RANGELANDS AND
NON CROP LANDS SUCH AS FENCE ROWS,
DRAINAGE DITCH BANKS, RIGHTS OF WAY

WEEDS	Amount BO-RID E-4 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
CANADA THISTLE AND PEREN NIAL SOW THISTLE	1.0	0.5	Apply in late summer or early fall when weeds are in flower.	
BIND WEED	1.0	0.5	Apply in late summer or early fall when weeds are in flower.	
WHITE TOP HOARY CRESS*	1.0	0.5	Apply in late summer or early fall when weeds are in flower.	
RUSSIAN KNAP WEED	1.0	0.5	Apply in late summer or early fall when weeds are in flower.	Stops top growth and kills top growth. If top growth is not killed, re-apply if necessary.

*Perennial weeds such as Canada Thistle may not be controlled by
this application. Top growth and reseedling usually will be pre-
vented by water treatment generally gives good control.

CAUTION

Keep out of contact with skin, eyes, nose, mouth, and clothing. Harmful if swallowed.
Avoid contact with eyes. Keep out of any body of water.
Do not use in areas where water may be used for drinking.
Do not use in areas where water may be used for irrigation.
Do not use in areas where water may be used for livestock.
Do not use in areas where water may be used for wildlife.
Do not use in areas where water may be used for recreation.
Do not use in areas where water may be used for agriculture.
Do not use in areas where water may be used for industry.


Do not use in areas where water may be used for drinking.
Do not use in areas where water may be used for irrigation.
Do not use in areas where water may be used for livestock.
Do not use in areas where water may be used for wildlife.
Do not use in areas where water may be used for recreation.
Do not use in areas where water may be used for agriculture.
Do not use in areas where water may be used for industry.

Do not use in areas where water may be used for drinking.
Do not use in areas where water may be used for irrigation.
Do not use in areas where water may be used for livestock.
Do not use in areas where water may be used for wildlife.
Do not use in areas where water may be used for recreation.
Do not use in areas where water may be used for agriculture.
Do not use in areas where water may be used for industry.



BO-RID K.O.

WEED KILLER E-4



LOW VOLATILE ESTER

2,4-D 4 LBS. PER GALLON

CAUTION Keep Out Of Reach Of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS

**KILLS WEEDS IN PASTURES, CROP and NON-CROP
AREAS, FENCE ROWS and RIGHT-OF-WAYS**

ACTIVE INGREDIENTS	
2,4-dichlorophenoxyacetic acid (ester) 67.0	
(Equivalent to 2,4-dichlorophenoxyacetic acid 44.4)	
INERT INGREDIENTS	
	33.0
	100.0*

EPA REG. NO. 100-100000 NET GALLONS

The R. H. Bogle Company

ALEXANDRIA, VA
JACKSONVILLE, FLA. MEMPHIS, TENN.
ALVA, OKLA.

WHEAT
BARLEY
OATS
RICE
CORN
SUGAR BEET
CORN
PRE
EMER
GENCE

CORN
SOR
GHUMS
POST
EMER
GENCE

PAS
TURES,
RANGE
LAND

**PARTIAL LIST OF WEEDS CONTROLLED
BY BO-RID E-4**

Canada Thistle	Leafy Spurge	Shepherd's Purse
Bindweed	Mustards	Wildflower
Burdock	Stinkweed	Wild Carrot
Carpet Weed	Purslane	Wild Mustard
Cocklebur	Radiata Vetch	
Corn Dock		
Handweed		
Ironweed		
Leafy Spurge		
Mustards		
Stinkweed		
Purslane		
Radiata Vetch		

MIXING INSTRUCTIONS

Determine the amount of BO-RID E-4 to be used by multiplying
the number of acres to be treated by the amount of
BO-RID E-4 per acre of the various weeds to be controlled. Mix the
amount of BO-RID E-4 with the amount of water required to make
the amount of water to be treated. BO-RID E-4 should be applied
to the weeds to be treated.


RENTAL WEEDING SERVICE...
 WEEDS...
 WEEDS...

WEEDS	RENTAL WEEDING SERVICE	STATE OF APPLICATION	REMARKS
CANADA			
THISTLE			
ANNUAL			
PERENNIAL			
SORGHUM			
THISTLE			
BIND WEED			
WHEAT			
HOARY CRAB			
RUSSIAN KNAP WEED			



BO-RID K.O.

WEED KILLER E-4



LOW VOLATILE ESTER

2.4 D 4 LBS PER GALLON

CAUTION Keep Out of Reach of Children

SEE DIRECTIONS FOR ADDITIONAL CAUTIONS

KILLS WEEDS IN PASTURES, CROP and NON CROP AREAS, FENCE ROWS and RIGHT OF WAYS

ACTIVE INGREDIENTS
 2,4-D (2,4-Dichlorophenoxyacetic acid) 4.0%
 2,4-D (2,4-Dichlorophenoxyacetic acid) 1.1%
 INERT INGREDIENTS 94.9%

NET GALLONS
The R. H. Bogle Company
 ALEXANDRIA VA
 JACKSONVILLE FLA MEMPHIS TENN
 ALVA OKLA

BO-RID
 WEED
 KILLER
 E-4
 ACCEPTED
 U.S. DEPARTMENT OF AGRICULTURE
 OFFICE OF PEST CONTROL AND PLANT QUARANTINE
 WASHINGTON, D.C.



BO-RID **K.O.**

WEED KILLER E-4

LOW VOLATILE ESTER

2.4 D 4 LBS PER GALLON

CAUTION Keep Out Of Reach Of Children

SEE OTHER PANEL FOR ADDITIONAL DIRECTIONS

KILLS WEEDS IN PASTURES, CROP and NON CROP AREAS, FENCE ROWS and RIGHT-OF WAYS

ACTIVE INGREDIENTS

2,4-D, 2,4-Dichlorophenoxyacetic acid (ester form).....	4.00
Ethanolamine.....	0.30
Inert Ingredients.....	5.70
100%	10.00

INERT INGREDIENTS

NET GALLONS

The R. H. Bogle Company

ALEXANDRIA, VA
JACKSONVILLE, FLA
MEMPHIS, TENN
ALVA, OKLA

DIRECTIONS BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY WEED CONTROL IN CROPS

Broad Leaf Weeds in	Amount Bo Rid E-4 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
SPRING WHEAT	2.0	0.5	Pre-emergence	
RYE	2.0	0.5	Pre-emergence	
BARLEY	2.0	0.5	Pre-emergence	
OATS	2.0	0.5	Pre-emergence	
FALL WHEAT	2.0	0.5	Pre-emergence	
FIELD CORN PRE-EMERGENCE	2.0	0.5	Pre-emergence	
CORN POST-EMERGENCE	2.0	0.5	Post-emergence	
SORGHUMS POST-EMERGENCE	2.0	0.5	Post-emergence	
PASTURES AND RANGE	2.0	0.5	Pre-emergence	

PARTIAL LIST OF WEEDS CONTROLLED BY BO RID E-4

Annual Weeds	Annual Weeds	Annual Weeds
Canada Thistle	Chenopods	Cupressus
Common Ragwort	Clover	Fern
Crabgrass	Croton	Giant Hogweed
Dandelion	Fern	Hemp
Field Bindweed	Goldenrod	Johnsongrass
Flower Spurge	Guinea Grass	Marestail
Hairy Bitterweed	Hardhead	Meadow Spurge
Lambsquarters	Ironweed	Milkweed
Marestail	Japanese Knotweed	Nutsedge
Plantain	Large Crabgrass	Ragwort
Pigweed	Red Top	St. Johns Wort
Red Top	Rigwort	Succowood
Rigwort	Succowood	Wormwood
Succowood	Wormwood	
Wormwood		

MIXING INSTRUCTIONS

1. Mix one gallon of BO-RID E-4 with 100 gallons of water in the sprayer. 2. Add 2.0 lbs of 2,4-D acid to the mixture. 3. Spray the mixture over the weeds. 4. Repeat the application every 7-10 days. 5. Do not use in wind. 6. Do not use on bare soil. 7. Do not use on lawns. 8. Do not use on shrubs. 9. Do not use on trees. 10. Do not use on animals.