

CAUTION

The product is a concentrated liquid containing 24% 2,4-D. Avoid contact with skin, eyes, or clothing. Use caution during application of water or oil used for mixing. Measure to quantity. The spray method of application can cause damage or injury to animals, personal property, or crops.

DO NOT USE AROUND THE HOME OR RECREATION AREAS


For personal safety, should this liquid product come in contact with your eyes, skin, or clothing, wash immediately with plenty of water. If eye irritation continues, seek medical advice. If skin irritation occurs, wash with soap and water. If any damage occurs, do not treat with any other agent except rinsing or water. If you suspect damage to trees, shrubs, or lawns, consult a professional arborist or landscape architect before applying. If damage occurs, determine the cause and take appropriate action. If damage from this product is suspected, contact your distributor for advice. Every precaution must be exercised to prevent damage to trees, shrubs, or lawns. If damage occurs, do not apply further amounts of Bo-Rid. If any damage is suspected, contact your distributor for advice. If damage occurs, do not apply further amounts of Bo-Rid. If any damage is suspected, contact your distributor for advice.



BO-RID K.O.[®]

BRUSH KILLER BKE - 2+2

LOW VOLATILE ESTER

CAUTION: KEEP OUT OF REACH OF CHILDREN

2,4 D 2 LBS - 2,4,5 T 2 LBS PER GALLON
ESPECIALLY COMPOUNDED for USE
ON BRUSH and WOODY PLANTS
IN PASTURES, RIGHT OF WAYS and TIMBERLAND

ACTIVE INGREDIENTS

2,4-D, 2,4,5-T
2 LBS - 2 LBS
PER GALLON
ESPECIALLY COMPOUNDED for USE
ON BRUSH and WOODY PLANTS
IN PASTURES, RIGHT OF WAYS and TIMBERLAND

INERT INGREDIENTS

NET	GALLONS
100	

NOTE:

When using this product, wear protective clothing, including rubber gloves and a mask. Use caution when mixing or spraying. Do not let spray drift onto nearby trees or shrubs. If you suspect damage, do not apply further amounts of Bo-Rid. If any damage is suspected, contact your distributor for advice.

DO NOT USE AROUND THE HOME OR RECREATION AREAS

For personal safety, should this liquid product come in contact with your eyes, skin, or clothing, wash immediately with plenty of water. If eye irritation continues, seek medical advice. If skin irritation occurs, wash with soap and water. If any damage occurs, do not treat with any other agent except rinsing or water. If you suspect damage to trees, shrubs, or lawns, consult a professional arborist or landscape architect before applying. If damage occurs, determine the cause and take appropriate action. If damage from this product is suspected, contact your distributor for advice. Every precaution must be exercised to prevent damage to trees, shrubs, or lawns. If damage occurs, do not apply further amounts of Bo-Rid. If any damage is suspected, contact your distributor for advice.

DO NOT USE AROUND THE HOME OR RECREATION AREAS

For personal safety, should this liquid product come in contact with your eyes, skin, or clothing, wash immediately with plenty of water. If eye irritation continues, seek medical advice. If skin irritation occurs, wash with soap and water. If any damage occurs, do not treat with any other agent except rinsing or water. If you suspect damage to trees, shrubs, or lawns, consult a professional arborist or landscape architect before applying. If damage occurs, determine the cause and take appropriate action. If damage from this product is suspected, contact your distributor for advice. Every precaution must be exercised to prevent damage to trees, shrubs, or lawns. If damage occurs, do not apply further amounts of Bo-Rid. If any damage is suspected, contact your distributor for advice.

DO NOT USE AROUND THE HOME OR RECREATION AREAS

The R. H. Bogle Company

ALEXANDRIA, VA
4430 N. TURNPIKE
ALEXANDRIA, VA
22302-1102

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

MIXING INSTRUCTIONS

Add the recommended amount of BO-RID BKE 2+2 to one-half to one-quarter of a gallon of water or oil to be used as spray or mix with water and the remaining water or oil and mix until spray mixture is uniform. Continue mixing while spraying. Do not let water get into tank or oil sprays used in stump or basal treatment.

BO-RID BKE 2+2 is a mixture of 2,4-D and 2,4,5-T which can be mixed with water or dissolved in oil. BO-RID BKE 2+2 is designed for weeds and woody growth which are controlled better by a mixture of 2,4-D and 2,4,5-T than either agent separately. Bramble, blackberry, some hawthorns, osage orange, poison ivy, ground cherries, wild grapevines are typical of the vegetation which can be controlled along fence rows, right-of-ways, highways in pastures and on rangelands and beneath power lines.

**WOODY PLANTS AND BRUSH
IN PASTURE AND RANGELANDS AND NON-CROPLAND
SUCH AS FENCE ROWS, RIGHT OF WAYS**

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FRUIT TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

Type of Spraying	Amount of Bo-Rid BKE 2+2 To Use	When to Apply	How to Apply
FOLIAGE TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.
FIELD TREATMENT	1 QUART OR 100 GALLONS OF WATER OR OIL	After frost or after the first killing frost in late autumn or early winter.	By air, by ground, by back pack, or by hand sprayer.

AUTION
Handle with care. Avoid contact with skin or eyes. If contact occurs, wash thoroughly with water. If irritation persists, get medical attention. Do not eat, drink or smoke while using product. Do not spray near food crops or areas where food may be stored. Do not spray on plants used for food or fiber. Do not spray on areas where food may be stored. Do not spray on areas where food may be stored.

DO NOT USE AROUND THE HOME OR RECREATION AREA.



BO-RID K.O.


BRUSH KILLER BKE - 2

LOW VOLATILE ESTER

CAUTION: KEEP OUT OF REACH OF CHILDREN.

SEE OTHER CANE FOR ALL FINAL CAUTIONS

2.4 D 2 LBS., 2.45 T 2 LBS. PER GALLON
ESPECIALLY COMPOUNDED for USE
ON BRUSH and WOODY PLANTS
in PASTURES, RIGHT OF WAYS and TIMBERLAND



ACTIVE INGREDIENTS	NET	GALLONS
2,4-D	100.00%	100.0
2,45-T	100.00%	100.0
INERT INGREDIENTS	100.00%	100.0
	100.0%	100.0

NOTICE

Product No. 1-9-69

NET GALLONS

The R. H. Bogle Company

ALEXANDRIA, VA
KISONVILLE, FLA. ALVA, OKLA. MEMPHIS, TENN.




BO-RID K.O.

BRUSH KILLER BKE - 2+2

LOW VOLATILE ESTER

CAUTION: KEEP OUT OF REACH OF CHILDREN



24 D. 2 LBS. 24 G. T. LBS. PER GALLON
ESPECIALLY COMPOUNDED FOR USE
ON FRUIT & WOODY PLANTS
ASTERS, RIGID ORNAMENTAL FLOWERS,

ACTIVE INGREDIENTS:

- 1. 10% Zinc Oxide
- 2. 10% Calamine Lotion
- 3. 10% Camphor Oil
- 4. 10% Salicylic Acid

INERT INGREDIENTS:

STAINLESS STEEL
NET 10 GALLONS

The R. H. Bogle Company

ALEXANDRIA VA

The R. H. Boole Company

ALEXANDRIA VA

DIRECTIONS

**DO NOT USE READ DIRECTIONS, CAUTION AND
NOTICE STATEMENTS CAREFULLY.**

MIXING INSTRUCTIONS

RECORDED AND INDEXED
BY THE BUREAU OF INVESTIGATION,
U. S. DEPARTMENT OF JUSTICE,
AT THE REQUEST OF THE ATTORNEY GENERAL,
FOR THE USE OF THE ATTORNEY GENERAL,
AND FOR THE USE OF THE SECRETARY OF STATE,
IN THE PREPARATION OF THE TREATY
WITH GERMANY.

... FLOODY PLANT, AND BRUSH
... THE WILDERNESS, AND SUN BURNED LAND
... SUCH AS FENCE ROWS, RIVER BANKS,

Type	Amount of Ba/Bd Pkg. To Use	When To Apply	How To Apply
FOLIAGE TREAT- MENT	1/2 oz.	Early Spring, just before new growth	Stir into water, wet brush, then apply spray to leaves and flowers.
FRIEZE TREAT- MENT (For large trees)	1/2 oz.	Early Spring, just before new growth	Make up 1/2 P.D. water, place spray gun in spraying line, turn on treat. Spray the trunk and limbs.
STUMP TREAT- MENT	1/2 oz.	Early Spring, just before new growth	Get a bucket full of water, dip brush in water and paint the stump.
SAP- BARK TREAT- MENT	1/2 oz.	Early Spring, just before new growth	Get a bucket full of water, dip brush in water and paint the bark.

1 quart contains 1 lb. total 2,4-D and 2,4,5-T acids
1 lb. does not contain 2,4,5-T acid.

At 14 years, the upper teeth will be reduced to one-half of their original size, and the lower teeth to one-third of their original size. The upper teeth will be reduced to one-half of their original size, and the lower teeth to one-third of their original size. The upper teeth will be reduced to one-half of their original size, and the lower teeth to one-third of their original size.