

NOV 27 1961
ACCEPT

021-100

AGGREG

GENERAL DIRECTIONS

CIOVAP® Insecticide is a specially prepared ready to use insecticide suitable for use in conventional hand, power or automatic spray equipment and in backrubbers. CIOVAP is approved for lactating dairy animals. No feed off period is necessary when used on beef animals. Apply with hand power or automatic equipment to moisten the hair only and not in excess of 2 fluid ounces daily per animal. Do not wet the skin due to possible skin irritation. Do not apply regularly to calves under 6 months of age. Brahman cattle should not be treated as they may show hypersensitivity to organic phosphates.

For the control of certain insects attacking animals, apply in accordance with State and Federal Recommendations.

VITHIN
OW.

CK

USE AND DIRECTIONS FOR APPLICATION

Lactating Dairy and Beef Animals: Flies (Horn Fly, Stable Fly, Face Fly, House Fly)

For the control of horn flies, stable flies, face flies, and house flies, make a light application of CIOVAP daily as a fine mist to thoroughly cover all parts of the animal, including the legs.

Lactating Dairy and Beef Animals: Lice

Make 1-2 applications of 3-3½ ounces of CIOVAP at 14-day intervals as a fine mist spray to thoroughly cover all parts of the animal. If material is applied by fogging or misting equipment, the maximum spraying time per animal is 20 seconds.

Back and Facerubbers:

For the control of horn flies and face flies on lactating dairy and beef animals, use CIOVAP in backrubbers and facerubbers. Use 1 gallon for each 20 linear feet of burlap backrubber. For automatic backrubbers recharge the reservoir as needed. Animals must rub face on rubbers for best face control.

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Trichlorofon.

ACD-13G

PRECAUTIONS IN USING

Handle with care. Avoid contact with eyes, nose, mouth, and skin. If contact occurs, wash immediately with plenty of water. If eyes get irritated, get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not reuse container. Destroy when empty.

SOLID RED



NOV 27 1961
ACCEPT

SHELL CHEMICAL COMPANY
A Division of Shell Oil Company
AGRICULTURAL DIVISION
HEALTH & NUTRITION
SAN RAMON, CA. 94583

CIOVAP[®]

Insecticide

SOLID RED

DAIRY & BEEF CATTLE SPRAY

Contains CIODRIN[®] & VAPONA[®] Insecticides

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

Controls **HORN FLIES - STABLE FLIES - HOUSE FLIES -
FACE FLIES & LICE**


CAUTION

READY-TO-USE

USE TIME



NET CONTENTS: 1 GALLON



CIOVAP

DAIRY & BEEF CATTLE SPRAY

Contains CIODRIN & VAPONA Insecticides

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED.

**Controls HORN FLIES - STABLE FLIES - HOUSE FLIES -
FACE FLIES & LICE**

ACTIVE INGREDIENTS

Dimethyl Phosphate of Alpha Methylbenzyl
3 Hydroxy Cis Crotonate
2,2 Dichlorovinyl Dimethyl Phosphate
Related Compounds

BY WEIGHT

1.00%
0.23%
0.02%
98.75%

INERT INGREDIENTS

TOTAL 100.00%

CIODRIN[®] Insecticide
Equivalent to 0.25% VAPONA[®] Insecticide

(SEE USE AND DIRECTIONS FOR APPLICATION ON BACK PANEL)

CAUTION

KEEP OUT OF REACH OF CHILDREN

(See left panel for additional cautions)

EPA Reg. No. 201-152-AA

READY-TO-USE



NET CONTENTS: 1 GALLON

