

1% VAPONA® INSECTICIDE **READY-TO-USE** ANIMAL SPRAY SOLUTION



For the Control of Horn Flies, Stable Flies, Face Flies, Mosquitoes and House Files by Direct Application on Dairy and Beef Cattle.

ACTIVE INGREDIENTS												(BY WEIGHT			
2, 2-Dichlorovinyl Dime	et	hy	/	P	h	01	P	ha	rte) .	•					. 0.93%*
Related Compounds		•	•				• .									. 0.07%
Deodorized Kerosene .	•															49.50%
Mineral Seal Oil																
																100.00%

*Equivalent to 1% VAPONA® Insecticide EPA Reg. No. 201-164AA

WARNING! KEEP OUT OF THE REACH OF CHILDREN

PRECAUTIONS IN USING

HAZARDOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. In case of contact with skin or eyes, flush with plenty of water, for eyes get medical attention. Keep away from open flame. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed, water, or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

DECONTAMINATE AND DESTROY USED CONTAINERS: Do not reuse for any purpose. Decontaminate containers by rinsing thoroughly with water and detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate potable water supplies.

GENERAL DIRECTIONS

1.0% VAPONA Insecticide animal spray solution is a specially prepared ready-to-use insecticide for use in conventional hand, power or automatic spray equipment, electric, fogging or misting equipment.

For the control of certain insects attacking animals, apply in accordance with State and Federal recommendations. Do not apply regularly to calves under 6 reontes or a treated as they may show hypersensitivity to organic phosphates. A

Do not treat animals within one day of slaughter.

Do not apply dermally in combination with Trichlorofon.

FOR ECONOMIC POLLIN REGIST 9. ED UNDER NO USE AND DIRECTIONS FOR APPLICATION AIS

Cattle, Dairy and Beef - Flies (Horn Fly, Stable Fly, 20/-

Face Fly, House Fly) Mosquitoes Apply 1 to 2 fluid ounces of 1.0% solution per animal as a mist spray daily with hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply

ANIMAL BUILDINGS

in excess of 2 fluid ounces per day per animal. No feed-off period is necessary when used on beef animals.

Horse Stables, Dairy Barns, Auction & Show Barns (Flies, Gnats, Mosquitoes)

Apply 1.0% VAPONA Insecticide as a fog or mist. 1 pint of a 1% solution will treat 8000 cubic feet (40 x 20 x 10). In dairy barns, before milking, the fog or mist can be applied over and around each animal to kill flies. Before applying reduce air movement as much as possible by closing doors, windows and other openings. Do not use this solution for fogging or misting purposes in areas where animals have received a direct application within 8 hours. Do not contaminate water, feed or foodstuffs, milk or milking utensils.

NOTICE OF WARRANTY

SHELL CHEMICAL COMPANY MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

NET CONTENTS: 30 GALLONS

SHELL CHEMICAL COMPANY, A Division of Shell Oil Company, AGRICULTURAL DIVISION, SAN RAMON, CA. 94583