

NET CONTENTS ONE POUND

Tobacco States

75% ZINEB

WETTABLE POWDER

For Control Of Certain Diseases
On Tobacco, Tomatoes, Flowers
and Certain Other Plants

ACTIVE INGREDIENT:	% by Wt.
Zineb (zinc ethylene bisdithiocarbamate)	75.00%
INERT INGREDIENTS:	25.00%
TOTAL	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED — Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting if given anything by mouth to or on that person. Get medical attention immediately.

IF INHALED — Remove victim to fresh air and insure freely respiration. If indicated.

IF ON SKIN — Wash with soap and water. Rinse thoroughly.

IF IN EYES — Rinse for 15 minutes with water and call a physician immediately.

SEE BACK PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 226-21

EPA Est. 226-KY-1

ACCEPTED

206-21
APR 13
Under Federal Pesticide and Fungicide Act as amended, for its registration under EPA Reg. No.

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. During the mixing and loading of this product, please wear the following protective clothing: long pants, long sleeve shirt, gloves, hat and boots. Do not enter treated area without protective clothing until dusts have settled. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. In case of contact, flush with plenty of water. For eyes, get medical attention. Keep away from fire and sparks.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

STORAGE and DISPOSAL

STORAGE: Do not store in areas of extreme heat or cold. Protect from moisture or extreme humidity.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag or can into application equipment. Then dispose of empty bag or can in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this Product in a manner inconsistent with its labeling.

GENERAL DIRECTIONS — Use 1½ to 2 pounds per 100 gallons of water (unless otherwise specified) in conventional sprayers, or equivalent in concentrate or aerial sprayers. Three pounds should be applied per acre on average size field crops.

Apply at first sign of, or prior to appearance of disease. Repeat at one or two week intervals as necessary unless otherwise noted.

BEANS — Downy Mildew, Rust, Anthracnose: Begin when first blossoms appear. Repeat applications at 4 to 7 day intervals. Do not apply within 7 days of harvest.

CABBAGE, BROCCOLI, CAULIFLOWER, COLLARDS — Downy mildew, Alternaria leaf blight: Do not apply within 7 days of harvest.

CANTALOUPE, CUCUMBERS AND WATERMELONS — Downy mildew, Alternaria (Macrosporium) blight, Anthracnose, Gummy stem blight: Direct spray especially at under-side of leaves, begin applications when plants appear above the ground.

TOMATOES — Early and late blight, Anthracnose, Stemphylium and Septoria leaf spots, Cladosporium gray mold: Begin spraying when disease is reported in area and repeat applications at 7 to 10-day intervals or more often if blight conditions are severe.

STRAWBERRIES — Leaf spot: Begin when new growth starts in spring. Repeat until fruit begins to form. Leaf scorch: Apply in new beds in late summer — 2 to 3 weekly applications. Do not apply within 7 days of harvest.

TOBACCO — Blue mold: Begin applications when plants emerge or when blue mold is first reported in area. Spray regularly 2 or 3 times per week during active blue mold weather. Repeat applications immediately after a rain. Use 3 lbs. 75% Zineb per 100 gal. water (2 level tablespoons per gallon). Apply at 3 to 6 gal. per 100 sq. yds. of plant bed, depending on size of plants.

SNAPDRAGONS, CHRYSANTHEMUMS, CARNATIONS, HOLLYHOCKS — Rust, Leaf spot and Anthracnose: Apply at weekly to 10-day intervals as soon as disease conditions threaten. For control of Botrytis petal spot and Ascochyta ray make application twice weekly during the blooming season.

ROSES — Black spot: Apply at weekly intervals. Applications should begin when new leaves unfold in spring.

WARRANTY STATEMENT

Tobacco States MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE OR OTHERWISE, EXPRESS OR IMPLIED concerning this product or its use, which extend beyond the statements on this label.

TOBACCO STATES CHEMICAL CO., INC.
LEWINGTON, KENTUCKY

M-2-10-84