

Net Contents

5 Pounds

7½ % METHOXY

GARDEN DUST

for the control of


MEXICAN BEAN BEETLES, LEAFHOPPERS, CUCUMBER BEETLES

*Methoxychlor, Technical 7½%
INERT INGREDIENTS 92½%

*7½% Methoxychlor, technical is equivalent to 6.6% 2,2-bis
(p-methoxyphenyl)-1, 1, 1-trichloroethane and .9% related compounds.

CAUTION:

Keep out of the reach of children. See other cautions on rear panel USDA Reg. No. 226-45



DIRECTIONS — do not dilute.

Use 7½ METHOXY GARDEN DUST throughout the garden for control of certain chewing insects. When dusting this insecticide it is very important to thoroughly cover the undersides as well as the tops of the leaves, for the average dusting use about 20 pounds of dust per acre. Repeat when necessary. Do not use on food crops later than 7 days of harvest: except on lettuce and broccoli do not use within 14 days of harvest: except on turnips do not use within 7 days of harvest: (14 days if tops are to be used as food or feed). This product should not be applied to beets within 14 days of harvest if the tops are to be used as food or feed.

CROPS	INSECTS
BEANS	Mexican Bean Beetle, Corn Earworm, Leafhoppers
TOMATOES	Flea Beetles, Leashoppers, Fruit Worm, Fall Armyworms, Suck Fly, Blister Beetle
CUCURBITS Cucumbers Melons Squash	Cucumber Beetles (various), Leathop- pers, Flea Beetles, Squash Borer
ASPARAGUS	Asparagus Beetle
BEETS	Flea Beetles, Leathoppers
EGGPLANT	Blister Beetles, Flea Beetles, Colorado Potato Beetles
BROCCOLL CABBAGE and CAULIFLOWER	Cross-striped Cabbage Worm, Dia- mond back Moth, Imported Cabbage Worm, Flea Beetle
LETTUCE	Leathoppers
PEAS	Pea Weevil
POTATOES	Leashopper, Flea Beetles, Colorado Potato Beetle
TURNIPS	Flea Beetles

CAUTION: This product is poisonous to fish. Keep out of lakes, streams and ponds.

S-1-68

ites Chemical Co., Inc., Lexington, Kentucky