

NET WEIGHT ONE POUND

**Tobacco States**



# Blight & Insect DUST

GENERAL PURPOSE GARDEN DUST for the control of certain GARDEN INSECTS AND DISEASES that prey on the home garden.

**ACTIVE INGREDIENTS:**

- 1. Deltamethrin 0.05%
- 2. Methoxychlor 4.0%
- 3. Cyfluthrin 0.05%
- 4. Imidacloprid 0.05%
- 5. Azoxystrobin 0.05%
- 6. Chlorantraniliprole 0.05%
- 7. Spinosad 0.05%
- 8. Lambda-cyhalothrin 0.05%
- 9. Fenprophate 0.05%
- 10. Permethrin 0.05%
- 11. Bacillus thuringiensis 0.05%
- 12. Azadirachtin 0.05%

**CAUTION:** Keep Out of Reach of Children  
See other Cautions on back panel

ACCEPTED  
Jul 8 1968  
226-65

**CAUTION:** Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals. Do not apply to fruit or vegetables to be used for food later than 15 days before harvest. Keep bag tightly closed and store in dry place. Misapplication upon repeated, prolonged, or careless use, may cause foliar damage inhibition. Should poisoning occur obtain prompt medical aid. Atropine is antidotal.

Toxic to fish. Use with care.  
**DIRECTIONS DO NOT DILUTE**

When dusting with this material it is important to thoroughly cover the underside of the leaves as well as the tops and stems. Most applications will start as soon as the plants appear and repeat every 5 to 10 days, longer intervals in wet weather. Repeat applications are not needed after washing rains. All residues should be removed from the vegetables by washing, brushing or other suitable means before using or marketing. On strawberries start dusting when insects appear and continue at weekly intervals for approximately 4 dustings. Use 20 to 40 lbs. per acre. Maximum use on beans should be 30 lbs. per acre.

| CROPS         | INSECTS  | DISEASES  |
|---------------|--|---|
| TOMATOES      | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite | Early and Late Blights, Anthracnose, and Septoria Leaf Spot |
| POTATOES      | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite | Early and Late Blights                                      |
| BEANS         | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite | Early Blight, Mosaic, and Rust                              |
| CHRYSAANTHEMS | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite |   |
| ROSES         | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite |   |
| CARNATIONS    | Colorado Beetle, Flea Beetle, Aphid, Armyworm, Cutworm, Leafminer, Spider Mite |   |