

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn bee-keepers well in advance to locate hives at a safe distance until one week after application. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

**DIRECTIONS**

Apply 1 3/4% BEAN DUST when you first see signs of insect infestation, preferably before damage from feeding occurs. For best results apply thoroughly to upper and lower leaf surfaces at the rate of 30 pounds per acre. If needed, repeat applications every 7-10 days. Apply up to and including the day of harvest. Do not feed treated bean vines to dairy or meat animals.

**CROPS**

Beans  
potatoes

**INSECTS**

Mexican Bean Beetle  
Colorado Potato Beetle

**CAUTION**

HARMFUL IF SWALLOWED OR INHALED. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of food and food stuffs.

USDA Reg. No. 226-143

ACCEPTED  
Aug 9, 1971  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
ED UNDER NO. 226-143 SUBJECT  
TO ATTACHED COMMENTS

226-143

**Tobacco States  
Pesticides**

NET CONTENTS—2 POUNDS



**1 3/4 % BEAN DUST**

CONTAINS SEVIN\*



For the control of  
MEXICAN BEAN BEETLES &  
COLORADO POTATO BEETLES

Active Ingredient:  
Carbaryl (1-Naphthyl N-Methylcarbamate) ..... 1.75%  
Inert Ingredients: ..... 98.25%

\*Trade Mark of UCC for 1-Naphthyl N-Methylcarbamate



**CAUTION:** Keep out of reach of children.  
See other cautions on back panel.

TOBACCO STATES CHEMICAL CO., INC.  
LEXINGTON KENTUCKY

