

NET CONTENTS ONE QUART

Tobacco States

THIODAN 2

EMULSIFIABLE CONCENTRATE

WARNING: Keep out of reach of children. See Antidote statement and other required warning statements on the back panel.

ACTIVE INGREDIENTS:

*Hexachlorohexahydromethano 2,4,3-benzodioxathiepin oxide	24.00%
Aromatic Petroleum Solvent	69.00%

INERT INGREDIENTS:	7.00%
---------------------------------	-------

*Thiodan 2 Emulsifiable Concentrate contains 2 pounds of this material per gallon. (R) Thiodan is a registered trademark of Farbwerke Hoechst A.G., U.S. Patent No. 2,799,685

USDA Reg. No. 226-159

WARNING:
Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, immediately remove contaminated clothing and flush attention. Wash thoroughly with soap and water after handling and commercial or prolonged exposure in spray-mixing and loading operation. Use a respirator of a type passed by the U.S. Department of Agriculture for protection.
This product is poisonous to fish and wildlife. Keep out of lakes, streams, or areas where drift of spray from areas treated, or areas occupied by unprotected animals.
Apply this product only as specified on this label.

DIRECTIONS FOR USE:

Complete coverage is necessary for a good control. Use at least 30 gallons per acre.

CROP	TO CONTROL	RATE OF APPLICATION
Beans	Mexican bean beetle	1 quart per acre (2 lbs. per 1000 sq. ft. water for)
Platoes	Flea beetle	EASTERN
	Colorado potato beetle	1 quart per acre
	Leafhopper	(2 lbs. per 1000 sq. ft. water for)
	Aphids	
	Southern Armyworm	1 to 2 qt.
	Green Stinkbug	(3 lbs. per 1000 sq. ft. water for)
	Colorado potato beetle	
	and other plant bugs	
	Potato tubeworm	
	Whitefly	
	Flea beetle	SEED BE
	Hornworm	1 qt. 100
	Budworm	FIELD
	Green peach aphid	1 to 2 qts
	Cabbage looper	
	Whitefly & Aphids	1 quart per
	Colorado potato beetle	
	Flea beetle	
	Green Stinkbug	
	Blister beetle	
	Hornworm	

ACCEPTED
MAY 11, 1971

FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 226-159 SUBJECT
TO ATTACHED COMMENTS.


CAUTION: Workers entering treated areas the day of application should wear protective clothing.
WARNING FLAMMABLE: Keep away from heat and open flame. Keep in a safe place. Do not store at temperatures below 0 F. Do not use in areas where fire is a hazard.
Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies.

S 2-68

PESTICIDE

TOBACCO STATES CHEMICAL

LEXINGTON, KENTUCKY



DAN 2

BLE CONCENTRATE

Keep out of reach of children. Statement and other required elements on the back panel.

S:

Hydromethano 2,4,3-	24.00%
epin oxide	69.00%
um Solvent	7.00%

le Concentrate contains 2 pounds of n. (R) Thiodan is a registered trademark of Tacco States Chemical Co., U.S. Patent No. 2,799,685

Reg. No. 226-159

STICIDE

ACCEPTED

MAY 11, 1971

FEDERAL INSECTICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 226-159 SUBJECT TO ATTACHED COMMENTS.

WARNING

Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Hazardous if swallowed, inhaled, or absorbed through skin! In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U.S. Department of Agriculture for Thiodan protection.

This product is poisonous to fish and wildlife. Keep out of lakes, streams, and ponds. Do not apply when weather conditions favor drift of spray from areas treated, or areas occupied by unprotected humans or beneficial animals.

Apply this product only as specified on this label.

DIRECTIONS FOR USE

Complete coverage is necessary for a good control. Use at least 30 gallons of water per acre.

CROP	TO CONTROL	RATE OF APPLICATION	REMARKS
	Mexican bean beetle	1 quart to acre (2 lbs. per gallon of water for small plots)	Do not apply after pods begin to form. Do not feed threshings to livestock. Do not use on lima beans.
	Flea beetle Colorado potato beetle Leafhopper Aphids	EASTERN AREAS 1 quart to acre (2 lbs. per gallon of water for small plots)	Apply as needed up to harvest.
	Southern Armyworm Green Stinkbug Leaf-footed plant bugs and other plant bugs Potato tubeworm Whitefly	1 to 2 qt. per acre (3 lbs. per gallon of water for small plots)	
	Flea beetle Hornworm Budworm Green peach aphid Cabbage looper	SEED BED 1 qt. 100 gallon FIELD 1 to 2 qts. per acre	In seed beds apply about 6 gallons of spray per 100 square yards. Do not apply within 5 days of harvest. Use minimum of 30 gal. of water per acre.
	Whitefly & Aphids Colorado potato beetle Flea beetle Green Stinkbug Blister beetle Hornworm	1 quart per acre	When applying in Greenhouses, wear mask or respirator of type passed by U.S.D.A. for Thiodan protection. Do not apply within 1 day of picking.

CAUTION: Workers entering treated areas the day of application should wear protective clothing.

WARNING FLAMMABLE: Keep away from heat and open flame. Keep container closed. Leaking packages should be removed to a safe place. Do not store at temperatures below 0°F. Do not use in undiluted form.

Rinse equipment and containers and dispose of wastes and spillage by burying. Destroy empty containers. Perforate, crush and bury in non-crop lands away from water supplies.

S 2-68

TACCO STATES CHEMICAL CO., INC.

LEXINGTON, KENTUCKY