

NET WEIGHT ONE POUND

Tobacco States

# 50% Chlordane WETTABLE POWDER

**ACTIVE INGREDIENT—**

Technical Chlordane\* ..... 50%

**INERT INGREDIENT** ..... 50%

\* Equivalent to 30% octachloro-4, 7-methanotetrahydroindane and 20% related compounds.

**CAUTION: Keep Out of Reach of Children**

**CAUTION:** Avoid contamination of feed and foodstuffs. Avoid inhalation of dusts or sprays. Harmful if swallowed. Contact with skin causes toxic symptoms. In case of skin contact, wash with soapy water.

Do not use on dairy animals, in dairy barns, or feed or forage to be fed to dairy animals or other animals being finished for slaughter. Do not use this product in poultry houses.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Destroy empty container by burning or burying in a safe place. (Keep out of smoke).

EPA Reg. No. 226-219

EPA Est. 226-KY-1

**ACCEPTED**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 226-219

**DIRECTIONS FOR SOIL INSECTS**

— Control Limited To Preplant Treatment Only —

**STRAWBERRIES** — White Grubs, Japanese Beetle Larvae, Mole Crickets and Cutworms. Mix 10 to 20 lbs. of this material with sufficient water to cover one acre, spraying until the soil surface is wet.

**CORN** — Cutworms. Mix 10 to 20 lbs. of this material with sufficient water to cover one acre, spraying until the surface is wet. Use suspended effective August 1, 1976.

For more effective control, if practical, disc 4 to 6 inches before seeding or transplanting corn or strawberries.

50% Chlordane can be mixed with fertilizer and applied same ways as above.

Do not treat when soil is wet. Apply before rain or sprinkling.

5176

**PESTICIDE**

**TOBACCO STATES CHEMICAL CO., INC**