

NET CONTENTS ONE GALLON

Tobacco States
BRAND

2-M ALFALFA SPRAY

Contains 2 lbs. Malathion and 2 lbs. Methoxychlor per gallon

ACTIVE INGREDIENTS:

*Malathion	22.73%
**Methoxychlor	22.73%
Xylene	48.34%

INERT INGREDIENTS	6.20%
TOTAL	100.00%

* O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

** Equivalent to 20.00% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 2.73% other isomers and reaction products.

CAUTION: Keep out of reach of children.

Do not use, pour, spill, or store near heat or open flame.

HARMFUL IF SWALLOWED. AVOID BREATHING OF SPRAY MIST. AVOID CONTACT WITH SKIN

Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. If swallowed, induce vomiting and get medical attention. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

GENERAL DIRECTIONS

2-M ALFALFA SPRAY may be used in conventional hydraulic sprayers, low-volume ground applicators, and airplane sprayers (use not less than 2 gallons diluted spray per acre with aircraft applications). Mix this product with water by agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

APPLICATION DIRECTIONS

ALFALFA -- To control Alfalfa weevil larvae apply 5 pints (in approximately 25 gallons of water) per acre. Make application in spring when most of the tips are beginning to show feeding. A second application may be necessary. If harvest infestation is severe, treat stubble as soon as crop is removed. To control Clover leaf weevil apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring when alfalfa is 2 to 6 inches tall.

To control Alfalfa caterpillar apply up to 2 quarts (in approximately 25 gallons of water) per acre. Apply if there are 10 or more caterpillars per sweep of net, usually when alfalfa is 1/3 grown.

To control meadow spittlebug apply 2 quarts (in approximately 25 gallons of water) per acre. Apply within a week after eggs begin to hatch.

To control potato leafhopper apply 1 quart (in approximately 25 gallons of water) per acre. Apply when alfalfa is about half grown or earlier if insects become abundant. Do not apply within 7 days of harvest. Do not apply during full bloom.

CLOVER -- To control clover leaf weevil apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring when clover is 2 to 6 inches tall. To control meadow spittlebug apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring within a week after eggs begin to hatch. As a fall treatment, apply 3 quarts (in approximately 25 gallons of water) per acre. Make application in early September.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agriculture Extension Service.

Treat clover in bloom only when bees and other pollinators are not visiting plants.

Do not apply to alfalfa within 7 days of harvest. Do not apply to clover within 7 days of harvest.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 226-229

PESTICIDE

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY