

NET CONTENTS: ONE GALLON



3-WAY DISINFECTANT

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

Recommended for cleaning, disinfecting, and sanitizing in hospitals, institutions, food plants, and commercial buildings.

ACTIVE INGREDIENTS:

Alpha-(p-Nonylphenyl)-omega-hydroxypoly(oxyethylene)-iodine complex (Providing 1.75% titratable iodine)	18.05%
Phosphoric Acid	18.00%

INERT INGREDIENTS	65.95%
TOTAL	100.00%

Manufactured under the following U. S. Patents:

2,931,777
2,977,315
3,028,299

3-WAY DISINFECTANT has its own built-in indicator of germicidal activity. When the amber color disappears, a fresh solution should be prepared. This product is not adversely affected by water hardness or low temperatures.

DANGER: KEEP OUT OF REACH OF CHILDREN

See back panel for additional precautions and First Aid

EPA Reg. No. 226-235

DIRECTIONS FOR USE:

Gross filth must be removed prior to application of the following recommended solutions.

FOR GENERAL CLEANING & DISINFECTING IN ONE OPERATION:

Use 3 ounces of 3-Way Disinfectant to 5 gallons of water. For porous or extremely dirty use 6 ounces to 5 gallons of water. For sanitizing previously-cleaned surfaces, use 3-Way Disinfectant in 5 gallons of water.

HOSPITALS:

Use as recommended above in animal rooms and cages, cafeterias, morgues, laboratory areas, patient rooms, bedframes, walls and floors, thermometers, instruments, rubber tubing & catheters. Allow at least 10 minutes contact time. Rinse all food-handling equipment with potable water before reuse. Before disinfecting, instruments should be cleaned to remove adhering blood and serous exudates.

COMMERCIAL BUILDINGS, INSTITUTIONS, AND GENERAL HOUSEKEEPING:

Use 3 ounces of 3-Way Disinfectant to 5 gallons of water for cleaning and disinfecting walls, lavatories, shower rooms, refuse containers, kitchen areas, and lunch room food-handling equipment with potable water before reuse.

FOOD PLANTS:

To sanitize previously-cleaned food-processing equipment and utensils, use 1 ounce of 3-Way Disinfectant to 5 gallons of water. Allow to drain but do not rinse. When used in food plants, 3-Way Disinfectant does not require a potable water rinse, in accordance with the Additive Regulation 121.2547.

DANGER: Do not get in eyes. Avoid contact with skin. Causes eye damage or skin irritation. In case of contact, flush immediately with plenty of water. For eyes, seek medical attention. Avoid contamination of food.

May be harmful if swallowed. If swallowed, do not induce vomiting. Drink milk, or plenty of water. Immediately get medical attention.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY

NOV 17 1973
UNLESS OTHERWISE SPECIFIED
FUNGICIDE
FOR INDUSTRIAL USE
REGISTERED UNDER EPA REG. NO. 226-235
TO ATTACH TO CONTAINER

226-235

ONE GALLON



3-WAY DISINFECTANT

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

for cleaning, disinfecting, and sanitizing in hospitals, food plants, and commercial buildings.

INGREDIENTS:

Trichlorophenyl-omega-hydroxypoly(oxyethylene)-	
iodine complex (Providing 1.75% titratable iodine)	18.05%
Hydrochloric Acid	16.00%
Surfactants	65.95%
.....	100.00%

Manufactured under the following U. S. Patents:

- 2,931,777
- 2,977,315
- 3,028,299

Disinfectant has its own built-in indicator of germicidal activity. As the amber color disappears, a fresh solution should be prepared. This product is not adversely affected by water hardness or high temperatures.

WARNING: KEEP OUT OF REACH OF CHILDREN

Check panel for additional precautions and First Aid

EPA Reg. No. 226-235

DIRECTIONS FOR USE:

Gross filth must be removed prior to application of the following recommended disinfecting solutions.

FOR GENERAL CLEANING & DISINFECTING IN ONE OPERATION:

Use 3 ounces of 3-Way Disinfectant to 5 gallons of water. For porous or extremely dirty surfaces use 6 ounces to 5 gallons of water. For sanitizing previously-cleaned surfaces, use 1 ounce of 3-Way Disinfectant in 5 gallons of water.

HOSPITALS:

Use as recommended above in animal rooms and cages, cafeterias, morgues, laboratories, operating areas, patient rooms, bedframes, walls and floors, thermometers, instruments, rubber gloves, tubing & catheters. Allow at least 10 minutes contact time. Rinse all food-handling surfaces with potable water before reuse. Before disinfecting, instruments should be cleaned to remove all adhering blood and serous exudates.

COMMERCIAL BUILDINGS, INSTITUTIONS, AND GENERAL HOUSEKEEPING:

Use 3 ounces of 3-Way Disinfectant to 5 gallons of water for cleaning and disinfecting floors, walls, lavatories, shower rooms, refuse containers, kitchen areas, and lunch rooms. Rinse all food-handling equipment with potable water before reuse.

FOOD PLANTS:

To sanitize previously-cleaned food-processing equipment and utensils, use 1 ounce of 3-Way Disinfectant to 5 gallons of water. Allow to drain but do not rinse. When used in this manner 3-Way Disinfectant does not require a potable water rinse, in accordance with the Federal Food Additive Regulation 121.2547.

DANGER: Do not get in eyes. Avoid contact with skin. Causes eye damage or skin irritation. In case of contact, flush immediately with plenty of water. For eyes, get medical attention. Avoid contamination of food.

May be harmful if swallowed. If swallowed, do not induce vomiting. Drink milk, egg white, or plenty of water. Immediately get medical attention.

RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD IT.

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY

NOV 02 1973
DEPARTMENT OF AGRICULTURE
OFFICE OF THE ASSISTANT SECRETARY FOR
ECONOMIC AFFAIRS
UNDER HO-226-235 SUBJECT
ATTACHED COMMENTS.

226-235