



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

228-333

Date of Issuance:

JUN 30 2009

NOTICE OF PESTICIDE:

Registration
[X] Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

Riverdale MCDA L.A.C.
Selective Herbicide

Name and Address of Registrant (include ZIP Code):

Nufarm Americas, Inc.
150 Harvester Drive, Suite 200
Burr Ridge, Illinois 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

- 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following revisions to the final printed labeling:
a. Per the acute toxicity review, the footnote "Note: Spanish language is optional" must be removed. In addition, as this product is listed as Category I for primary eye, the following text must also be added to the label in close proximity to the signal word:

"Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Continued on Page 2

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

JUN 30 2009

- b. Per the acute toxicity review, the Agency recommends that the Note to Physician be revised to address the Toxicity Category I eye irritation concerns. The following are examples the types of text that are appropriate:

- technical information on symptomatology;
- use of supportive treatments to maintain life functions;
- medicine that will counteract the specific physiological effects of the pesticide;
- company telephone number to specific medical personnel who can provide specialized medical advice

- c. Per the acute toxicity review and the Dicamba RED, the handler PPE must be revised to read:

“Some materials that are chemical-resistant to this product are **made of any waterproof material**. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

Goggles or face shield,
Long-sleeved shirt and long pants,
Shoes plus socks, and
Chemical-resistant gloves.”

- d. The NON-AGRICULTURAL USE REQUIREMENTS box and the text “The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to product agricultural plants on farms, forests, nurseries, or greenhouses” are not needed and may be deleted from the label.

The text “**Do not enter or allow others to enter the treated area until sprays have dried**” must remain on the label.

- e. Add “**Broadcast Treatment**” before the application restriction “Do not apply more than 1.5 lbs ae MCPA per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.”

Additionally, add the following directly after the Broadcast treatment restrictions:

“**Spot Treatment (For Right-of-Ways):**
Do not apply more than 3.0 lbs MCPA ae per acre per year.
A spot treatment is defined as an area no greater than 1000 ft. sq.”

- f. The following spray drift text as required on Page 91 of the MCPA RED must be added to the label and any conflicting text must be deleted:

“SPRAY DRIFT MANAGEMENT”

-Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

-Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.”

-Apply only when the wind speed is 2-10 mph at the application site.

-Do not make applications into temperature inversions.

Additional requirements for aerial applications:

-The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

-Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

-When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

-Do not apply with a nozzle height greater than 4 feet above the crop canopy.

- g. Revise the heading “Ornamental Lawns and Turf” to “Non-Residential Ornamental Lawns and Turf”

- h. The following restrictions are required by the source technical Clopyralid:

Non-Agricultural Uses

Turfgrass and lawn uses are restricted to non-residential sites.

Note: In the states of California, Oregon and Washington, turfgrass and lawn uses are restricted to golf courses only.

This product **must not** be formulated into manufacturing-use herbicide products or end-use herbicide products for use on residential turfgrass.

- i. If the product is labeled for reformulation into products bearing labeling for turfgrass applications other than residential turfgrass, the label must contain the following statements:

General Precautions:

- Do not use on residential turfgrass.
- Do not send grass clippings to a compost facility.
- Do not collect grass clippings for mulch or compost.
- Applicator must give notice to landowners/property managers to not use grass clippings for composting.
- In the states of California, Oregon and Washington, turfgrass and lawn uses are restricted to golf courses only.

- 3. A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

Sample 5-25-09

5/11

ACCEPTED
with COMMENTS
In EPA Letter Dated:

JUN 30 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

MCDA L.A.C.

Selective Herbicide

228-333

FOR SELECTIVE BROADLEAF WEED CONTROL IN NON-RESIDENTIAL
ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, CLOPYRALID AND DICAMBA

CONTROLS: Dandelion, Chickweed, Black medic, Knotweed, Plantain, Oxalis, Clover, Cocklebur, Thistle and
many other species of broadleaf weeds; some of which are listed on this label.

ACTIVE INGREDIENT:

Dimethylamine Sal. of 2-Methyl-4-Chlorophenoxyacetic Acid*	48.13%
Monmethanolamine Salt of 3, 6-Dichloro-2-Pyridinecarboxylic Acid**	2.58%
Dimethylamine Salt of Dicamba (3, 6-Dichloro-o-Anisic Acid)***	4.73%

OTHER INGREDIENTS:	44.56%
TOTAL	100.0%

By Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	39.30%, 3.75 lbs./gal.
**3, 6-Dichloro-2-Pyridinecarboxylic Acid	1.96%, 0.18 lbs./gal.
***3, 6-Dichloro-o-Anisic Acid	3.93%, 0.37 lbs./gal.

FOR USE BY PROFESSIONAL TURF MAINTNEANCE PERSONNEL,
LANDSCAPING OR COMMERCIAL APPLICATORS ONLY.

**KEEP OUT OF REACH OF CHILDREN
DANGER - PELIGRO**

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

NOTE: Spanish language is optional.

EPA REG. NO. 228-333
EPA EST. NO. 228-IL-1

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527



1/6
=

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER - PELIGRO**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. Do not get in eyes, or on clothing. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on the EPA chemical resistant category selection chart.

Mixers, loaders, applicators and other handlers must wear: long sleeved shirt and long pants, shoes plus socks and chemical resistant gloves when mixing, loading or using any hand-held equipment.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washable exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS	
Users should:	
<ul style="list-style-type: none"> • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

MCPA has properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. This product is for use only on Non-Residential Ornamental Turf (Athletic and Recreational Sports Fields, Cemeteries, Golf Courses**, Industrial Sites, Non-Cropland, Parks, Rights-of-Way and Roadsides). **Excluding Bentgrass Tees.

Do not use on residential turfgrass. Do not collect grass clippings for mulch or compost. Do not send clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for mulching or composting.

Not for use on turf being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes.

NOT FOR SALE OR USE IN NASSAU AND SUFFOLK COUNTIES IN N.Y. STATE.

IN CALIFORNIA AND WASHINGTON, TURFGRASS AND LAWN USES ARE RESTRICTED TO GOLF COURSES ONLY.

IN OREGON, THE USE OF THIS PRODUCT IS LIMITED TO THE SITES STATED ON THIS LABEL WHICH ARE AGRICULTURAL, FOREST, RIGHT-OF-WAY, GOLF COURSE OR CEMETERY SITES.

*This product cannot be used on agricultural or forest sites.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

USE PRECAUTIONS

Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on Centipede, St. Augustine, Dichondra, nor on lawns or turf where desirable clovers are present. Avoid fine mists. Use lawn type sprayer with coarse spray as wind drift is less likely. Avoid contact with exposed feeder roots of ornamentals and trees. Maximum control of weeds will be obtained from Spring or early Fall applications when weeds are actively growing. The degree of weed control and duration of effect will vary with weed size and density, spray rate and coverage, and growing conditions before, during, and after the time of treatment. Use the higher rate for hard-to-control weeds. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F.

The suitable use of this product on non-recommended turf species may be determined by treating a small area at any rate/acre which does not exceed 3 pints/acre. The treated area should be observed for any sign of turf injury for a period of 30 days of normal growing conditions to determine the phytotoxicity and efficacy to the treated area.

For optimum results: (1) avoid applying during excessively dry or hot periods unless irrigation is used; (2) turf should not be mowed 1 to 2 days before and following application; (3) reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water, and rinse thoroughly before reuse for other sprays.

GENERAL APPLICATION RESTRICTIONS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

WEEDS CONTROLLED

Alder	Bur ragweed	English daisy	Hop clover
Annual yellow sweet clover	Burweed	Fall dandelion	Horsenettle
Artichoke	Buttercup	False dandelion	Horsetail
Aster	Canada thistle	False flax	Indiana mallow
Austrian fieldcress	Carolina geranium	False sunflower	Ironweed
Bedstraw	Carpetweed	Fiddleneck	Jewelweed
Beggartick	Catchweed bedstraw	Field bindweed	Jimsonweed
Biden	Catsear	Field pansy	Kochia
Bindweed	Catnip	Flea bane (daisy)	Knawel
Bird vetch	Chickweed	Flixweed	Knotweed
Bitterweed	Chicory	Florida betony	Lambsquarter
Bitter wintercress	Cinquefoil	Florida pusley	Lespedeza
Black-eyed Susan	Clover	Frenchweed	Locoweed
Black medic	Cockle	Galinsoga	Lupine
Black mustard	Cocklebur	Garlic mustard	Mallow
Blackseed plantain	Coffeebean	Goathead	Marshelder
Blessed thistle	Coffeeweed	Goatsbeard	Matchweed

Blue lettuce
Blue vervain
Box elder
Bracted plantain
Brassbuttons
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Duckfoot plantain
Bulbous buttercup
Bull nettle
Bull thistle
Burdock
Earning nettle

Common chickweed
Common mullein
Common sowthistle
Corn Chamomile
Creeping jenny
Crimson clover
Croton
Cudweed
Curly dock
Curly indigo
Dandelion
Dead nettle
Dock
Dogbane
Dogfennel
Elderberry

Goldenrod
Ground ivy
Gumweed
Hairy bittercress
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Honeysuckle

Mexicanweed
Milk vetch
Milkweed bloodflower
Mugwort
Morningglory
Mouseear chickweed
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oxalis
Oxeye daisy
Parsley-piert

WEEDS CONTROLLED (CONTINUED)

Parsnip	Rough cinquefoil	Stinkweed	White clover
Pearlwort	Rough fleabane	Stitchwort	White mustard
Pennycress	Roundleaved marigold	Strawberry clover	Wild mustard
Pennywort	Rush	Sumac	Wild aster
Peppergrass	Russian pigweed	Sunflower	Wild buckwheat
Pepperweed	Russian thistle	Sweet clover	Wild carrot
Pigweed	St. Johnswort	Tall nettle	Wild four-o'clock
Pineywoods bedstraw	Scarlet pimpernel	Tall vervain	Wild garlic
Plains coreopsis	Scotch thistle	Tansy mustard	Wild geranium
Plantain	Sheep sorrel	Tansy ragwort	Wild lettuce
Poison hemlock	Shepherdspurge	Tanweed	Wild marigold
Poison ivy	Slender plantain	Tarweed	Wild onion
Poison oak	Small flower galinsoga	Thistle	Wild parsnip
Pokeweed	Smartweed	Tick trefoil	Wild radish
Porycle	Smooth dock	Toadflax	Wild rape
Porycleweed	Smooth pigweed	Trailing Crownvetch	Wild strawberry
Prickly lettuce	Sneezeweed	Tumble mustard	Wild sweet potato
Prickly sida	Southern wild rose	Tumble pigweed	Wild vetch
Primrose	Sowthistle	Tumbleweed	Willow
Prostrate knotweed	Spanishneedle	Velvet leaf	Witchweed
Prostrate pigweed	Spatdock	Venice mallow	Woolly morningglory
Prostrate spurge	Speedwell	Veronica	Woodsorrel
Prostrate vervain	Spiny Amaranth	Vervain	Woolly croton
Puncture vine	Spiny cocklebur	Vetch	Woolly plantain
Purslane	Spotted catsear	Virginia buttonweed	Wormseed
Ragweed	Spotted knapweed	Virginia creeper	Yarrow
Red clover	Spotted spurge	Virginia pepperweed	Yellow rocket
Redroot pigweed	Spurge	Wavyleaf bullthistle	Yellow flower pepperweed and other broadleaf weeds.
Red sorrel	Spurweed	Western clematis	
Redstem filaree	Stinging nettle	Western salsify	

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill (or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 to 3 pints in 20 to 240 gallons of water per acre (0.73 to 1.10 fl. oz. in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Ryegrass, Bentgrass (excluding golf course greens and tees), Bahia, Bermudagrass and Zoysia grass. **Controlled Droplet Applicators - (CDA), Atomizers, and Spinning Disk Applicators:** Use this product at the rate of 2 to 3 pints per acre (0.73 to 1.10 fluid ounces per 1,000 square feet) in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment).

Lower Volume Equipment: Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface. **NOTE:** For all grasses (1) Do not overlap spray patterns; (2) Use reduced rates if grass is stressed from heat, drought, etc.; and (3) Follow CDA equipment spray instructions.

Aerial applications are prohibited.

Do not apply more than 1.5 lbs ai MCPA per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

COMPATIBILITY

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test (given below) is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed. The following compatibility test should always be performed prior to full-scale mixing.

1. Pour 18 ounces of water into a quart jar.
2. Add 1 ounce of either the liquid fertilizer or liquid iron to be used.
3. Add 1 ounce of this product.
4. Close jar and shake well.
5. Watch the mixture for several seconds after shaking and check again after 30 minutes.
6. If the mixture does not show signs of separating, the combination may be used. If the mixture foams excessively, gels, separates or gets very thick, do not combine for field application.
7. Compatibility may be improved by the use of a compatibility agent. Some suggested compatibility agents to try are Kalo Laboratories Complex, Hopkins Chemical's Unite, Farm Chemicals Inc.'s Compat, Harcros Chemicals' T-Mulz 734-2, Rigo Company's Rigo Compatibility Agent, Witco Chemical's Sponto 168D, Amoco Oil's Amoco Spray Mate and Universal Coop.'s

Chem-Link. These agents are all used in the same manner. Follow the previously outlined test procedures and add 1/6 ounce of the compatibility agent between steps (the compatibility agent must be added to the fertilizer or iron before adding this product).
8. If the mixture does not separate, gel, foam or get very thick, it may be used for field application. Mix only the amount to be sprayed. Do not allow to stand overnight.

STORAGE AND

DISPOSAL Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be stored in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Or

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

11
/

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV1 02406)

Editor's Note:

The State Sale and Use Prohibitions of this product must be part of printed label text for any product which might possibly enter the channels of trade in the states of California, New York, Oregon and/or Washington. The label prohibitions are optional for all other states.