

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

July 25, 2023

Carrie M. Tackema Sr. Regulatory Manager Nufarm, Inc. 11901 S. Austin Ave. Alsip, IL 60803

Subject: Registration Review Label Mitigation for Bifenthrin Product Name: Bifenthrin Granule Insecticide EPA Registration Number: 228-483 Application Date: 04-26-2021 Decision Number: 573410

Dear Carrie Tackema:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Bifenthrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2 EPA Reg. No. 228-483 Decision No. 573410

If you have any questions about this letter, please contact Samantha Thomas by phone at 202-566-2368, or via email at <u>thomas.samantha@epa.gov</u>.

Sincerely,

Cathury Sitter

Cathryn Britton, Chief Risk Management and Implementation Branch 5 Pesticide Re-Evaluation Division Office of Pesticide Programs

Enclosure

BIFENTHRIN GRANULE INSECTICIDE

FOR OUTDOOR RESIDENTIAL LAWN, HOME PERIMETER, ORNAMENTAL & FLOWER GARDENS, AND HOME GARDEN USE ONLY

ACTIVE INGREDIENT: Bifenthrin*	By Wt. 0.1%	
OTHER INGREDIENTS:		
	TOTAL:	ō

*Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

[SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON BACK OF BAG.] SEE [BACK PANEL] [BOOKLET,] [BACK PANEL] FOR [FIRST AID] [AND] ADDITIONAL PRECAUTIONARY STATEMENTS.

> For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840



EPA REG. NO. 228-483 EPA EST. NO._____

NOTE: Spanish language is optional

NET WEIGHT ____ lbs / (____kg)

A C C E P T E D 07/25/2023

Manufactured [By] For

Alsip, IL 60803

NUFARM AMERICAS INC.

11901 South Austin Avenue

Under the Federal Insecticide, Fungicide

and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2000 4000

228-483

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION - PRECAUCION

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum or using tobacco.

FIRST AID					
IF SWALLOWED	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. 				
IF IN EYES • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.					
HOT LINE NUMBER					
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.					

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and should be avoided.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on the driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a high-performance product and is:

- Virtually Odor-free.
- Fast Acting.
- Easy-to-Use.
- Broad Spectrum.
- Kills Ants, Fleas, Spiders, Ticks & Other Insects.
- Not harmful to grasses.
- For use on lawns, home perimeter, ornamental & flower gardens.
- For Outdoor Multi-insect control.
- Kills Lawn & Garden pests/insects/bugs.
- Kills/controls pests/insects/bugs in lawns & landscapes.

This product is designed for use as a ready-to-use granular insecticide to control various insects and spiders on lawns, home perimeter, ornamental & flower gardens. Please read the following "Directions for Use" carefully and fully before applying this product. For outdoor use only.

In New York State, **DO NOT** apply to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. **DO NOT** water to the point of run-off.

For applications at rates above 0.34 lbs bifenthrin/A ([insert lbs product/A]), users must water the treated area immediately after application. Spot treatments must not exceed two square feet in size (2 ft. by 1 ft. or 4 ft. by 0.5 ft.)

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

DO NOT make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

DO NOT apply when the wind speed is greater than 15 mph.

DO NOT apply the product into fish pools, ponds, streams, or lakes. DO NOT apply directly to sewers or storm drains, or to any area

like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

DO NOT allow the product to enter any drain during or after application.

DO NOT apply to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

DO NOT apply or irrigate to the point of runoff.

INSECT RESISTANCE MANAGEMENT

For resistance management, this product contains bifenthrin, a Group 3A pyrethroid insecticide. Repeated and exclusive use of this product or other Group 3A insecticides may lead to the buildup of resistant strains of insects in some crops.

Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, this product may be used as part of a resistance management strategy established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode-of-action classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local or state agricultural authorities for details.

Unless directed otherwise in the specific crop/pest sections of this label, the best practices are to follow these instructions to delay the development of insecticide resistance:

- Avoid using the same mode of action (IRAC group number) on consecutive generations of insect pests.
- Apply this product using a "treatment window" approach to avoid exposure of successive insect pest generations to the same mode of action.
- A "treatment window" is defined as the period of residual activity provided by single or sequential applications of products with the same mode of action. This "treatment window" should not exceed approximately the length of one generation of the target pest.
- Within the "treatment window", make no more than 2 applications of this product or other product that contains a Group 3A
 insecticide within a single generation of the target pest on a crop or within a 30-day period to the same insect species on a crop.
- Following a "treatment window", rotate to a treatment window of effective products with a different mode of action. This window should approximate the duration of one generation of the target pest.
- Target the most susceptible insect life stages, whenever possible.
- DO NOT use less than the labeled rates of this product when applied alone or in tank mixtures.
- Monitor insect populations for product effectiveness.

If resistance to this product develops in your area, this product or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local Nufarm representative or agricultural advisor for the best alternate method of control for your area. For additional information on insect resistance monitoring, visit the Insecticide Resistance Action Committee (IRAC) on the web at http://www.irac-online.org.

LAWN, HOME PERIMETER, ORNAMENTAL & FLOWER GARDENS

SURFACE-FEEDING INSECTS

Use this product for the control of ants, including Carpenter, Red Harvester, Odorous Pyramid, Pharaoh (Sugar) and Fire Ants^{*}, Foraging Fire Ants, Armyworms (including Fall Armyworms), Billbug (including Bluegrass Billbug), Chiggers, Chinch Bugs, Crickets, Cutworms (including Black Cutworms), Earwigs, Grasshoppers, Mealybugs, Sod Webworms (Lawn Moth), and Weevils (including Hyperodes and Bluegrass). Uniformly apply 1.2 to 2.4 pounds over a 1,000 square feet area by using any hand- or power-operated gravity or rotary type spreader designed for the application of granular materials. A 12-pound bag will treat a 5,000 to 10,000 square feet area. *Optional language:* [XX pound bag should treat XX,XXX square feet. This would be used for a bag size other than 12 pounds. The rate of application i.e. 1.2 to 2.4 pounds of product over1,000 square feet will remain the same.] Best results are obtained if the treated area is thoroughly watered immediately after application. Treat when insects first appear and retreat as necessary. Repeat application is necessary only if there are signs of renewed insect activity. Limit repeat application to no more than once per seven days.

*Aids in control

Argentine Ants, Black Turf Ants, Black Turfgrass Ataenius, Fleas (including Cat Fleas), Leafhoppers, Spiders (including Brown Recluse spiders) and Ticks (including Brown Dog Ticks)

Uniformly apply 2.4 pounds over a 1,000 sq. ft. area. Best results are obtained if the treated area is thoroughly watered immediately after application.

Centipedes, Millipedes, Pavement Ants

Uniformly apply 4.8 pounds over a 1,000 sq. ft. area. Best results are obtained if the treated area is thoroughly watered immediately after application.

Mole Crickets (including Tawny Mole Crickets)

Uniformly apply 2.4 to 4.8 pounds over a 1,000 sq. ft. area. For best results: treat in late spring and water the treated area immediately after application.

For Fire Ant Mounds

To control Fire ants, uniformly apply 1 tablespoon or 3 teaspoons** of this product to the surface of each mound. For best results, 1) apply in cool weather, 65 to 80°F, or in early morning or late evening hours. 2) avoid disturbing ants, 3) treat new mounds as they appear and 4) thoroughly water the treated area immediately after application.

**Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring pesticides.

Home Vegetable Gardens

How to Apply

Apply a single application just prior to planting or after plants emerge and work into the top 4 to 6 inches of soil. Use 5 cups (1.2 lbs.) of product for a 500 sq. ft. area.

Vegetables	Days to wait to harvest	Pests
Green Peas, Edible Pods (Sugar Snap Peas, Snow Peas) Green Beans (Wax Beans, Snap Beans) Dry Beans (Black-eyed Peas, Cowpea)	3 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.
Sweet Corn	1 Day	Ants, armyworm, cutworms, crickets, earwigs, European corn borer, fall armyworm, grasshoppers, southwestern corn borer and weevils.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cavalo Broccoli, Chinese Broccoli, Chinese Cabbage, Kohlrabi	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.
Cucurbit Vegetables Chayote, Citron melon, Cucumber, Edible Gourd Muskmelon (Cantaloupe, casaba, Crenshaw melon, Golden melon, Honeydew Melon, Pineapple melon, Santa Claus melon, Snake melon) Pumpkin, Summer squash (Crookneck, Scallop, Straightneck, Zucchini) Winter squash (Butternut, Calabaza, Hubbard including Acorn and Spaghetti) Watermelon	3 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.
Peppers	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.
Eggplant	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.
Head Lettuce	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.

Home Vegetable Garden Spreader Tips For best results, use a Scotts[®] spreader.

Spreader	5 cups (1.2 lbs.) of product for 500 sq. ft.	Directions	
Scotts AccuGreen® Drop Spreader	3-1/4	Operate spreader the longest way of the garden area. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.	
Scotts SpeedyGreen [®] Rotary	2	Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs.	

When using other spreaders, calibrate the spreader settings based on the manufacturer's instructions.

OPTION I. Directions for Use for the ORTHO Group of The Scotts Company

[Optional Text]

Bifenthrin Granular Insecticide is a high-performance product and is:

- [Virtually Odor-free]
- [Fast Acting]
- [Easy-to-Use]
- [Not harmful to grasses]
- For use on lawns, ornamentals, home perimeter and flower gardens.
- For Outdoor Multi-insect control.
- Kills/controls pests/insects/bugs in lawns & landscapes.
- Kills Ants, Fleas, Ticks and other pests/insects/bugs.

- Controls/kills lawn-destroying/lawn-invading pests/insects/bugs.
- Covers/Treats up to 8,300 sq. ft. per 10 lbs.
- Covers up to 16,600 sq. ft. per 20 lbs.
- Covers up to 5,000 sq. ft. per 6 lbs.
- Creates a pest/insect/bug barrier around your home.
- Exceptional Value!
- Kills Chinch Bugs, Mole Crickets and Cutworms, plus other listed insects.
- Kills surface and sub-surface pests/insects/bugs that ruin your lawn.
- Ready-to-use.
- Easy to spread.
- Soil-penetrating formula reaches deep below ground.
- Deep-reaching formula that kills lawn-destroying insects.
- Provides a barrier of protection around your home.
- Guaranteed to work or your money back.
- Satisfaction Guaranteed.
- The smart solutions to pest/insect/bug problems.
- Solves your pest/insect/bug problems fast.
- Outdoor use only (icon).
- Keeps [Lawn] Insects Away for Up to 2 Months [8 weeks].
- Kills Tough Lawn Insects [and Keeps Them Away for Up to 2 Months [8 weeks]].

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For outdoor use only. Do apply when insect activity or damage is first observed in the lawn (lawn thins and turns brown or moths fly from the grass when you mow). Don't treat just the browned areas of lawn. To prevent further damage, treat entire lawn. Retreat as necessary. Repeat application is necessary only if there are signs of renewed insect activity. Repeat application should be limited to not more than once per seven days.

In New York State **DO NOT** apply to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch). In New York State, make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. **DO NOT** water to the point of run-off. For applications at rates above 0.34 lb. bifenthrin/A (170 lbs. product/A), users must water the treated area immediately after application. For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

DO NOT make applications during rain. **DO NOT** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. **DO NOT** allow the product to enter any drain during or after application.

Product Facts	
Surface Feeding Lawn Insects	Use this product for the control of: Ants (including Argentine, Black Turf, Carpenter, Red Harvester, Odorous, Pyramid, Pharoah), Armyworms (including Fall Armyworms), Billbugs (including Bluegrass Billbugs), Black Turfgrass Ataenius, Chinchbugs, Crickets, Cutworms (including Black Cutworms), Earwigs, Fleas (including Cat Fleas), Grasshoppers, Leafhoppers, Mealybugs, Sod Webworms (Lawn Moth) and Weevils (including Hyperodes and Bluegrass).
How To Use	 Insects (Listed above): Uniformly apply 2.4 pounds over a 1,000 sq. ft. area of lawn using any hand- or power-operated gravity or rotary type spreader designed for the application of granular materials. A 10 pound bag will treat up to 4,167 sq. ft. area of lawn. A XX pound bag will treat up to XXX sq. ft. area of lawn. Best results are obtained if the treated area is thoroughly watered immediately after application. Centipedes, Millipedes, Pavement Ants: Uniformly apply 4.8 pounds over a 1,000 sq. ft. area of lawn. Best results are obtained if the treated area is thoroughly watered immediately after application. For Mole Crickets (including Tawny Mole Crickets): Uniformly apply 2.4 pounds over a 1,000 sq. ft. area of lawn. For best results, treat in late spring and water thoroughly immediately after application. For Fire Ant Mounds: Uniformly apply 1 tablespoon or 3 teaspoons** of granules to the surface of each mound. For best results: 1) apply in cool weather, 65 to 80° F. In early morning or late evening hours, 2) Avoid disturbing ants, 3) treat new mounds as they appear, 4) thoroughly water the treated area after application.

**Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring pesticides.

Product Facts Table (Option 2)

Product Facts – Treats up to 830 sq. ft. (for 1 lb. bag)				
KILLS INSECTS	Ants, including Argentine, Black Turf, Carpenter, Red Harvester, Odorous, Pyramid, Pharoah (sugar) and pavement ants, Armyworms (including Fall Armyworms), Billbugs (including Bluegrass Billbugs), Black Turfgrass Ataenius, Centipedes, Chiggers, Chinchbugs, Crickets, Cutworms (including Black Cutworms), Earwigs, Fleas (including Cat Fleas), Grasshoppers, Leafhoppers, Mealybugs, Millipedes, Mole Crickets (including Tawny Mole Crickets), Sod Webworms (Lawn Moth), Spiders (including Brown Recluse), Ticks (including Brown Dog Ticks) and Weevils (including Hyperodes and Bluegrass).			
WHERE TO USE	On residential lawns, ornamental (flower) gardens, and home vegetable gardens.			

Ornamental Flower Beds (Outdoors)

Pests Controlled: Ants, crickets, earwigs, grasshoppers, spiders and ticks.

How to Apply: Evenly apply 1.2 lb. of product for every 500 sq. ft. area. Repeat application is necessary only if there are signs of renewed insect activity. Limit repeat application to no more than once per seven days.

Home Invading Insects: Barrier Treatment Around House Foundation (Outdoors)

Pests Controlled: Ants, chiggers, crickets, earwigs, grasshoppers, spiders and ticks. **How to Apply:** Evenly apply a 2-foot band of granules next to the house foundation. Use 1.2 lb. for each 500 sq. ft. (2 ft. X 250 ft.). If necessary, repeat applications every 2 weeks only if there are signs of renewed insect activity.

SPREADER TIPS				
For best results, use a Scotts [®] spreader				
Spreader	Spreader Settings covers x,xxx sq. ft.	Directions		
Scotts Drop Spreaders	?	Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.		
Scotts Easy Green [®] Rotary	??	Side shut-off lever to the right; application automatically starts and stops when you do. For complete coverage make each pass about 30 inches from the previous one so some overlapping occurs.		
Scotts SpeedyGreen [®] Rotary	?	Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs. Avoid applying this product on flower, vegetable and shrub plantings.		

(Bag Size specific)

SPREADER TIPS

For best results, use a Scotts[®] spreader

Spreader	Spreader Settings Pounds of Product per 500/1,000 sq. ft. 05/12 1.2/2.4 2.4/4.8		duct per q. ft.	Directions
Scotts AccuGreen [®] Drop Spreader	3-1/4	4-1/2	6	Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.
Scotts SpeedyGreen [®] Rotary	2	2-3/4	4-1/2	Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs. Avoid applying this product on flower, vegetable and shrub plantings.

When using other spreaders, calibrate the spreader settings based on the manufacturer's instructions.

OPTION 1. (The ORTHO Group)

Notice: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

Questions, Comments or Medical Information call 1-800-225-2883

www.ortho.com

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

OPTION II. Directions for Use for the ORTHO Group of The Scotts Company

Bifenthrin Granular Insecticide is a high-performance product, and is: [Virtually Odor-free] [Fast Acting] [Easy-To-Use] [Not harmful to grasses] For Outdoor Multi-insect control. Kills unwanted pests/insects/bugs on contact. Exceptional Value! Ready-to-use. Easy to spread. Guaranteed to work or your money back. Satisfaction Guaranteed The smart solutions to pest/insect/bug problems. Solves your pest/insect/bug problems fast. Keeps [Lawn] Insects Away for Up to 2 Months [8 weeks]. Kills Tough Lawn Insects [and Keeps Them Away for Up to 2 Months [8 Weeks]]. Vegetable Garden Insect Control How to Apply

Apply a single application just prior to planting or after plants emerge and work into the top 4 to 6 inches of soil. Use 5 cups (1.2 lbs.) of product for a 500 sq. ft. area.

Vegetables	Days to wait to harvest	Pests	
Green Peas, Edible Pods (Sugar Snap Peas, Snow Peas) Green Beans (Wax Beans, Snap Beans) Dry Beans (Black-eyed Peas, Cowpea)	3 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	
Sweet Corn	1 Day	Ants, armyworm, cutworms, crickets, earwigs, European corn borer,	
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cavalo Broccoli, Chinese Broccoli, Chinese Cabbage, Kohlrabi	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	
Cucurbit Vegetables Chayote, Citron melon, Cucumber, Edible Gourd Muskmelon (Cantaloupe, casaba, Crenshaw melon, Golden melon, Honeydew Melon, Pineapple melon, Santa Claus melon, Snake melon) Pumpkin, Summer squash (Crookneck, Scallop, Straightneck, Zucchini) Winter squash (Butternut, Calabaza, Hubbard including Acorn and Spaghetti) Watermelon	3 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	
Peppers	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	
Eggplant	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	
Head Lettuce	7 Days	Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	

HOME VEGETABLE GARDEN SPREADER TIPS For best results, use a Scotts [®] spreader.			
	5 cups (1.2 lbs.) of product for 500 sq. ft.	Directions	
Scotts AccuGreen [®] Drop Spreader	3-1/4	Operate spreader the longest way of the garden area. Overlap wheel marks to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.	
Scotts SpeedyGreen [®] Rotary	2	Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs.	

When using other spreaders, calibrate the spreader settings based on the manufacturer's instructions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefuly open containers. After partial use replace lids and close tightly. For bags after partial use, reseal open bag by folding top down and securing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Do not pour or dispose down-the-drain or sewer. Call your local solid waste agency for local disposal options.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

All other trademarks are the property of their respective owners.

(RV[mmddyy])

is a registered trademark of Nufarm Americas Inc.