



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

April 22, 2016

Gregory C. Mattern  
Nufarm Americas Inc.  
4020 Aerial Center Parkway, Suite 101  
Morrisville, NC 27560

Subject: Label Amendment – To update use directions and make other minor labeling changes.  
Product Name: NUP06237 Herbicide  
EPA Registration Number: 228-551  
Application Date: 11/25/2015  
Decision Number: 511618

Dear Mr. Mattern:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.


A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 228-551  
Decision No. 511618

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Karen Samek by phone at (703) 347-8825, or via email at [samek.karen@epa.gov](mailto:samek.karen@epa.gov).

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending to the right.

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

**ACCEPTED**  
 04/22/2016  
 Under the Federal Insecticide, Fungicide  
 and Rodenticide Act as amended, for the  
 pesticide registered under  
 EPA Reg. No. 228-551

# NUP06237

# HERBICIDE

[Alternate Brand Names: Trupower3 Selective Herbicide]

**SELECTIVE BROADLEAF WEED CONTROL FOR TURFGRASSES INCLUDING USE ON SOD FARMS. TO CONTROL CLOVER, DANDELION, HENBIT, PLANTAINS, WILD ONION, AND MANY OTHER BROADLEAF WEEDS. ALSO FOR HIGHWAYS AND RIGHTS-OF-WAY.**

**CONTAINS 2,4-D MECOPROP-p AND DICAMBA**

**ACTIVE INGREDIENTS**

Triisopropanolamine Salt of 2,4-Dichlorophenoxyacetic Acid\* ..... 47.77%  
 Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid \*\*† ..... 7.74%  
 Dicamba Acid (3,6-Dichloro-o-anisic Acid)\*\*\* ..... 3.20%

**OTHER INGREDIENTS** ..... 41.29%  
**TOTAL** ..... 100.00%

Isomer Specific Method, Equivalent to:

\*2,4-Dichlorophenoxyacetic Acid ..... 25.60%, 2.56 lbs./gal.  
 \*\*(+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid ..... 6.40%, 0.64 lbs./gal.  
 \*\*\*3,6-Dichloro-o-anisic Acid ..... 3.20%, 0.32 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p

**KEEP OUT OF REACH OF CHILDREN**  
**DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300**  
**For Medical Emergencies Only, Call (877) 325-1840**

EPA REG. NO. 228-551  
 EPA EST. NO.

**MANUFACTURED BY**  
**NUFARM AMERICAS INC.**  
 11901 S. AUSTIN AVE  
 ALSIP, IL 60803

**NET CONTENTS \_\_\_\_\_ GAL. (\_\_\_\_\_ Liters)**  
**[Designation as "NONREFILLABLE" or "REFILLABLE" for containers > 5 GAL]**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER - PELIGRO**

Corrosive, causes irreversible eye damage. Do not get in eyes, or on skin or clothing. Harmful if swallowed.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are made of any waterproof material.

**All mixers, loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks, and
- Protective eyewear (Goggles or face shield or shielded safety glasses)\*
- Chemical-resistant gloves (except for applicators using groundboom equipment).
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

\* Once this product has been diluted with water at a dilution factor of at least 13.6X or more [3 pints of product in 5 gallons], eye protection is not required.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
Wash hands, face, and arms with soap and water before eating, smoking, drinking or using the toilet.	
Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.	
Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.	

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers, other persons, or pets, either directly or through drift. Only protected handlers may be in the area during application. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

Aerial application is prohibited.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls worn over short-sleeve shirt and short pants, chemical-resistant footwear plus socks, chemical-resistant gloves made of any waterproof material, chemical-resistant headgear for overhead exposure, and protective eyewear.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people or pets to enter the treated area until sprays have dried.

## SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Do not allow the herbicide solution to drift onto desirable broadleaf plants, as small amounts of product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

### Droplet Size

Use a medium or coarser spray nozzles according to (ASAE standard 572) definition for standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

### Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

## USE RESTRICTIONS

Do not apply this product through any type of irrigation system. Aerial application of this product is prohibited. Product should not be used in or near greenhouses. For turf use, the maximum number of broadcast applications per treatment site is 2 per year.

## USE PRECAUTIONS

This product is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs). Also for use on Sod Farms.

Avoid mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Avoid fine mists. Except as noted, use only lawn-type sprayers. Coarse sprays are less likely to "wind-drift". Use coarse spray droplets. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the dripline of trees and other species. Do not apply to newly seeded grasses until well established. When air temperature exceeds 90°F, temporary turf injury may result from broadcast applications. Limit applications to spot or directed applications.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only to avoid temporary turf injury.

Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results: (1) turf should not be mowed for 1 to 2 days before and after application; (2) do not apply if rainfall is expected within 4 hours; delay irrigation cycle for 24 hours. Reseed no sooner than three to four weeks after application of this product. Failure to observe all precautions may result in injury to turf and/or susceptible plants.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizer and liquid iron differ in pH, free ammonia content, density, salt concentration and percentage of water, a compatibility test is recommended prior to mixing in application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

## WEEDS CONTROLLED

Bedstraw	Dandelion	Lespedeza	Poison oak	Wild carrot
Black medic	Dock	Mallow	Purslane	Wild garlic
Buckhorn	Ground ivy	Morningglory	Ragweed	Wild lettuce
Burdock	Healall	Peppergrass	Sheep sorrel	Wild onion
Chicory	Henbit	Pigweed	Shepherdspurse	Yarrow
Chickweed	Knotweed	Plantain	Speedwell	
Clover	Lambsquarter	Poison ivy	Spurge	

**TURF, ORNAMENTAL**  
**(golf courses, cemeteries, parks, sports fields, turfgrass, lawns )**  
**AND**  
**TURF (grown for seed or sod)**

**APPLICATION RATES**

SITE	USE RATE (Fluid Ounces)	SPRAY VOLUME (Gallons)	USE RATE (Pints)	SPRAY VOLUME (Gallons)	USE DIRECTIONS
	PER 1,000 SQ. FT.		PER ACRE		
Bahagrass, Bluegrass, Common Bermudagrass, Fescue, Ryegrass, Zoysiagrass	1.1 to 1.5	0.5 to 5	3 to 4	5 to 220	<b>Normal Applications</b>
					<b>For Professional Lawn Maintenance</b> - Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.
St. Augustinegrass and Centipedegrass	0.75 to 0.90 Maximum	1 to 5	2 to 2.5 Maximum	40 to 200	Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses. <b>If Bermudagrass is dormant</b> , up to 4 pints per acre may be used. However, some hybrid Bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.
Bentgrass (Putting and Bowling Greens)	0.67 Maximum	3.3	1.8 Maximum	145	Apply on closely mowed Bentgrass, preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

**NOTE:** Care should be taken to avoid overdosing Bentgrass, St. Augustine, and Centipede grasses or injury may result. Large volumes of spray water (i.e. one fluid ounce in 2 to 4 gallons of water per 1,000 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

Ornamental Turf: Do not apply more than 2 broadcast applications per year, excluding spot treatments, with a minimum of 30 days between applications. Do not apply more than 4.68 pints of this product per acre per application. Do not apply more than 9.37 pints of this product per acre per year, including all broadcast and spot treatments combined.

Turf Grown for Seed/Sod: Do not apply more than 2 broadcast applications per year, excluding spot treatments, with a minimum of 30 days between applications. Do not apply more than 6.25 pints of this product per acre per application. Do not apply more than 12.5 pints of this product per acre per year, including all broadcast and spot treatments combined.

**LOW VOLUME SPRAY APPLICATION EQUIPMENT**

EQUIPMENT	APPLICATION	USE RATE		USE DIRECTIONS
		Fluid Ounces per 1,000 Square Feet	Pints per Acre	
<b>Controlled Droplet Applicators (CDA), Atomizers and Spinning Disk Applicators</b>	Cool season grasses	1.1 to 1.5	3 to 4	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment). Do not overlap spray patterns.
	Listed warm season grasses	0.75 to 0.9	2 to 2.5	Use in sufficient water to assure coverage (1 to 4 gallons of water per acre is normal for this type of equipment); and follow CDA spray instructions for cool season grasses. Use reduced rates if grass is stressed from heat, drought etc.
<b>Lower Volume Equipment</b>	Cool Season and Listed Warm Season Grasses	1.1 to 1.5	3 to 4	Use as little as 5 gallons of water per acre. Use only application equipment that is capable of spreading a uniform droplet, wetting each weed surface.

**NON-CROPLAND**

**(fencerows, hedgerows, roadsides, ditches, rights-of-way, utility, powerlines, railroads, airports, and similar industrial sites)**

**Roadsides (including aprons and guardrails), rights-of-way:** For control of broadleaf weeds, mix at a rate of 1.5 to 3.125 quarts of this product per acre in enough water to wet all parts of the foliage. This may require 50 to 300 gallons of water per acre. This mixture will cover 1 acre (43,560 square feet). Thoroughly saturate all weeds with spray mixture. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with the high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of this product per gallon of water and spray to thoroughly wet all foliage, but do not exceed 6.25 pints of product per acre per application.

**Annual/Perennial Control:** Do not apply more than 2 broadcast applications per year, excluding spot treatments, with a minimum of 30 days between applications. Do not apply more than 6.25 pints of this product per acre per application. Do not apply more than 12.5 pints of this product per acre per year, including all broadcast and spot treatments combined.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species.

**Woody Brush:** Add 1 to 1.2 gallons of this product per acre to wet all parts of the brush foliage, stem and bark. This may require up to 200 to 600 gallons of water per 43,560 square feet depending upon the height and thickness of the brush. Mix thoroughly before spraying.

**Woody Plant Control:** Make no more than 1 broadcast application per year, excluding spot treatments. Do not apply more than 9.37 pints per acre per application, and do not apply more than 12.5 pints of this product per acre per year, including all broadcast and spot treatments combined.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions for application.

**Maximum Seasonal Application Rate to non-cropland sites is 4 pounds 2,4-D Acid equivalent per acre per application site.**

**Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. al v. EPA, C01-0132C, (W.D.WA). For further information, please refer to <http://www.epa.gov.espp>.**



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER DISPOSAL:

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

**Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Or**

**Refillable containers larger than 5 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

## TANK MIXES

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**IMPORTANT: PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.**

## COMPATIBILITY

Before full-scale mixing of this product with other pesticides, fertilizers, secondary plant nutrients, adjuvants, surfactants or oils, you must determine the compatibility of the proposed mixture. Use proportionate quantities of each ingredient and mix in a small container. Always mix one product thoroughly with the diluent before adding another product. If no incompatibility is evident after 30 minutes, the mixture is generally compatible for spraying. To evaluate potential short term effects of applying the mixture, test the tank mix combination on a few plants or a small area before larger-scale treatments. Wait at least 2 to 3 days for problems to become apparent.

IMPORTANT: MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES, REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS. ANY LIABILITY FOR LOSS, INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER.

#### **WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER, AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, CROP OR PLANT DAMAGE, OR LOSS OF YIELD, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE WEATHER, WIND, AND TEMPERATURE, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER, AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER OR MANUFACTURER, AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY OTHER REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS. NO WARRANTIES SHALL BE CREATED BY COURSE OF DEALING, USAGE OF TRADE, OR COURSE OF PERFORMANCE. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THE FACE HEREOF. THE SELLER OR MANUFACTURER ASSUMES NO RESPONSIBILITY THAT THE GOODS WILL BE FIT FOR ANY PARTICULAR PURPOSE FOR WHICH YOU MAY BE BUYING OR USING THE GOODS, EXCEPT AS OTHERWISE PROVIDED IN THE CONTRACT.

#### **LIMITATION OF LIABILITY WARNING**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NEITHER MANUFACTURER NOR SELLER SHALL BE LIABLE TO BUYER OR USER OR TO CUSTOMERS OF BUYER, IF ANY, FOR INDEMNIFICATION OR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE USE, MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE GOODS AND, IF BUYER OR USER WISHES, THE RETURN OF THE GOODS BY BUYER TO SELLER.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the goods, and return it unopened to the Seller, and the purchase price will be refunded.**

*By using the goods, you expressly agree to all of the terms and conditions of this contract.*

(RV\_\_\_\_\_)

Trupower and The Optical Advantage is a registered Trademark of Nufarm Americas Inc.

**OPTIONAL MARKETING CLAIMS – FRONT PANEL**

**GET THE OPTICAL ADVANTAGE®**

This product contains a purified form of MCPP known as the optical isomer, which is the portion of the molecule active as an herbicide. The Optical Advantage ® is not associated with user eye safety.

Nufarm Grow a better tomorrow.  
Grow a better tomorrow.