



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

March 15, 2022

Carrie Takema
Regulatory Manager
Nufarm Americas, Inc.
4020 Aerial Parkway, Suite 101
Morrisville, NC 27560

Subject: Label Amendment – To add a California Proposition 65 Statement
Product Name: NUP-13026 Abamectin 0.15 Miticide/Insecticide
EPA Registration Number: 228-734
Application Date: July 11, 2018
Decision Number: 542745

Dear Ms. Takema:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 228-734
Decision No. 542745

with FIFRA section 6. If you have any questions, please contact Joseph Belsky by phone at 202-566-2495, or via email at belsky.joseph@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Gene Benbow".

Gene Benbow, Product Manager 7
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

RESTRICTED USE PESTICIDE

**TOXIC TO FISH, MAMMALS, AND AQUATIC ORGANISMS
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER
THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED
APPLICATOR'S CERTIFICATION.**

ABAMECTIN GROUP 6 INSECTICIDE

NUP-13026 Abamectin 0.15 Miticide/Insecticide [ABN: Abamex Miticide/Insecticide]

ACTIVE INGREDIENT:

Abamectin (CAS No.71751-41-2):..... 2.0%*

OTHER INGREDIENTS:.....98.0%

TOTAL:.....100.0%

*Contains 0.15 pounds Abamectin per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

SEE [NEXT PAGE] [BELOW] FOR ADDITIONAL FIRST AID AND PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

EPA REG. NO. 228-734
EPA EST. NO.:

MANUFACTURED FOR
NUFARM AMERICAS INC.
11901 South Austin Avenue
Alsip, IL 60803



[Nufarm Grow a better tomorrow]
[Grow a better tomorrow]

NET CONTENTS _____ GALS. (_____ Liters)

000228.00734.20220315

ACCEPTED
03/15/2022
Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 228-734

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made out of barrier laminate or butyl rubber \geq 14 mils
- Socks plus shoes

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscle tremors. Toxicity following accidental ingestion of this product can be minimized by early administration of chemical adsorbents (e.g., activated charcoal). If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parenteral fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since Abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic Abamectin exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from target areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. **DO NOT** apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a medium potential for reaching both surface water and aquatic sediment via runoff for several weeks to months after application. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of Abamectin from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecast to occur within 48 hours.

Use of this product may pose a risk to threatened and endangered species of fish, amphibians, crustaceans (including fresh water shrimp), and insects. All use of this product in the state of California should comply with the recommendations of the California Endangered Species Project. Before using this product in California, consult with your county agriculture commissioner to determine use limitations that apply in your area.

Runoff Prevention: To protect the environment, **DO NOT** allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

ATTENTION: This product contains chemicals known to the state of California to cause birth defects or other reproductive harm.

DIRECTIONS FOR USE

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: For grape girdling cane turning, and tying in grapes. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made out of barrier laminate or butyl rubber \geq 14 mils
- Shoes plus socks

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, AND/OR ILLEGAL RESIDUES.

PRODUCT INFORMATION

When applied following the **Directions for Use** on this label, this product controls listed mites and insects in various crops. Optimum control of target insects requires this product to be mixed with sufficient amounts of water to provide a thorough coverage of foliage. Spray volume in gallons will vary with the size and number of trees per acre or with the density of target crop foliage. Optimum insect control is achieved only if the foliage receives a thorough coverage

of this product. Calibrate all spray equipment to ensure delivery of sufficient spray solution to thoroughly cover foliage of target crops.

Phytotoxicity: Although this product has been tested for phytotoxicity and compatibility and is safe to use on a variety of crops and as a tank mix in combination with many commonly used products, it is not possible to test all crop varieties grown under a variety of growing conditions or to test all possible combination of products. Before use, test potential mixtures with this product to ensure physical compatibility and lack of phytotoxic effects to the target crop.

New York State Restrictions: Aerial applications of this product are prohibited in the State of New York.

DO NOT apply this product through any type of irrigation system.

[OPTIONAL TEXT FOR TIP-N-MEASURE CONTAINERS]

Use Directions for Tip-N-Measure Container:

1. Remove the measuring chamber cap and seal over container opening.
2. Replace the cap and securely tighten.
3. Tip the container until the measuring chamber contains the desired amount of product.
4. Return container to its level position.
5. Remove measuring chamber cap and pour product into proper application equipment.
6. Replace the cap and securely tighten.

SPRAY DRIFT MANAGEMENT

NOTE: When states have more stringent regulations, they must be observed.

Spray Drift Precautions for Application with Aircraft or Ground Application Equipment

- **DO NOT** apply with ground application within 25 feet of or with aircraft within 150 ft. of lakes, reservoirs, rivers, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish farm ponds.
- **DO NOT** cultivate within 25 ft. of the aquatic area to allow growth of a vegetative filter strip.
- Observe the following precautions when using ground application to spray tree crops or hops in the vicinity of aquatic areas such as lakes, reservoirs, permanent streams, marshes, potholes, natural ponds, estuaries, or commercial fish farm ponds:
 - **DO NOT** apply this product when weather conditions favor drift to aquatic areas.
 - **DO NOT** apply within 110 ft. upwind of aquatic areas or when wind speed is above 8 mph.
 - Spray last 3 rows windward of aquatic areas using nozzles on one side only, with spray directed away from the aquatic areas.
 - Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove/orchard when spraying the outside row. Shut off nozzles when turning at ends of rows and passing tree gaps in rows.

SPRAY DRIFT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must not exceed 65% of the wingspan for fixed wing aircraft or 75% of the rotor diameter for helicopters. Otherwise, the boom length must not exceed 75% of the wingspan for fixed wing aircraft or 90% of the rotor diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Airblast Applications:

- Sprays must be directed into the canopy.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at the row end and when spraying the outer row.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1)
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications:

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 15 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

SPRAY DRIFT

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

SPRAY DRIFT

Handheld Technology Applications:

- Take precautions to minimize spray drift.

RESISTANCE MANAGEMENT

For resistance management, this product is a Group 6 insecticide. Any insect population may contain or develop plants naturally resistant to this product and other Group 6 insecticides. The resistant biotypes may dominate the insect population if these insecticides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 6 insecticides within a growing season sequence or among growing seasons with different insecticide groups that control the same insects.
- Use tank mixtures with insecticides from a different group if such use is permitted; where information on resistance in target insect species is available, use the less resistance-prone partner at a rate that will control the target insect(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated insect-management program for insecticide use that includes scouting and uses historical information related to insecticide use and rotation and that considers mechanical control methods, cultural, biological and other management practices.
- Scout before and after insecticide application to monitor insect populations for early signs of resistance development. Indicators of possible insecticide resistance include: (1) failure to control a insect species normally controlled by the insecticide at the dose applied, especially if control is achieved on other insects; (2) a spreading patch of non-controlled insects of a particular insect species; (3) surviving insects mixed with controlled individuals of the same species. If resistance is suspected, prevent insect production in the affected area by an alternative insecticide from a different group or by a mechanical method. Prevent movement of resistant insects to other areas by cleaning equipment.
- If a insect pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or insecticide with a different mode of action, if available.

- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated insect-management recommendations..
- For further information or to report non-performance or suspected resistance, contact Nufarm at 1-800-345-3330.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant insects to this MOA have been found in your region. **DO NOT** assume that each listed insect is being controlled by this mechanism of action. Co-formulated active ingredients are intended to broaden the spectrum of insects that are controlled. Some insects may be controlled by only one of the active ingredients in this product.

CROP USE DIRECTIONS

AVOCADOS

Target Insects	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Avocado thrips (<i>Scirtothrips perseae</i>) Persea Mites (<i>Oligonychus perseae</i>)</p>	<p>DO NOT use less than 10 fl. oz./A of this product</p> <p>Low to moderate insect infestations: 10.0-15.0 Heavy pest infestations: 16.0-20.0</p> <p>ADD a horticultural oil for best results</p>	<p>If using 400 gal/A or more, apply 2.5 fl. oz. of this product/100 gal of water and 1.0-4.0% of NR (narrow range) 415 oil.</p> <p>Although aerial applications are permitted when this product is applied in 50 or more gal water/A, better control of target insects is obtained using ground applications. Users accept all liability for level and duration of thrip control from aerial applications of this product.</p> <p>Apply this product when thrips first appear but prior to finding 5 immature thrips per leaf or fruit. For low to moderate infestations, include 1.0-4.0% of NR 415 oil in combination with this product and the horticultural spray oil approved for use on avocados.</p> <p>Make a second application, if needed, to maintain control, but wait at least 30 days after the first application to retreat. SEE RESTRICTIONS.</p>	<p>Apply using ground equipment in 100 or more gal water/A.</p> <p>Pre-Harvest Interval – 14 days</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 40 fl. oz./A (0.047 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT allow livestock to graze in treated orchards.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil), minimum 1 to 4%, v/v (minimum)</p> <p>Application Interval: Wait at least 30 days before repeating application.</p>
<p>IMPORTANT:</p> <ul style="list-style-type: none"> • Low infestations are defined as 1-2 immature thrips per leaf/fruit • Moderate infestations are defined as 3-4 thrips per leaf/fruit; • Heavy/severe infestations are defined as more than 5 thrips per leaf/fruit. <p>To prevent crop injury, observe all precautions and restrictions on the oil label. When using oil concentrations above 2.0%, treat a small test area before making a large-scale application.</p>			

BEAN, DRY AND SUCCULENT

Crops in this group include *Cicer arietinum* (chickpea, garbanzo bean); *Lupinus* spp. (including grain lupin, sweet lupin, white lupin, and white sweet lupin); *Phaseolus* spp. (including kidney bean, lima bean, mung bean, navy bean, pinto bean, snap bean, and waxbean); *Vicia faba* (broad bean, fava bean); *Vigna* spp. (including asparagus bean, blackeyed pea and cowpea*)

* For use on cowpeas that are grown only for dry seed. Do not allow livestock to graze cowpea forage and do not harvest cowpea forage or hay for use as livestock feed.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Spider mite</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 13.0-16.0</p>	<p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p> <p>Leafminers and Spider mites: Apply when adult leafminers or spider mites first appear, and repeat as needed to maintain control within constraints of a sound resistance management program.</p> <p>Make a second application, if needed, but wait at least 6 days after the first application. SEE RESTRICTIONS.</p>	<p>Do not apply in less than 10 gal. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft.</p> <p>Pre-Harvest Interval – 7 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Application Interval: Wait at least 6 days before repeating application.</p>

CELERIAC (*Apium graveollens*)

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Twospotted spider mite	16.0	<p>Apply when mites first appear.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p> <p>For best results, use in combination with a non-ionic surfactant.</p>	<p>Ground application only</p> <p>Apply using ground equipment using conventional dilute or concentrate sprayers.</p> <p>Use sufficient volumes (greater than 20 gal of water per acre) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 7 days</p> <p>To avoid insect resistance to this product, do not apply more than two sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year after transplanting.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

CITRUS FRUIT CROP GROUP 10-10

Crops in this group include Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; mount white lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; Satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin); tangor; trifoliate orange; uniq fruit; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
Broad mite Citrus bud mite Twospotted spider mite	Concentrate Sprays: 10-20 fl. oz./A (Dilute Sprays: 1.0-2.0 fl. oz./100 gal)		Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough coverage (or outside coverage for Asian citrus psyllid, citrus leafminer, or citrus thrips).
Citrus rust mite	Concentrate Sprays: 5-20 fl. oz./A (Dilute Sprays: 0.5-2.0 fl. oz./100 gal)	Apply using conventional dilute or concentrate ground equipment.	Apply this product with a minimum of 0.20% horticultural oil spray (not a dormant oil) in the spray mixture or not less than 1 gallon of oil/A (except when specified differently below).
Citrus leafminer	Concentrate Sprays: 5-20 fl. oz./A (Dilute Sprays: NR)	Aerial application is permitted only for control of citrus leafminer. Aerial application to citrus is not approved in California. ADD a horticultural oil for best results. Asian citrus psyllid: Apply to protect newly expanding foliage flush during the spring, summer or fall. This product will only control adults and nymphs present at the time of application. Mites: Apply when mites first appear during spring, summer, and/or fall.	To control citrus leafminer -Ground or aerial application is permitted: however, do not apply with aircraft to citrus in California. For Aerial application use a minimum of 10 gallons of finished spray volume per acre. Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures) use a greater volume of water to insure adequate coverage.
Asian citrus psyllid Citrus thrips	Concentrate Sprays: 10-20 fl. oz./A (Dilute Sprays: NR)	Citrus bud mite: For best results, time the spray at "bud swell". Citrus leafminer: Apply to protect new growth during spring, summer, or fall.	All other pests: Ground application only. Do not apply with aircraft. Pre-Harvest Interval – 7 days To avoid insect resistance to this product, do not apply to citrus grown in nurseries.

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
			<p>DO NOT apply this product, or any other Abamectin-containing product within 30 days of the last treatment.</p> <p>DO NOT apply more than 40 fl. oz./A (0.047 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make more than 3 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT allow livestock to graze in treated orchards.</p> <p>Application is not permitted from onset of flowering until after petal fall is complete.</p>

NR = Not recommended.

IMPORTANT: The rate of this product per 100 gallons is based on a volume of 1,000 gal/A dilute spray. For concentrate sprays, adjust the dosage to apply the amount/A equal to that used in full cover dilute spray. Ground application: For best results use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 mph. For best results, use 100-250 gal/A of spray mix applied to achieve outside coverage.

COTTON

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Carmine spider mite Pacific spider mite Strawberry spider mite Twospotted spider mite	Early season: 4.0-6.0	<p>Apply this product when mites first appear.</p> <p>Rate will depend on the cotton plant size and thickness of foliage. Use the 4-6 fl. oz./A rate only early in season and for cotton plants 10 inches or less in height. Do not use less than 4 fl. oz./A.</p> <p>West of the Rockies: The lower use rates may only be used on cotton less than 10 inches in height and applied only with ground equipment.</p> <p>Note: This product may be used without a wetting agent. Spreading and penetrating surfactants can improve mite control. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended. Do not use binder or sticker-type surfactants.</p>	<p>Applications can be made by air or ground using a minimum of 5 gallons of water/A. Use sufficient volume to ensure thorough coverage.</p> <p>DO NOT apply more than 16 fl. oz/A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 32 fl. oz./A (0.038 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>Pre-Harvest Interval – 20 days</p> <p>DO NOT graze livestock in treated fields or feed cotton foliage to livestock.</p> <p>Application Interval: Wait at least 21 days before repeating application.</p>
	Mid-season/Lay-by: 8.0-16.0	<p>Apply this product when mites first appear.</p> <p>Rate will depend on the cotton plant size and thickness of foliage.</p>	

CUCURBIT VEGETABLE CROP GROUP (CROP GROUP 9)

Crops in this group include chayote, Chinese waxgourd, citron melon, cucumbers, gherkin, edible gourds (includes hyotan, cucuzza, hechima, Chinese okra); *Momordica* spp., (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon, (hybrids and/or cultivars of *Cucumis melo* includes true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon), pumpkin, summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), winter squash, (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), watermelon (includes hybrids and/or varieties of *Citrullus lanatus*)

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Leafminers Spider mite</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 16.0</p>	<p>Users accept all liability for level and duration of mite control from aerial applications of this product.</p> <p>Apply when insects first appear.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p>	<p>Use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications.</p> <p>Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 7 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

FRUITING VEGETABLES CROP GROUP (except Cucurbits) (CROP GROUP 8-10)

Crops in this group include African eggplant; bush tomato; bell pepper; cocona; currant tomato; eggplant; garden huckleberry; goji berry; groundcherry; martynia; naranjilla; okra; pea eggplant; pepino; nonbell pepper; roselle; scarlet eggplant; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Broad mite Colorado potato beetle <i>Liriomyza</i> leafminers Potato psyllid Spider mite Thrips palmi Tomato psyllid Tomato Russet mite	Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0 ADD a nonionic surfactant for best results	Apply when insects first appear. Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS. Users accept all liability for level and duration of mite control from aerial applications of this product. A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray. Mites: For best control apply with ground equipment. Broad, russet, and spider mites: Apply when mites first appear. Thrips palmi: Apply when thrips are first observed. Leafminers: Apply when adult flies are first observed Potato Psyllids: Sequential application 10-14 days apart may be needed for infestations with heavy egg and nymph populations.	Do not apply in less than 20 gal water/A with ground equipment. Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure adequate coverage. Pre-Harvest Interval – 7 days DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application. DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year. DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product. DO NOT apply by air in the State of New York. For resistance management, DO NOT use on fruiting vegetables grown for transplanting. Application Interval: Wait at least 7 days before repeating application.
Tomato pinworm	16.0	Tomato pinworm: Application may be made from the time moth activity is detected up to, but no later than, the time when newly emerged larvae are present.	

**GRAPES AND SMALL FRUIT VINE CLIMBING SUBGROUP (except Fuzzy Kiwifruit)
(CROP SUBGROUP 13-07F)**

Crops in this group include Amure river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Pacific spider mite Twospotted spider mite Variegated leafhoppers* Western grapeleaf skeletonizers Western grape* leafhoppers Willamette spider mite</p> <p>*NOTE: This product only provides knock-down control of pest.</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 16.0</p> <p>ADD a nonionic surfactant for best results</p>	<p>Spider mites: Apply when mites first appear (or before more than 5 motiles per leaf are observed).</p> <p>Western grapeleaf skeletonizer: Apply with a non-ionic surfactant when the larvae are initially seen. For best results, apply shortly after eggs hatch.</p> <p>Leafhoppers: For contact knock-down control of these pests, apply with a non-ionic surfactant when pest first appears.</p> <p>European Grapevine Moth: Apply to aid in control of first generation of pest on grapes.</p> <p>Make a second application, if needed, but wait at least 21 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic (not binder or sticker-type) surfactant must be used in combination with this product to provide improved wetting of foliage and an even deposition of the spray.</p> <p>It is impossible to test the safety of this product plus non-ionic surfactants on all grape varieties under the various conditions which lead to phytotoxicity. Therefore carefully follow the directions for use on the nonionic surfactant label.</p>	<p>Use conventional ground application equipment in 50 gal or more of water/A to ensure thorough foliage coverage, or with electrostatic spray equipment in 5 gal or more of water/A.</p> <p>DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: For grape girdling cane turning, and tying in grapes. DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.</p> <p>Pre-Harvest Interval – 28 days</p> <p>DO NOT exceed 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application</p> <p>DO NOT apply more than 32 fl. oz./A (0.038 lb ai/A) of this product or any other Abamectin-containing product per growing year.</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT graze livestock in treated vineyards.</p> <p>DO NOT apply by air.</p>
<p>European Grapevine Moth (<i>Lobesia botrana</i>)</p>	<p>10-16</p>		<p>Application Interval: Wait at least 21 days before repeating application.</p>

GUAVA

Crops in this group include guava; feijoa; jaborcaba; wax jambu; starfruit; passionfruit; and acerola

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
Twospotted spider mite	<p style="text-align: center;">Low to moderate pest infestations: 10.0-15.0</p> <p style="text-align: center;">Heavy pest infestations: 16.0-20.0</p>	<p style="text-align: center;">ADD a horticultural oil for best results</p> <p style="text-align: center;">Apply when mites first appear during spring, summer, and/or fall.</p>	<p>Ground application only.</p> <p>Do not apply in less than 50 gal water/A with conventional ground application equipment. When using an electro-static sprayer, less than 50 gal water/A may be used; however, do not use less than 5 gal water/A.</p> <p>Pre-Harvest Interval – 7 days</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT make more than 3 applications of this product, or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 60 fl. oz./A (0.07 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>Application Interval: wait at least 14 days before repeating application.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.20%, v/v).</p>
<p>IMPORTANT: Concentrate Spray rates (this product /A) are adjusted to apply the amount/A equal to that used in full cover dilute spray. Rates of this product/100 gal are based on 1,000 gal/A of dilute spray. Ground application: For best results, use 150-300 gal/A spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 mph.</p>			

HERB CROP SUBGROUP (CROP SUBGROUP 19A)

Crops in this group include angelica, balm, basil, borage, burnet, camomile (or chamomile), catnip, chervil (dried), chives, chives (Chinese), cilantro (leaf), clary, coriander (leaf), costmary, culantro (leaf), curry (leaf), dill weed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram (*Origanum* species), nasturtium, parsley (dried), pennyroyal, rosemary, rue, sage, summer and winter savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff, and wormwood

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Spider mite</p>	<p>Low to moderate pest infestations: 8.0</p> <p>Heavy pest infestations: 16.0</p> <p>ADD a nonionic surfactant for best results</p>	<p>Apply when adult leafminer flies first appear.</p> <p>Make a second application, if necessary, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p> <p>Do not use a surfactant when applying this product to chives.</p> <p>Although this product has been tested in combination with a nonionic surfactant for safety to herbs, it is impossible to test on all herb types and varieties under the variety of conditions that may cause crop injury.</p>	<p>Apply using ground equipment using conventional dilute or concentrate sprayers.</p> <p>Use sufficient volumes (greater than 20 gal of water per acre) to ensure thorough foliage coverage.</p> <p>Application is not permitted from onset of flowering until after petal fall is complete.</p> <p>Pre-Harvest Interval for listed herbs except chive– 14 days.</p> <p>Pre-Harvest Interval for chives - 7 days.</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any Abamectin-containing product per application.</p> <p>DO NOT make more than 2 applications of this product or any Abamectin-containing product per single cutting (harvest).</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any Abamectin-containing product per year.</p> <p>DO NOT apply by air.</p> <p>DO NOT graze livestock in treated fields or feed treated foliage to livestock.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

HOPS (EXCLUDING CALIFORNIA)

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Twospotted spider mite	<p style="text-align: center;">8.0-16.0</p> <p style="text-align: center;">ADD a nonionic surfactant for best results</p>	<p>Use conventional ground application as dilute or concentrate sprays. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. If trellis height is 6-8 feet, use 40 gal of water/A at the 8-16 fl. oz./A rate; if height is greater than 6-8 feet, use 100 gal/A at the 16 fl. oz./A rate.</p> <p>Apply when twospotted spider mites reach treatment thresholds.</p> <p>Make a second application, if needed, but wait at least 21 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Ground application only.</p> <p>Pre-Harvest Interval – 28 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any Abamectin-containing product per application.</p> <p>DO NOT apply more than 32 fl. oz./A (0.038 lb ai/A) of this product or any Abamectin-containing product per year.</p> <p>DO NOT apply by air.</p> <p>DO NOT apply more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT graze livestock in treated hop yards.</p> <p>Application Interval: Wait at least 21 days before repeating application and repeat application only after using an alternative miticide with a different mode of action.</p>

LEAFY VEGETABLE (Except Brassica Vegetables) CROP GROUP (CROP GROUP 4)

Crops in this group include amaranth (leafy amaranth, Chinese spinach, tampala), arugula (rocket), cardoon, celery, celtuce, chervil, Chinese celery, chrysanthemum (edible leaved and garland), corn salad, cress (garden and upland (yellow rocket and winter cress)), dandelion, dock (sorrel), endive (escarole), fennel, Florence fennel, lettuce (head and leaf), New Zealand spinach, orach, parsley, purslane (garden and winter), radicchio (red chicory), rhubarb, spinach, Swiss chard and vine spinach

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Carmine spider mite <i>Liriomyza</i> leafminers Twospotted spider mite</p>	<p>Low to moderate pest infestations: 10.0 – 15.0</p> <p>Heavy pest infestations: 16.0</p> <p>ADD a nonionic surfactant for best results</p>	<p>Users accept all liability for level and duration of insect control from aerial applications of this product.</p> <p>Spider mites: Apply when mites first appear, and repeat as needed to maintain control.</p> <p>Leafminers: Apply when adult flies are first observed and repeat applications as needed to maintain control.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Apply this product when mites or adult leafminer flies are observed and use ground equipment in 20 or more gal water/A, or with electrostatic spray equipment in 5 gal or more of water/A.</p> <p>Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications.</p> <p>Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 7 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>To avoid insect resistance to this product, do not apply to vegetables grown for transplants.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

LYCHEE

Crops in this group include lychee, longan, Spanish lime, rambutan, and pulasan

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
<p>Twospotted spider mite</p> <p><i>Liriomyza</i> leafminers</p> <p>Thrips</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 20.0</p>	<p>ADD a horticultural oil for best results</p> <p>Mites: Apply when mites first appear during spring, summer, and/or fall.</p> <p>Leafminers: Apply to protect new growth during spring, summer, or fall.</p> <p>Thrips: Applications targeted for thrips will only control the current generation and must be correctly timed. Apply when economic thresholds have been reached (after egg hatch has begun – preferably early to mid-hatch).</p>	<p>Apply using conventional dilute or concentrate ground equipment.</p> <p>Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 14 days</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 40 fl. oz./A (0.046 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>Application Interval: wait at least 30 days before repeating application.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.20%, v/v)</p>
<p>IMPORTANT: Concentrate Spray rates (this product/A) are adjusted to apply the amount/A equal to that used in full cover dilute spray. Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 mph. Rates of this product/100 gal are based on 1,000 gal/A of dilute spray.</p>			

MINT (PEPPERMINT AND SPEARMINT)

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Twospotted spider mite	<p align="center">8.0-12.0</p> <p align="center">ADD a nonionic surfactant for best results</p>	<p>Apply this product when mites first appear. Users accept all liability for level and duration of insect control from aerial applications of this product.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic surfactant (not binder or sticker-types) used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray on the foliage.</p>	<p>Use ground equipment in 20 or more gal water/A. Although aerial applications are permitted when this product is applied in 5 or more gal water/A, better control of target insects is obtained using ground applications.</p> <p>Increased water volumes are required under certain situations (severe pest infestations, highly dense foliage, elevated temperatures) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 28 days</p> <p>DO NOT exceed 12 fl. oz./A (0.014 lb ai/A) of this product or any Abamectin-containing product per application.</p> <p>DO NOT apply more than 36 fl. oz./A (0.042 lb ai/A) of this product or any Abamectin-containing product per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply by air in the State of New York.</p> <p>DO NOT graze livestock in treated fields or feed treated foliage to livestock.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

ONION, BULB (CROP SUBGROUP 3-07A)

Crops in this group are bulb onion, including bulb Chinese, bulb daylily, bulb fritillaria, bulb garlic, bulb great-headed garlic, bulb lily, bulb onion, pearl onion, potato onion, bulb serpent garlic, bulb shallot, cultivars, varieties, and/or hybrids of bulb onion, subgroup 3-07A.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Thrips</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 13.0-16.0</p>	<p>Leafminers: Apply when adult leafminers first appear, and repeat as needed to maintain control within constraints of a sound resistance management program.</p> <p>Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program.</p> <p>Do not use this product as a rescue treatment for thrips control.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p>	<p>Do not apply in less than 20 gal. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p> <p>Make 2 consecutive applications of this product then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional applications of this product.</p> <p>Pre-Harvest Interval – 30 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any Abamectin-containing product per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply by air in the State of New York.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>
<p>IMPORTANT: Add a non-ionic activator type wetting, spreading and/or penetrating adjuvant approved for use on bulb onion. Do not use binder sticker type adjuvants. Tank mixing with products that have binder sticker type formulations can reduce performance of this product.</p>			

ONION, GREEN (CROP SUBGROUP 3-07B)

Crops in this group are chive, fresh leaves; chive, Chinese, fresh leaves; *Elegans hosta*; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p><i>Liriomyza</i> leafminers Thrips</p>	<p>Low to moderate pest infestations: 8.0-12.0</p> <p>Heavy pest infestations: 13.0-16.0</p>	<p>Leafminers: Apply when adult leafminers first appear, and repeat as needed to maintain control within constraints of a sound resistance management program.</p> <p>Thrips: Apply in a thrips management program and when thrips are at economic threshold. Repeat application, as needed, to maintain control within constraints of a sound resistance management program.</p> <p>Do not use this product as a rescue treatment for thrips control.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p>	<p>Do not apply in less than 20 gal. of water/A with ground application equipment. Do not apply in less than 5 gal. of water/A with aircraft.</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures), use a greater volume of water to ensure adequate coverage.</p> <p>Pre-Harvest Interval – 7 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 64 fl. oz./A (0.076 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply by air in the State of New York or California.</p> <p>Application Interval: Wait 7 days before repeating application. Make 2 consecutive applications then rotate to a chemistry with a different mode of action. Make at least 2 applications of a chemistry with a different mode of action before making additional applications of this product.</p>
<p>IMPORTANT: Add a non-ionic activator type wetting, spreading and/or penetrating adjuvant approved for use on bulb onion. Do not use binder sticker type adjuvants. Tank mixing with products that have binder sticker type formulations can reduce performance of this product.</p>			

PAPAYA

Crops in this group include papaya, star apple, black sapote, sapodilla, canistel, and mamey sapote

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
Twospotted spider mite	<p align="center">Low to moderate pest infestations: 10.0-15.0</p> <p align="center">Heavy pest infestations: 16.0-20.0</p> <p align="center">Dilute Sprays: 1.0 – 2.0 fl. oz./100 gal)</p>	<p>Apply using conventional dilute or concentrate ground equipment.</p> <p>ADD a horticultural oil for best results</p> <p>Apply when mites first appear during spring, summer, and/or fall.</p>	<p>Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. When using an electro-static sprayer, less than 50 gal water/A may be used; however, do not use less than 5 gal water/A.</p> <p>Pre-Harvest Interval – 3 days</p> <p>DO NOT apply to mango.</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT make more than 3 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 60 fl. oz./A (0.070 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>Application Interval: wait at least 14 days before repeating application.</p> <p>Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.20%, v/v)</p>
<p>IMPORTANT: Concentrate Spray rates (this product/A) are adjusted to apply the amount/A equal to that used in full cover dilute spray. Rates of this product/100 gal are based on 1,000 gal/A of dilute spray. Ground application: For best results, use 150-300 gal/A of spray mix with a minimum of 3 gal/A of horticultural oil (not a dormant oil) and apply at a ground speed of 1 to 1.5 mph.</p>			

PINEAPPLE

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
Twospotted spider mite	<p>Low to moderate pest infestations: 10.0-15.0</p> <p>Heavy pest infestations: 16.0-20.0</p>	Apply using conventional dilute or concentrate ground equipment.	<p>Ground application only.</p> <p>Use sufficient volumes (greater than 50 gal of water/A) with conventional ground application equipment. When using an electro-static sprayer, less than 50 gal of water/A may be used; however, do not use less than 5 gal of water/A.</p> <p>Pre-Harvest Interval – 16 weeks</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply more than 40 fl. oz./A (0.046 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>Application Interval: wait at least 7 days before repeating application.</p>

POME FRUIT (CROP GROUP 11-10)

Crops in this group include apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
European red mite McDaniel spider mite Pear rust mite Pear psylla Tentiform leafminers Twospotted spider mite White apple leafhoppers Yellow mite	Concentrate Sprays: 10-20 fl. oz./A (Dilute Sprays: 2.5-5.0 fl. oz./100 gal) Add a horticultural oil for best results	Gallons of spray will vary with size and number of trees per acre and density of foliage. For best results, apply in combination with horticultural spray oil (not a dormant oil) as directed. Applications without horticultural spray oil may result in reduced efficacy and/or residual control. Applying this product plus horticultural spray oil fewer than 14 days before or after applying Captan® can result in phytotoxicity and crop loss. Make a second application, if needed, but wait at least 21 days after the first application. SEE RESTRICTIONS. Spider Mites: Apply when spider mite or insect thresholds are reached.	Apply using conventional dilute or concentrate ground equipment. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. Application is not permitted from onset of flowering until after petal fall is complete. Pre-Harvest Interval – 28 days DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application. DO NOT apply more than 40 fl. oz./A (0.047 lb ai/A) of this product or any other Abamectin-containing product per year. DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year. DO NOT apply by air. DO NOT graze livestock in treated orchard. Horticultural Spray Oil (DO NOT use a dormant oil): For dilute sprays use 0.25% or 1 gal./A. For concentrate sprays use a minimum of 1 gal./A. Application Interval: Wait at least 21 days before repeating application.
<p>IMPORTANT: Concentrate Spray rates (this product/A) are determined by selecting the amount needed for a full cover dilute spray. The amount of this product/A as a concentrate spray is the same amount as required for dilute sprays to the orchard/grove and could be less than 10 fl oz. this product/A for small trees. Rates of this product/100 gal are based on 400 gal/A of dilute spray.</p>			

STONE FRUIT CROP GROUP (CROP GROUP 12-12)

Crops in this group include apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry (sweet and tart); Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum (chickasaw, damson, and Japanese); plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
<p>European red mite Pacific spider mite Twospotted spider mite</p>	<p>Concentrate Sprays: 10-20 fl. oz./A (Dilute Sprays: 2.5- 5.0 fl. oz./100 gal)</p> <p style="text-align: center;">PLUS</p> <p>a horticultural oil or nonionic surfactant</p>	<p>Make applications when mites first are seen.</p> <p>Make a second application, if needed, but wait at least 21 days after the first application. SEE RESTRICTIONS.</p> <p>Always apply this product in combination with either a nonionic surfactant, or with horticultural spray oil (not a dormant oil), but use caution since oil may cause phytotoxicity to foliage and fruit.</p>	<p>Apply using conventional dilute or concentrate ground equipment.</p> <p>Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 21 days</p> <p>DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 40 fl. oz./A (0.047 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year.</p> <p>DO NOT make applications from onset of flowering until after petal fall is complete.</p> <p>DO NOT apply by air.</p> <p>DO NOT graze livestock in treated orchards.</p> <p>Horticulture Spray Oil (NOT a dormant oil) 1 gal/A (minimum) (0.25%, v/v)</p> <p>Nonionic Surfactant – labeled rate</p>
<p>IMPORTANT: Concentrate Spray rates (this product/A) are determined by selecting the amount needed for a full cover dilute spray. The amount of this product/A as a concentrate spray is the same amount as required for dilute sprays to the orchard and could be less than 10 fl oz. of this product/A for small trees. Rates of this product/100 gal are based on 400 gal/A of dilute spray.</p> <ol style="list-style-type: none"> 1) A nonionic surfactant that spreads on and/or penetrates the leaf cuticle can improve insect control. It is impossible to test on all stone fruit varieties under the variety of conditions that may cause crop injury. Therefore, when using this product in combination with a non-ionic surfactant, carefully follow the Directions for Use and Precautions on the surfactant label and in official spray guides. 2) A horticultural oil (not a dormant oil) may improve efficacy and may be used but it may increase the possibility of phytotoxicity to foliage and fruit. 			

STRAWBERRIES, LOW GROWING BERRY (CROP SUBGROUP 13-07G)

Crops in this group include bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lignonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Strawberry spider mite Twospotted spider mite</p>	<p align="center">16.0</p>	<p>Optimum coverage of the leaves (tops and bottoms) can be achieved by adjusting the spray volumes and placement of the nozzles.</p> <p>Apply when mites first appear.</p> <p>Make a second application at 7-10 day intervals. Repeat the two step application process, if needed. SEE RESTRICTIONS.</p> <p>A non-ionic or silicone-based (not a sticker or binder type) surfactant used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray.</p>	<p>Ground application only.</p> <p>Apply using conventional ground equipment or electrostatic sprayers.</p> <p>Use sufficient volumes (greater than 50 gal of water/A for conventional sprayers, greater than 10 gal of water/A for electrostatic sprayers) to ensure thorough foliage coverage.</p> <p>Pre-Harvest Interval – 3 days</p> <p>To avoid insect resistance to this product, do not apply to strawberries grown in nurseries.</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 64 fl. oz./A (0.075 lb ai/A) of this product or any other Abamectin-containing product per year.</p> <p>DO NOT apply by air.</p> <p>Application Interval: Wait at least 21 days after the second application before repeating application.</p>

TREE NUTS CROP GROUP (CROP GROUP 14-12)

Crops in this group include African nut-tree; almond; beech nut; Brazil nut; Brazilian pine; bunya; bur oak; butternut, Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnuts (black and English); yellowhorn; cultivars, varieties, and/or hybrids of these.

Target Pests	Application Rate fl. oz./A (fl. oz./100 gal)	Application Instructions and Timing	Restrictions
European red mite Pacific spider mite Strawberry spider mite Twospotted spider mite	Concentrate Sprays: 10-20 fl. oz./A (Dilute Sprays: 2.5- 5.0 fl. oz./100 gal) <p align="center">Plus</p> a horticultural spray oil	Apply this product when spider mites first appear. Greater residual spider mite control is achieved when spray is deposited on new foliage growth. Make a second application, if needed, but wait at least 21 days after the first application. SEE RESTRICTIONS. To manage resistance, if pest infestation requires additional treatments, use other alternative insecticides with different modes of action which are labeled for use on these pests and crops.	Apply using conventional dilute or concentrate ground equipment. Use sufficient volumes (greater than 40 gal of water/A) to ensure thorough foliage coverage. Pre-Harvest Interval – 21 days DO NOT apply more than 20 fl. oz./A (0.023 lb ai/A) of this product or any other Abamectin-containing product per application. DO NOT apply more than 40 fl. oz./A (0.047 lb ai/A) of this product or any other Abamectin-containing product per year. DO NOT make more than 2 applications of this product or any other Abamectin-containing product per year. DO NOT apply by air. DO NOT graze livestock in treated groves/orchards. Horticultural Spray Oil (DO NOT use a dormant oil) 1 gal./A (minimum) (0.25%, v/v)

IMPORTANT: Rates of this product/100 gal are based on 400 gal/A of dilute spray. Concentrate Spray rates (this product/A) are determined by selecting the amount needed for a full cover dilute spray. Use the same amount of product per acre in concentrate sprays that would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl oz. of this product/A on small trees.

TUBEROUS AND CORM VEGETABLES CROP SUBGROUP (CROP SUBGROUP 1 C)

Crops in this group include arracacha, arrowroot, artichoke, Chinese artichoke, edible canna, Jerusalem artichoke, cassava (bitter and sweet), chayote (root), chufa, dasheen, ginger, leren, potato, sweet potato, tanager, turmeric, yam bean, and yam (true)

Target Pests	Application Rate fl. oz./A	Application Instructions and Timing	Restrictions
<p>Colorado potato beetle <i>Liriomyza</i> leafminers Potato psyllid Spider mite</p>	<p>Low to moderate pest infestations: 8.0-12.0 Heavy pest infestations: 16.0</p>	<p>Apply when mites and adult leafminer flies first appear.</p> <p>Colorado potato beetle: Begin applications when 50% of the egg masses have hatched or larvae are observed. If a second application is made, ensure it is to control a single Colorado potato beetle generation.</p> <p><i>Liriomyza</i> leafminers: Make the first application when adult flies are first observed. Repeat applications as needed to maintain control.</p> <p>Spider mites: Make the first application when mites first appear. Repeat application as needed to maintain control.</p> <p>Make a second application, if needed, but wait at least 7 days after the first application. SEE RESTRICTIONS.</p> <p>A non-ionic or silicone-based (not a sticker or binder type) surfactant used in combination with this product provides improved wetting of foliage and allows an even deposition of the spray.</p>	<p>Apply using conventional ground or aerial application equipment.</p> <p>Use sufficient volumes (greater than 20 gal of water/A for ground applications and greater than 5 gal of water/A for aerial applications).</p> <p>Under conditions such as high pest populations, dense foliage, or adverse application conditions (such as high temperatures) use a greater volume of water to ensure adequate coverage.</p> <p>Pre-Harvest Interval – 14 days</p> <p>DO NOT apply more than 16 fl. oz./A (0.019 lb ai/A) of this product or any other Abamectin-containing product per application.</p> <p>DO NOT apply more than 32 fl. oz./A (0.038 lb ai/A) of this product or any other Abamectin-containing product per year for control spider mites or Colorado potato beetle.</p> <p>DO NOT apply more than 48 fl. oz./A (0.056 lb ai/A) of this product or any other Abamectin-containing product per year for control of leafminers.</p> <p>DO NOT make more than 2 applications to control Colorado potato beetle per year.</p> <p>DO NOT make more than 2 sequential applications of this product or any other Abamectin-containing product.</p> <p>DO NOT apply by air in the State of New York.</p> <p>DO NOT graze or feed treated foliage to livestock.</p> <p>Application Interval: Wait at least 7 days before repeating application.</p>

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place.

PESTICIDE DISPOSAL: Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

SPILLS: For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-424-9300, day or night.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV[yymmdd])