

Dusting Sulfur



ORTHO

(FUNGICIDE-MITICIDE)

Active Ingredient	By Wt.
Sulfur	98%
Inert Ingredients	2%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

DIRECTIONS

ALFALFA (Seed): Spider Mites—25 to 50 lbs. per acre. Do not apply during blooming period in desert areas.

ALMONDS, CURRANTS, FIGS, HOPS, PLUMS, PRUNES, WALNUTS: European Red Mite, Pacific Mite—50 lbs. per acre. Dust when Spiders begin to multiply in large numbers. Since the effectiveness of sulfur for these Red Spiders varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.

BEANS, PEAS: Common Red Spiders, Pacific Mites—15 to 20 lbs. per acre. Dust when Spiders begin to multiply in large numbers. Since the effectiveness of sulfur for these Red Spiders varies in different localities, State Agricultural Experiment Stations should be consulted as to the effectiveness before application.

ASPARAGUS: Rust—15 to 20 lbs. per acre. Dust after cutting stops. Irrigate, cultivate, then apply the sulfur. Repeat if disease is severe.

AVOCADOS: Mites—25 to 50 lbs. per acre. Dust when mites are numerous.

BEANS, DAHLIAS, PEAS: Powdery Mildew—15 to 20 lbs. per acre. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

CURRANTS, PEACHES, ROSES: Powdery Mildew—50 lbs. per acre on Currants and Peaches, 30 to 40 lbs. per acre on Roses. Dust when disease appears and re-treat as necessary. Check for plant safety before treating commercially.

CHERRIES: Leaf Spot and/or Brown Rot—40 to 45 lbs. per acre. Make first application at pink bud stage, repeat at petal fall and shuck fall stages. Repeat as necessary through growing season.

CITRUS: Citrus Red Spider, Two-Spotted Mite, Thrips—80 to 100 lbs. per acre. Dust when an infestation threatens.

COTTON: Atlantic Mite, Pacific Mite, Two-Spotted Mite—25 to 35 lbs. per acre.

GRAPES: Powdery Mildew—10 lbs. per acre. Dust first when shoots are 6 inches long. Dust again just before the blossoms open. In severe cases, later dusting may be necessary.

MELONS: Powdery Mildew—10 to 20 lbs. per acre. May be controlled by dusting, but the sulfur often injures the vines, particularly if the temperature reaches 95° F. Treat when disease first appears and re-treat as necessary. Do not use on sulfur-sensitive varieties.

PEACHES, PLUMS, PRUNES: Brown Rot—50 lbs. per acre. Put on a preventive dusting when the fruit starts to ripen. This often reduces the severity of the disease. **Rust**—Apply 50 lbs. per acre during summer when disease appears. Repeat after 2 weeks if necessary.

ACCEPTED

OCT 4 1974

239-67

UNDE... IDEL... TICIDE

FUNGICIDE... MITICIDE

SUGAR BEETS: Spider Mites

50 lbs. per acre.

TOMATOES: Russet Mite, Two Spotted Spider Mite

50 lbs. per acre.

Because of sulfur, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

CAUTION: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90°F. or above). Do not make sulfur applications at such times. Do not use sulfur on Eastern American Type Grapes, Cucumbers, Apricots or other sulfur susceptible plants. Never use sulfur before, with or immediately following foliage oil sprays.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5992-D Product 353-2 Made in U.S.A.

EPA Reg. No. 239-67-AA



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN