



# ORTHO

**DANGER - POISON - PRECAUTIONS**



**DO NOT TOUCH**

**POISONOUS BY SKIN CONTACT**

POISONOUS if touched by hands or applied or splashed on skin, in eyes or in or on clothing, shoes, or on clothes.



**DO NOT BREATHE**

**POISONOUS IF BREATHED**

POISONOUS if inhaled as a gas, mist, or dust.

**POISONOUS TO FISH AND WILDLIFE**

POISONOUS TO FISH AND WILDLIFE. Do not use in or near water. Do not use in or near fish ponds, streams, rivers, or lakes. Do not use in or near wildlife habitats. Do not use in or near areas where fish and wildlife are present.

POISONOUS TO HUMANS. Do not use in or near homes, schools, or day care centers. Do not use in or near areas where children are present.

# Parathion 15 Wetttable

**STOP • READ LABEL BEFORE USING!**

**POISON**

See Side Panels for Antidote & Precautions



**DANGER**

Keep Out of Reach of Children

**NOT FOR HOME USE**

USE ONLY EQUIPMENT (1) Wear work clothes (2) Wash hands after use. Approved by the U.S. Bureau of Entomology and Plant Quarantine. WORK SAFETY AND HEALTH ADMINISTRATION. Do not use in or near water. Do not use in or near fish ponds, streams, rivers, or lakes. Do not use in or near wildlife habitats. Do not use in or near areas where fish and wildlife are present. Do not use in or near homes, schools, or day care centers. Do not use in or near areas where children are present.



**DO NOT USE!**

**DANGER**

Keep Out of  
Reach of  
Children

#### USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING

(1) Wear waterproof pants, coat, hat, rubber boots or rubber overboots. (2) Wear safety goggles. (3) Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11. (4) Wear heavy duty, natural rubber gloves.

#### WORK SAFETY RULES

Keep a first-aid kit and a copy of this label on site. Do not eat, drink, or smoke while using this product.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

**In case of contact** ...

#### NOTE TO PHYSICIANS

Emergency Information: (411) (36) (37) ...

#### POST TREATED AREA

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

#### CONTAINER DISPOSAL

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

#### CROP USE DIRECTIONS

Do not use eye protection if the applicator is not spraying. Do not use eye protection if the applicator is not spraying.

**Chevron Chemical Company**



# ORTHO

## Parathion 15 Wettable (INSECTICIDE — Contains 15% Parathion)

**DIRECTIONS** This product is not to be used around homes or  
house gardens.

USE INDICATED AMOUNT IN 100 GALS. WATER — UNLESS  
OTHERWISE SPECIFIED.

**APPLES** Codling Moth; Plum Curculio; Orange Tortrix; San  
Jose, Forbes and Scurry Scales, Red-Banded Leaf Roller  
(Second and Third Brood); Woolly Apple Aphid, Mealybug —  
1½ lbs. Bud Moth, Red Bug, Fruit Tree Leaf Roller, Rosy  
Apple Aphid, Green Apple Aphid, Red-Banded Leaf Roller  
(First Brood), Grasshoppers, European Red Mite, Two-  
Spotted Mite, Pacific Mite, Schoenii Mite, Clover Mite  
— 1½ lbs. (Certain Mites, such as Two Spotted, Repeat at 7 to 10  
day intervals during summer months). Apply at petal fall and every  
10 to 14 days thereafter until control is achieved. Directions for  
Mites are applicable to other deciduous fruit, such as Peach,  
Apple, Pear, etc., as well as to fruit and Europe of McIntosh  
Apples, and related varieties. Do not apply within 14 days of  
harvest.

**BEANS** Mexican Bean Beetle, Leafhopper, Aphids — 1 lb.,  
Red Spider Mites, Two Spotted Spider Mite, Leafminer — 1½  
lbs. Armyworms, Leaf Roller — 1 lb. (250 gals. per acre). Repeat  
at 7 to 10 day intervals. Do not apply within 15 days of harvest.

**BEEF** Webworm — 1½ lbs. (250 gals. per acre). Aphids — 1½  
lbs. (250 gals. per acre). Do not apply within 15 days of harvest.  
(Do not apply to alfalfa hay used for hay or feed.)

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, KALE, MUS-  
TARD GREENS, TURNIPS** Diamond back Moth, Imported  
Cabbageworm, Armyworms — 1½ lbs. (250 gals. per acre).

**Aphids, Thrips** — 1 lb. (250 gals. per acre). Do not apply to  
Broccoli and Brussels Sprouts within 7 days of harvest and to  
Cabbage, Kale, Mustard Greens and Turnips within 10 days of  
harvest.

**CARROT** Aphids — 3½ lbs. (300 gals. per acre). Do not apply  
within 15 days of harvest. Do not use treated carrot tops for  
food or feed.

**CELERY** Celeryworms, Aphids — 1½ lbs. per acre. Spiders, Mites  
— 2¾ lbs. Do not apply within 15 days of harvest.

**CHERRY** Plum Curculio, Cherry Fruit Worm, Oriental Fruit  
Moth, Bud Moth, Tortrix, Cankerworms — 1½ lbs. Aphids,  
Mites — 1½ lbs. Apply at petal fall and every 11 to 14 days  
thereafter until control is achieved. Do not apply within 14 days  
of harvest.

**CITRUS (Florida)** Purple Scale, Florida Red Scale, Chaff  
Scale, Snow Scale, Mealybugs, White Fly, Aphids — 1½ to 2  
lbs. when needed. Best results are obtained when used in the  
post bloom spray and in summer spray during June through Sep-  
tember. Make thorough coverage. Cottony Cushion Scale — 2  
lbs. Note: Parathion does not control Soft Brown Scale. Do not  
apply within 14 days of harvest. Do not apply rates exceeding 4  
lbs. (actual Parathion) per acre within 30 days of harvest.

**CORN** European Corn Borer — 3 lbs. per acre in water two or  
three times during growing season. Consult state agricultural experiment station for proper  
timing of treatment. Armyworms, Corn Rootworm Adults —  
1½ lbs. per 100 gals. water. Do not apply over 3 lbs. per acre.  
European Corn Borer — apply when adult beetles are im-  
mature. Do not apply over 12 days of harvest in cutting for  
hay.

**CUCUMBER, SQUASH, MELONS** Cucumber Beetle, Melon  
worm, Pickleworm, Serpentine Leaf Miner, Aphids, Thrips,  
Spider Mite — 1½ lbs. (250 gals. per acre). Do not apply to  
Melons, Cucumbers, or Squash until 10 days after harvest. Do not  
apply to Cucumbers, Squash, and Melons until 10 days  
after harvest.

**ONIONS** Thrips — 1 lb. (250 gals. per acre). Repeat at 7 to 10

day intervals, if necessary. Do not apply  
to Onions within 10 days of harvest.

**PEACH** Plum Curculio, Oriental  
Peach Twig Borer, Cat-facing  
Leaf Roller, Cottony Peach Scale,  
Spider Mites — 1 lb. California  
days of harvest. Do not apply more  
than 5 lbs. (actual Parathion) per  
acre. Do not apply more than 5 lbs. (actual  
Parathion) per acre within 10 to 14 days  
thereafter until control is achieved.  
Do not apply within 14 days of harvest.  
Do not apply more than 5 lbs. (actual  
Parathion) per acre per year.

**PECANS** Black Pecan Aphid,  
May Beetle, Pecan Leaf Casella,  
Spittlebug — 2 lbs. Twig Girdler —  
1½ lbs. Do not apply within 15  
days of harvest or after shucks are  
open.

**PEPPERS** Aphids, Serpentine  
Leaf Miner — 1½ lbs. Do not apply  
within 15 days of harvest.

**POTATO, TOMATO** Colorado  
Leafhopper, Serpentine Leaf Miner,  
Aphids — 1½ lbs. (250 gals. per  
acre). Do not apply to Potatoes,  
Tomatoes within 10 days of harvest.

**SPINACH** Aphids — 1 lb. (250  
gals. per acre). Do not apply within  
14 days of harvest.

**STRAWBERRY** Leaf Roller, Red  
Strawberry Weevil — 1½ lbs. (250  
gals. per acre). Repeat at 7 to 10  
day intervals. Do not apply within  
14 days of harvest.

**TOBACCO (Field)** Aphids — 1½  
lbs. (250 gals. per acre). Do not apply  
within 5 days of harvest. Do not  
apply to tobacco plants within 14  
days of harvest. Do not apply to  
tobacco plants within 14 days of  
harvest.

**CONDITIONS OF SALE** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under liability resulting from handling, storage and use of this material.

**Chevron Chemical Company**



# ORTHO

## Parathion 15 Wettable

(INSECTICIDE)

**POISON**

SEE CONTAINER LABEL FOR ANTIDOTE AND DANGER STATEMENTS

READ ENTIRE LABEL AND BOOKLET. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. COMPLETELY EMPTY CONTAINER. DISPOSE OF WASTE PESTICIDE AND CONTAINER BY BURYING. DO NOT BURN. NEVER RE-USE.

**DIRECTIONS FOR USE**