



BUG-GETA[®] Meal

For Control of SNAILS & SLUGS

**Also Kills: Cutworms,
Brachyrhinus Weevil**

**NET WEIGHT
2½ POUNDS**

Active Ingredients		By Wt.
Calcium Arsenate	12.5%	5.16%
Metaldehyde	3.4%	1.75%
Inert Ingredients	84.1%	93.09%
Arsenic expressed as metallic	1.95%	
Water soluble arsenic As ₂ maximum	1.95%	

BUG-GETA Meal is a composition of specially balanced ingredients, including metaldehyde, Calcium Arsenate and Metaldehyde as the active ingredients.

It is highly effective against a wide variety of insects and foreign pests such as SNAILS, SLUGS, CUTWORMS, and BRACHYRHINUS WEEVILS. BUG-GETA is used on Flower Gardens and Lawn Areas, around Ornamental Flowering Plants and in Lath Houses and Greenhouses.

Do not apply directly to plants.

KEEP AWAY FROM CHILDREN, PETS AND LIVESTOCK.

A METALDEHYDE ARSENICAL BAIT

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP FEEDER IN ORIGINAL CONTAINER



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

WARNING: May be fatal if swallowed. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Avoid contamination of feed and feed-stuffs. Keep out of reach of children and domestic animals. Dispose of container promptly when empty. This product is toxic to wildlife. Birds feeding in treated areas may be killed.

DIRECTIONS

SNAILS, SLUGS, CUTWORMS around Ornamental Flowering Plants—The preferred method of treating with BUG-GETA is to have the area thoroughly watered in the area to be treated. Then broadcast BUG-GETA Meal over this area, or place it in small mounds of about 1 heaping teaspoonful every 2 feet. In case weather the mounds of BUG-GETA should be somewhat larger, about 1 pound to each 300 square feet. Use broadcast method if children, dogs, cats or birds are apt to disturb the bait.

Heavy watering or rains will cause some of the BUG-GETA to drift into the ground and become ineffective. By having the ground watered thoroughly before applying the bait, the next watering can be postponed for a longer time, thus allowing the bait to stay on the surface longer.

Make first application at beginning of spring planting season before plants reach maturity. Repeat at 2 week intervals, after heavy rains, or as needed.

CAMELIAS, AZALEAS, RHODODENDRONS AND OTHER PLANTS: BRACHYRHINUS WEEVILS (Strawberry Root Weevils) Place 2 to 4 teaspoonfuls around base of each plant depending on size of plant.

GREENHOUSES AND LATH HOUSES: Distribute meal on and beneath benches but do not place on pets or fruits unless plants are well established. Keep meal from coming in contact with plants. Insulate well before distributing meal.

Note: This product is sold by weight and package is full when packed, not due to fluffy nature, the material is included to weight.

NOTICE: Buyer assumes all responsibility for safety or loss; not in accordance with directions.

CHEVRON CHEMICAL COMPANY

FORM 6963-E

EPA Reg. No. 239-111

ORTHO DIVISION—San Francisco, California 94119

Product 175

RICHMOND, CALIF.

LAFAYETTE, CALIF.

DES PLAINES, ILL.



PERTH AMBOY, N.J.

ATLANTA, G.A.