

[239-113]
**Rotenone 1
 Dust**

ACCEPTED
 AUG 23 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 239-113 SUBJECT
 TO SECTION 200000



(INSECTICIDE)

Active Ingredients	By Wt.
Rotenone from Cubé	1%
Other Resins from Cubé	1%
Inert Ingredients	98%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.
 Dust repeatedly as soon as insects appear. Do not apply to fruits and vegetables within 1 day of harvest.

- X ASPARAGUS: Asparagus Beetle—Use 25 to 30 pounds per acre.
- X BEANS: ~~Twelve-Spotted Beetle~~, Thrips, Flea Beetles—Use 15 to 25 pounds per acre.
- X BLUEBERRIES: Blueberry Maggot—Use 30 pounds per acre.
- X CABBAGE: Cabbage Worms, Flea Beetles—Use 25 to 35 pounds per acre.

NET WEIGHT 50 LBS.


Chevron Chemical Company
 Ortho Division/San Francisco, Calif. 94119
 Richmond, California Fresno, California Honolulu, Hawaii
 Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
 Form 2337-R EPA Reg. No. 239-113 Product 1297 Made in U.S.A.

- X CHERRIES: Fruit Fly—Use 40 to 60 pounds per acre.
- X CRANBERRIES: Cranberry Fireworm, Cranberry Fruitworm—Use 35 lbs. per acre.
- CUCUMBERS: Cucumber Beetle, Flea Beetles — Use 20 to 25 lbs. per acre.
- CURRENTS, GOOSEBERRIES: Gooseberry Fruit Worm — Use 30 pounds per acre.
- MELONS: Cucumber Beetles, Flea Beetles—Use 20 to 25 lbs. per acre.
- PEAS: Pea Weevil—Apply 20 to 25 pounds per acre. The dusting should be completed before the pods have set and before the Weevils have had time to move into the fields, thus necessitating the dusting of a wider border. Fields of ten acres or less should be dusted over the entire area with ORTHO Rotenone 1 Dust. In larger fields the field edges should be dusted to a depth of not less than 150 to 200 feet, unless actual infestations are known to be concentrated in a narrower zone.
- POTATOES: Potato Flea Beetle—Use 15 to 20 pounds per acre. Repeat as necessary.
- PRUNES: Pear Thrips—Use 30 to 40 pounds per acre.
- X RASPBERRIES and other brambles: Raspberry Fruit Worm—Use 40 pounds per acre.


- STRAWBERRIES: Spittle Bug—Use 30 pounds per acre.
- SWEET CORN: European Corn Borer—Use 30 pounds per acre. Apply when eggs start to hatch. Repeat treatments at 5 to 7-day intervals as necessary. Additional applications should be made for first and second generation borer when necessary. Consult your State Agricultural Experiment Stations for the number and timing of treatments.
- FLEAS: Against Fleas in houses—Sweep or dust ORTHO Rotenone 1 Dust into the cracks of the floors around baseboards, etc. If animals are kept in the basement, ORTHO Rotenone 1 Dust should be liberally sprinkled around the floor. Repeat the treatment as often as necessary and most frequently where the animals are permitted to run at large.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above), resulting from handling, storage, and use of this material.


READ THE LABEL
 KEEP OUT OF REACH OF CHILDREN
CAUTION: Harmful if swallowed.

[239-113]
**Rotenone 1
 Dust**



(INSECTICIDE)

Active Ingredients	By Wt
Rotenone from Cuba	1%
Other Resins from Cuba	1%
Inert Ingredients	98%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

TURN OFF WATER WHEN EMULSIONING. DO NOT SMOKE.

DIRECTIONS

Use a sprayer or blower for application by hand or mechanical means.

Do not reapply until insects appear. Do not apply to plants until 10 days after last day of harvest.

ASPARAGUS: Asparagus Beetle—Use 25 to 30 lbs. per acre.

BEANS: Twelve-Spotted Beetle, Thrips, Flea Beetles—Use 10 to 20 lbs. per acre.

BLUEBERRIES: Blueberry Maggot—Use 30 pounds per acre.

CABBAGE: Cabbage Worms, Flea Beetles—Use 25 to 30 lbs. per acre.

CHERRIES: Fruit Fly—Use 40 to 60

CRANBERRIES: Cranberry Fireworm—Use 35 lbs. per acre.

CUCUMBERS: Cucumber Beetle, Flea Beetles—Use 25 lbs. per acre.

CURRENTS, GOOSEBERRIES: Gooseberry Flea Beetle—Use 30 pounds per acre.

MELONS: Cucumber Beetles, Flea Beetles—Use 25 lbs. per acre.

PEAS: Pea Weevil—Apply 20 to 25 lbs. per acre. Dusting should be completed before the Weevils have had time to begin nesting. The dusting of a treacher or bed should be dusted with ORTHO Rotenone 1 Dust. In larger fields, dust to a depth of no less than 2 inches and to a depth of no less than 2 inches.

POTATOES: Potato Flea Beetle—Use 25 to 30 lbs. per acre as necessary.

PRUNES: Pear Thrips—Use 20 to 30 lbs. per acre.

RASPBERRIES and other brambles: Use 25 to 30 lbs. per acre.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Orto Division, San Francisco, Calif. 94119

Richmond, California

Fresno, California

Honolulu, Hawaii

Des Moines, Iowa

Edith, New Jersey

Orlando, Florida

Form 100-2, EPA Reg. No. 239-113, Product 1274, Made in U.S.A.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: This product is highly flammable. Do not use near open flames or other sources of ignition.